

4-manifolds with inequivalent symplectic forms and 3-manifolds with inequivalent fibrations

Curtis T. McMullen and Clifford H. Taubes*

September 28, 2003

Abstract

We exhibit a closed, simply connected 4-manifold X carrying two symplectic structures whose first Chern classes in $H^2(X, \mathbb{Z})$ lie in disjoint orbits of the diffeomorphism group of X . Consequently, the moduli space of symplectic forms on X is disconnected.

The example X is in turn based on a 3-manifold M . The symplectic structures on X come from a pair of fibrations $\pi_0, \pi_1 : M \rightarrow S^1$ whose Euler classes lie in disjoint orbits for the action of $\text{Diff}(M)$ on $H_1(M, \mathbb{R})$.

1 Introduction

Symplectic 4-manifolds. A *symplectic form* ω on a smooth manifold X^{2n} is a closed 2-form such that $\omega^n \neq 0$ pointwise. Given a pair of symplectic forms ω_0 and ω_1 on X , we say:

- (i) ω_0 and ω_1 are *homotopic* if there is a smooth family of symplectic forms ω_t , $t \in [0, 1]$, interpolating between them;
- (ii) ω_0 is a *pullback* of ω_1 if $\omega_0 = f^*\omega_1$ for some diffeomorphism $f : X \rightarrow X$; and
- (iii) ω_0 and ω_1 are *equivalent* if they are related by a combination of (i) and (ii).

*This research was supported in part by the NSF. 1991 Mathematics Subject Classification: Primary 57R57, Secondary 57M25, 57N13.

Any symplectic form ω admits a compatible almost complex structure $J : TX \rightarrow TX$ (satisfying $\omega(v, Jv) > 0$ for $v \neq 0$). Let $c_1(\omega) \in H^2(X, \mathbb{Z})$ denote the first Chern class of the (canonical) complex line bundle $\wedge_{\mathbb{C}}^n TX$ determined by J . It is easy to see that the first Chern class is a deformation invariant of the symplectic structure; that is, $c_1(\omega_0) = c_1(\omega_1)$ if ω_0 and ω_1 are homotopic.

The purpose of this note is to show:

Theorem 1.1 *There exists a closed, simply-connected 4-manifold X which carries a pair of inequivalent symplectic forms. In fact, ω_0 and ω_1 can be chosen such that $c_1(\omega_0)$ and $c_1(\omega_1)$ lie in disjoint orbits for the action of $\text{Diff}(X)$ on $H^2(X, \mathbb{Z})$.*

One can also formulate this result by saying that the moduli space $\mathcal{M} = (\text{symplectic forms on } X) / \text{Diff}(X)$ is disconnected.

Fibered 3-manifolds. To construct the 4-dimensional example X , we first produce a compact 3-dimensional manifold M^3 that fibers over the circle in two unrelated ways.

To describe this example, we recall the correspondence between closed 1-forms and measured foliations. Let α be a closed 1-form on M , such that α and its pullback to ∂M are pointwise nonzero. Then α defines a *measured foliation* \mathcal{F} of M^3 , transverse to ∂M , with $T\mathcal{F} = \text{Ker } \alpha$ and with transverse measure $\mu(T) = \int_T |\alpha|$. Conversely, a (transversally oriented) measured foliation \mathcal{F} determines such a 1-form α . If α happens to have integral periods, then we can write $\alpha = d\pi$ for a *fibration* $\pi : M \rightarrow S^1 = \mathbb{R}/\mathbb{Z}$, and the leaves of \mathcal{F} are then simply the fibers of π .

The *Euler class* of a measured foliation,

$$e(\mathcal{F}) = e(\alpha) \in H_1(M, \mathbb{Z}) / (\text{torsion}),$$

is represented geometrically by the zero set of a section $s : M \rightarrow T\mathcal{F}$, such that the vector field $s|_{\partial M}$ is inward pointing and nowhere vanishing.

Just as for symplectic forms, we say:

- (i) α_0 and α_1 are *homotopic* if they are connected by a smooth family of closed 1-forms α_t , nonvanishing on M and ∂M ;
- (ii) α_0 is a *pullback* of α_1 if $\alpha_0 = f^*\alpha_1$ for some $f \in \text{Diff}(M)$; and
- (iii) α_0 and α_1 are *equivalent* if they are related by a combination of (i) and (ii).

In the 3-dimensional arena we will show:

Theorem 1.2 *There exists a compact link complement $M = S^3 - \mathcal{N}(K)$ which carries a pair of inequivalent measured foliations α_0 and α_1 . In fact α_0 and α_1 can be chosen to be fibrations, with $e(\alpha_0)$ and $e(\alpha_1)$ in disjoint orbits for the action of $\text{Diff}(M)$ on $H_1(M, \mathbb{Z})$.*

(Here and below, $\mathcal{N}(K)$ denotes an open regular neighborhood of a link K in a 3-manifold.)




Figure 1. An axis added to the Borromean rings.

Description of the manifolds. For the specific examples we will present, the link K is obtained from the Borromean rings $K_1 \cup K_2 \cup K_3$ by adding a fourth component K_4 ; see Figure 1. The fourth component is the *axis* of a rotation of S^3 cyclically permuting $\{K_1, K_2, K_3\}$; it can be regarded as a vertical line in \mathbb{R}^3 , normal to a plane nearly containing the rings.

Alternatively, we can also write $M = T^3 - \mathcal{N}(L)$, where

- $T^3 = \mathbb{R}^3/\mathbb{Z}$ is the flat Euclidean 3-torus,
- $L \subset T^3$ is a union of 4 disjoint, oriented, closed geodesics,
- (L_1, L_2, L_3) gives a basis for $H_1(T^3, \mathbb{Z})$, and
- $L_4 = L_1 + L_2 + L_3$ in $H_1(T^3, \mathbb{Z})$.

The 4-manifold X of Theorem 1.1 is the fiber-sum of $T^3 \times S^1$ with 4 copies of the elliptic surface $E(1) \rightarrow \mathbb{C}\mathbb{P}^1$, with the elliptic fiber $F \subset E(1)$

glued along $L_i \times S^1$. The key to the example is that $\text{Diff}(X)$ preserves the *Seiberg–Witten norm*

$$\|s\|_{\text{SW}} = \sup\{|s \cdot t| : \text{SW}(t) \neq 0\}$$

on $H^2(X, \mathbb{R})$, just as $\text{Diff}(M)$ preserves the Alexander norm on $H^1(M, \mathbb{R})$. The Seiberg–Witten norm manifests the rigidity of the smooth structure on X , allowing us to check that the Chern classes $c_1(\omega_1), c_1(\omega_2)$ lie in different orbits of $\text{Diff}(X)$.

On the other hand, using Freedman’s work one can see that these two Chern classes *are* related by a homeomorphism of X . In fact, using the 3-torus we can write $H^2(X, \mathbb{Z})$ with its intersection form as a direct sum

$$(H^2(X, \mathbb{Z}), \wedge) = (\mathbb{Z}^6, \begin{pmatrix} 0 & I \\ I & 0 \end{pmatrix}) \oplus (V, q),$$

where the Chern classes $c_1(\omega_1), c_1(\omega_2)$ lie in the first factor and are related by an integral automorphism preserving the hyperbolic form. By Freedman’s result [FQ, §10.1], this automorphism of $H^2(X, \mathbb{Z})$ is realized by a homeomorphism of X .

Many more examples can be constructed along similar lines. For a simple variation, one can replace L_4 with a geodesic homologous to $L_1 + L_2 + (2m + 1) \cdot L_3$, $m \in \mathbb{Z}$, and replace the elliptic surface $E(1)$ with its n -fold fiber sum, $E(n)$. The manifolds M and X resulting from these variations also satisfy the Theorems above.

| 3-manifolds | 4-manifolds |
|--|--|
| Measured foliations \mathcal{F} of M | Symplectic forms ω on X |
| Fibrations $M \rightarrow S^1$ | Integral symplectic forms |
| Fibers minimize genus | Pseudo-holomorphic curves minimize genus |
| Euler class $e(T\mathcal{F})$ | First Chern class $c_1(\wedge_{\mathbb{C}}^2 TX)$ |
| Alexander polynomial $\Delta_M \in \mathbb{Z}[H_1]$ | Seiberg–Witten polynomial $\sum \text{SW}(t) \cdot t \in \mathbb{Z}[H^2]$ |
| Alexander norm on $H^1(M, \mathbb{R})$ | Seiberg–Witten norm on $H^2(X, \mathbb{R})$ |

Table 2.

Notes and references. Our examples exploit a dictionary between 3 and 4 dimensions, some of whose entries are summarized in Table 2.

The connection between the Thurston norm and the Seiberg–Witten invariant was developed by Kronheimer and Mrowka in [KM], [Kr2], [Kr1], while the work of Meng–Taubes and Fintushel–Stern brought the Alexander polynomial into play [MeT], [FS1], [FS2], [FS3]. Inasmuch as the Alexander polynomial is tied to the Thurston norm in [Mc2], [Mc1], (see also [Vi]), there is an intriguing circle of ideas here which might be better understood.

2 The Alexander and Thurston norms

In this section we recall the Alexander and Thurston norms for a 3-manifold, and prove that Theorem 1.2 holds for the link complement pictured in the Introduction.

The Thurston norm. Let M be a compact, connected, oriented 3-manifold, whose boundary (if any) is a union of tori. For any compact oriented n -component surface $S = S_1 \sqcup \cdots \sqcup S_n$, let

$$\chi_-(S) = \sum_{\chi(S_i) < 0} |\chi(S_i)|.$$

The *Thurston norm* on $H^1(M, \mathbb{Z})$ measures the minimum complexity of a properly embedded surface $(S, \partial S) \subset (M, \partial M)$ dual to a given cohomology class; it is given by

$$\|\phi\|_T = \inf\{\chi_-(S) : [S] = \phi\}.$$

The Thurston norm extends by linearity to $H^1(M, \mathbb{R})$.

Let $B_T = \{\phi : \|\phi\|_T \leq 1\}$ denote the unit ball in the Thurston norm; it is a finite polyhedron in $H^1(M, \mathbb{R})$. A basic result is:

Theorem 2.1 *Suppose $\phi_0 \in H^1(M, \mathbb{Z})$ is represented by a fibration $M \rightarrow S^1$ with fiber S . Then:*

- $\|\phi_0\|_T = \chi_-(S)$;
- ϕ_0 is contained in the open cone $\mathbb{R}_+ \cdot F$ over a top-dimensional face F of the Thurston norm ball B_T ;
- every cohomology class in $H^1(M, \mathbb{Z}) \cap \mathbb{R}_+ \cdot F$ is represented by a fibration;
- the classes in $H^1(M, \mathbb{R}) \cap \mathbb{R}_+ \cdot F$ are represented by measured foliations; and

- the Euler class $e = e(\phi_0) \in H_1(M, \mathbb{Z})$ is dual to the supporting hyperplane to F . More precisely, $\phi(e) = -1$ for all $\phi \in F$.

In this case we say F is a *fibred face* of the Thurston norm ball. For more details, see [Th2] and [Fr].

The Alexander norm. Next we discuss the Alexander polynomial and its associated norm. Let $G = H_1(M, \mathbb{Z})/(\text{torsion}) \cong \mathbb{Z}^{b_1(M)}$. The *Alexander polynomial* Δ_M is an element of the group ring $\mathbb{Z}[G]$, well-defined up to a unit and canonically determined by $\pi_1(M)$. It can be effectively computed from a presentation for $\pi_1(M)$ (see e.g. [CF]). Writing

$$\Delta_M = \sum_G a_g \cdot g,$$

the *Newton polygon* $N(\Delta_M) \subset H_1(M, \mathbb{R})$ is the convex hull of the set of g such that $a_g \neq 0$. The *Alexander norm* on $H^1(M, \mathbb{R})$ measures the length of the image of the Newton polygon under a cohomology class $\phi : H_1(M, \mathbb{R}) \rightarrow \mathbb{R}$; that is,

$$\|\phi\|_A = |\phi(N(\Delta_M))|.$$

From [Mc2] we have:

Theorem 2.2 *If M is a 3-manifold with $b_1(M) \geq 2$, then we have*

$$\|\phi\|_A \leq \|\phi\|_T$$

for all $\phi \in H^1(M, \mathbb{R})$; and equality holds if ϕ is represented by a fibration $M \rightarrow S^1$.

Links in the 3-torus. We now turn to the Thurston norm for link-complements in the 3-torus. Let $T^3 = \mathbb{R}^3/\mathbb{Z}^3$ denote the flat Euclidean 3-torus. Every nonzero cohomology class $\phi \in H^1(T^3, \mathbb{Z})$ is represented by a fibration (indeed, a group homomorphism) $\Phi : T^3 \rightarrow S^1$.

Consider an n -component link $L \subset T^3$, consisting of disjoint, oriented, closed geodesics $L_1 \cup \cdots \cup L_n$. Define a norm on $H^1(T^3, \mathbb{R})$ by

$$\|\phi\|_L = \sum |\phi(L_i)|, \tag{2.1}$$

where the L_i are considered as elements of $H_1(M, \mathbb{Z})$. Let M be the link complement $T^3 - \mathcal{N}(L)$, equipped with the natural inclusion $M \subset T^3$.

Theorem 2.3 Given $\phi \in H^1(T^3, \mathbb{Z})$, let ψ denote its pullback to $M = T^3 - \mathcal{N}(L)$. Then we have:

$$\|\phi\|_L = \|\psi\|_T = \|\psi\|_A. \quad (2.2)$$

Moreover:

- (a) ψ is represented by a fibration $\Psi : M \rightarrow S^1 \iff$
- (b) $\phi(L_i) \neq 0$ for all $i \iff$
- (c) ϕ belongs to the open cone over a top-dimensional face of the norm ball $B_L = \{\phi : \|\phi\|_L \leq 1\} \subset H^1(T^3, \mathbb{R})$.

Proof. We begin by showing (a-c) are equivalent. If ψ is represented by a fibration $\Psi : M \rightarrow S^1$, then the fibers are transverse to ∂M and thus $\phi(L_i) \neq 0$ for all i . On the other hand, the latter condition insures that the linear fibration $\Phi : T^3 \rightarrow S^1$ associated to ϕ restricts to a fibration of M representing ψ , so we have (a) \iff (b). Finally $\|\phi\|_L$ behaves linearly on $H^1(T^3, \mathbb{R})$ unless one of the terms $\phi_i(L)$ changes sign, and thus the cone on the top dimensional faces is exactly the locus where $\phi(L_i) \neq 0$ for all i , showing (b) \iff (c).

To establish equation (2.2), first suppose ψ is represented by a fibration $\Psi : M \rightarrow S^1$ with fiber S . Since we may take $\Psi = \Phi|_M$, we see S is a union of tori with $\sum |\phi(L_i)|$ punctures, and thus

$$\chi_-(S) = \|\psi\|_T = \sum |\phi(L_i)| = \|\phi\|_L.$$

Equality with the Alexander norm holds by Theorem 2.2.

Thus (2.2) holds on the cone over the top-dimensional faces of B_L . Since this cone is dense, (2.2) holds throughout $H^1(T^3, \mathbb{Z})$ by continuity. \blacksquare

The Borromean rings plus axis. We now turn to the study of the 4-component link $K \subset S^3$ pictured in Figure 1. Let $M = S^3 - \mathcal{N}(K)$, and let m_i denote the meridian linking K_i positively. Then (m_1, m_2, m_3, m_4) forms a basis for $H_1(M, \mathbb{Z}) \cong \mathbb{Z}^4$, and the Alexander polynomial Δ_M can be written as a Laurent polynomial in these variables.

Lemma 2.4 The Alexander polynomial of $M = S^3 - \mathcal{N}(K)$ is given by

$$\begin{aligned} \Delta_M(x, y, z, t) = & -4 + \left(t + \frac{1}{t}\right) - \left(xy + \frac{1}{xy} + yz + \frac{1}{yz} + xz + \frac{1}{xz}\right) \\ & + \left(xyz + \frac{1}{xyz}\right) + \left(x + \frac{1}{x} + y + \frac{1}{y} + z + \frac{1}{z}\right), \end{aligned}$$




Figure 3. The Newton polygon of $\Delta_M(x, y, z, 1)$ (top), and its dual.

where $(x, y, z, t) = (m_1, m_2, m_3, m_4)$.

Proof. The projection in Figure 1 yields the Wirtinger presentation

$$\begin{aligned} \pi_1(M) = \langle a, b, c, d, e, f, g, h, i, j, k, l : \\ aj = jb, bi = ic, gc = ag, dc = ce, ae = fa, fj = jd, \\ ge = eh, hj = ji, di = gd, jg = gk, kc = cl, le = ej \rangle. \end{aligned}$$

Here (a, b, c) , (d, e, f) , (g, h, i) and (j, k, l) are the edges of K_1 , K_2 , K_3 and K_4 respectively. Given this presentation, the calculation of Δ_M is a straightforward application of the Fox calculus [Fox]. \blacksquare

Figure 3 shows the intersection of the Newton polygon $N(\Delta_M)$ with the (x, y, z) -hyperplane.

To bring the 3-torus into play, recall that 0-surgery along the Borromean rings determines a diffeomorphism

$$S^3 - \mathcal{N}(K_1 \cup K_2 \cup K_3) \cong T^3 - \mathcal{N}(L_1 \cup L_2 \cup L_3),$$

where (L_1, L_2, L_3) are disjoint closed geodesics forming a basis for $H_1(T^3, \mathbb{Z})$. Under this surgery, the meridians (m_1, m_2, m_3) go over to longitudes of (L_1, L_2, L_3) . On the other hand, K_4 goes over to the isotopy class of a geodesic $L_4 \subset T^3$, with

$$L_4 = L_1 + L_2 + L_3 \quad \text{in } H_1(T^3, \mathbb{Z}).$$

(To check the homology class of L_4 , note that in S^3 we have $\text{lk}(K_i, K_4) = 1$ for $i = 1, 2, 3$.)

The meridian m_4 goes over to a meridian of L_4 , so unlike (m_1, m_2, m_3) it becomes trivial in $H_1(T^3, \mathbb{Z})$. Thus we have:

$$H^1(M, \mathbb{R}) \supset H^1(T^3, \mathbb{R}) = (\mathbb{R} \cdot m_4)^\perp.$$

Lemma 2.5 *The action of $\text{Diff}(M)$ on $H^1(M, \mathbb{R})$ preserves the subspace $H^1(T^3, \mathbb{R})$.*

Proof. Consider the Newton polygon

$$N = N(\Delta_M) \subset H_1(M, \mathbb{R}),$$

where Δ_M is given by Proposition 2.4. Since $(t + 1/t)$ is the only expression in Δ_M involving t , we have $N = N_0 + [-1, 1] \cdot t$ where

$$N_0 = N(\Delta_M(x, y, z, 1))$$

is the polyhedron in (x, y, z) -space shown in Figure 3. The vertices $\pm t$ of N are thus combinatorially distinguished: they are the endpoints of 14 edges of N (coming from the 14 vertices of N_0), whereas all other vertices of N have degree 5. Since $\text{Diff}(X)$ preserves N , it also stabilizes the special vertices $\{\pm t\}$, and thus $\text{Diff}(X)$ stabilizes $H^1(T^3, \mathbb{R}) = (\mathbb{R} \cdot t)^\perp = (\mathbb{R} \cdot m_4)^\perp$. ■

Proof of Theorem 1.2. For our chosen link $L \subset T^3$, we have

$$\|\phi\|_L = |\phi(m_1)| + |\phi(m_2)| + |\phi(m_3)| + |\phi(m_1 + m_2 + m_3)|.$$

The unit ball $B_L \subset H^1(T^3, \mathbb{R})$ of this norm is shown in Figure 3 (bottom); it is dual to the convex body N_0 .

Note that B_L has both triangular and quadrilateral faces. Pick integral classes $\phi_0, \phi_1 \in H^1(T^3, \mathbb{Z})$ lying inside the cones over faces F_0 and F_1 of different types, and let $\alpha_0, \alpha_1 \in H^1(M, \mathbb{Z})$ denote their pullbacks to M .

By Theorem 2.3, the classes α_0 and α_1 correspond to fibrations $M \rightarrow S^1$. On the other hand, $\text{Diff}(M)$ preserves the subspace $H^1(T^3, \mathbb{R}) \subset H^1(M, \mathbb{R})$ as well as the norm $\|\phi\|_L = \|\alpha\|_T$ on this subspace. Thus $\text{Diff}(M)$ preserves B_L , so it cannot send the face F_0 to F_1 . The supporting hyperplanes for α_0 and α_1 in B_T thus lie in different orbits of $\text{Diff}(M)$. But these supporting hyperplanes are represented by $e(\alpha_0)$ and $e(\alpha_1)$, so their Euler classes are in different orbits as well. ■

The Thurston norm. As was shown in [Mc2], the Alexander and Thurston norms agree for many simple links. The norms agree for the Borromean rings plus axis $K \subset S^3$ as well.

To see this, note that K can be presented as the closure of a 3-strand braid wrapping once around the axis $K_4 \subset K$. A disk spanning K_4 and transverse to $K_1 \cup K_2 \cup K_3$ determines a fibered face F of the Thurston norm ball B_T . As observed by N. Dunfield, one can use the Teichmüller polynomial [Mc1] to show that for any 3-strand braid, the fibered face F coincides with a face of the Alexander norm ball B_A . In the example at hand, all the vertices of B_A are contained in $\pm F$, so we have $B_A \subset B_T$ by convexity. The reverse inclusion comes from the general inequality $\|\phi\|_A \leq \|\phi\|_T$.

Further example: a closed 3-manifold. To conclude, we describe a *closed* 3-manifold N which fibers over the circle in two inequivalent ways.

Let $M = T^3 - \mathcal{N}(L) = S^3 - \mathcal{N}(K)$ be the link complement considered above. Note that the longitudes of K_1, K_2 and K_3 are all homologous to the meridian m_4 of K_4 , since the components of the Borromean rings are

unlinked, while each component links K_4 once. Since T^3 is obtained by 0-surgery on K , all the meridians of L are homologous to m_4 .

Now let $N \rightarrow T^3$ be the 2-fold covering, branched over L , determined by the homomorphism

$$\xi : H_1(M, \mathbb{Z}) \rightarrow \{-1, 1\}$$

satisfying $\xi(m_1) = \xi(m_2) = \xi(m_3) = 1$ and $\xi(m_4) = -1$.

The pullback map $H^1(T^3, \mathbb{R}) \rightarrow H^1(N, \mathbb{R})$ is easily seen to be injective. We claim it is an isomorphism. To see surjectivity, let $N' \subset N$ be the preimage of $M \subset T^3$. Decomposing $H^1(N', \mathbb{R})$ into eigenspaces for the action of the $\mathbb{Z}/2$ deck group for $N' \rightarrow M$, we obtain an isomorphism

$$H^1(N', \mathbb{R}) \cong H^1(M, \mathbb{R}) \oplus H^1(M, \mathbb{R}_\xi),$$

where the last term represents cohomology coefficients twisted by the character ξ of $\pi_1(M)$. Since $\Delta_M(\xi) = \Delta_M(1, 1, 1, -1) = 4 \neq 0$, we have $H^1(M, \mathbb{R}_\xi) = 0$ (cf. [Mc2, §3]). Thus any cohomology class in $H^1(N, \mathbb{R})$ restricts to a $\mathbb{Z}/2$ -invariant class on N' , so it is the pullback of a class on T^3 .

Moreover, every fibration of T^3 transverse to L lifts to a fibration of N , so we find:

Theorem 2.6 *The Thurston norm ball $B_T \subset H^1(N, \mathbb{R})$ agrees with the norm ball $B_L \subset H^1(T^3, \mathbb{R})$, and every face is fibered.*

Picking fibrations in combinatorially inequivalent faces of B_T as before, we have:

Corollary 2.7 *The closed 3-manifold N admits a pair of fibrations α_0, α_1 such that $e(\alpha_0), e(\alpha_1)$ lie in disjoint orbits for the action of $\text{Diff}(N)$ on $H^2(N, \mathbb{Z})$.*

3 Fiber sum and symplectic 4-manifolds

In this section we recall the fiber sum construction, which can be used to canonically associate a 4-manifold $X = X(P, L)$ to a link L in a 3-manifold P . Under this construction, suitable fibrations of P give symplectic forms on $X(P, L)$, and the Alexander polynomial Δ_M of $M = P - \mathcal{N}(L)$ determines Seiberg–Witten invariants of X . It is then straightforward to prove Theorem 1.1 by taking $X = X(T^3, L)$, where $L \subset T^3$ is the 4-component link discussed in previous sections.

Fiber sum. Let $f_i : T^2 \times D^2 \rightarrow X_i$, $i = 1, 2$ be smooth embeddings of the torus cross a disk into a pair of smooth closed 4-manifolds. Let

$$X'_i = X_i - f(T^2 \times \text{int } D^2);$$

it is a smooth manifold whose boundary is marked by $T^2 \times S^1$. The *fiber sum* Z of X_1 and X_2 is the closed smooth manifold obtained by gluing together X'_1 and X'_2 along their boundaries, such that $(x, t) \in \partial X'_1$ is identified with $(x, -t) \in \partial X'_2$. We denote the fiber sum by

$$Z = X_1 \#_{T_1=T_2} X_2,$$

where $T_i = f(T^2 \times \{0\}) \subset X_i$; note that there is an implicit identification between the normal bundles of the tori T_i .

The fiber sum of symplectic manifolds along symplectic tori is also symplectic. More precisely, if ω_i are symplectic forms on X_i with $\omega_i > 0$ on T_i and $\int_{T_1} \omega_1 = \int_{T_2} \omega_2$, then Z carries a natural symplectic form ω with $\omega = \omega_i$ on X'_i .

For more details, see [Go], [MW], [FS1], [FS2], [FS3].

The elliptic surface $E(1)$. A convenient 4-manifold for use in the fiber-sum construction is the *rational elliptic surface* $E(1)$. The complex manifold $E(1)$ is obtained by blowing up the base-locus for a generic pencil of elliptic curves on $\mathbb{C}\mathbb{P}^2$. Thus $E(1)$ is isomorphic to $\mathbb{C}\mathbb{P}^2 \# 9\overline{\mathbb{C}\mathbb{P}^2}$; it is simply-connected and unique up to diffeomorphism. The pencil provides a holomorphic map $E(1) \rightarrow \mathbb{C}\mathbb{P}^1$ with generic fiber F an elliptic curve, and the canonical bundle of $E(1)$ is represented by the divisor $-F$.

The projection $E(1) \rightarrow \mathbb{C}\mathbb{P}^1$ gives a natural trivialization of the normal bundle of the fiber torus F . Since $F \subset E(1)$ is a holomorphic curve in a projective variety, there is a symplectic (Kähler) form on $E(1)$ with $\omega|_F > 0$.

Each of the nine exceptional divisors gives a holomorphic section

$$s : \mathbb{P}^1 \rightarrow E(1).$$

In particular, a meridian for the fiber F is contractible in $E(1) - \mathcal{N}(F)$, since it bounds the image of a disk under s . Since $E(1)$ is simply-connected, any loop in the complement of F is homotopic to a product of conjugates of meridians, so $E(1) - \mathcal{N}(F)$ is also simply-connected.

For a detailed discussion of the topology of elliptic surfaces, see [HKK, §1] or [GS].

From links to 4-manifolds. Now let $L \subset P^3$ be a framed n -component link in a closed, oriented 3-manifold. Such a link determines:

- a 3-dimensional *link complement* $M = P - \mathcal{N}(L)$, and
- a 4-dimensional *fiber-sum* $X = X(P, L) = (P \times S^1) \#_{L \times S^1 = nF} nE(1)$.

To describe the fiber-sum in more detail, note that each component L_i of L determines a torus

$$T_i = L_i \times S^1 \subset P \times S^1,$$

and the framing of L_i provides a trivialization of the normal bundle of T_i . Take n copies of the elliptic surface $E(1)$ with fiber F ; as remarked above, the projection $E(1) \rightarrow \mathbb{C}\mathbb{P}^1$ provides a natural trivialization of the normal bundle of F . Finally, choose an orientation-preserving identification between $L \times S^1$ and nF . The fiber-sum $X(P, L)$ is then defined using these identifications.

It turns out that every orientation-preserving diffeomorphism of F extends to a diffeomorphism of $E(1)$, preserving the normal data; indeed, the monodromy of the fibration $E(1) \rightarrow \mathbb{C}\mathbb{P}^1$ is the full group $SL_2(\mathbb{Z})$. Thus the diffeomorphism type of $X(P, L)$ is the same for any choice of identification between $L \times S^1$ and nF .

Proposition 3.1 *The fiber-sum X is simply-connected if $\pi_1(M)$ is normally generated by $\pi_1(\partial M)$ (e.g. if M is homeomorphic to a link complement in S^3).*

Proof. When the simply-connected manifolds $n(E(1) - \mathcal{N}(F))$ are attached to $M \times S^1$ along $\partial M \times S^1$, they kill $\pi_1(\partial M \times S^1)$ by van Kampen's theorem. Since the latter groups normally generate $\pi_1(M \times S^1)$, the resulting manifold X is simply-connected. ■

Promotion of cycles. The fiber-sum construction furnishes us with an inclusion $M \times S^1 = (P \times S^1)' \subset X$.

Proposition 3.2 *The map*

$$i : H_1(M, \mathbb{R}) \rightarrow H^2(X, \mathbb{R}),$$

sending a 1-cycle $\gamma \subset M$ to the Poincaré dual of $\gamma \times S^1 \subset X$, is injective.

Proof. The map i is a composition of three maps:

$$H_1(M) \rightarrow H_2(M \times S^1) \rightarrow H_2(X) \rightarrow H^2(X).$$

The first arrow is part of the Künneth isomorphism, and the last comes from Poincaré duality, so they are both injective. As for the middle arrow

$$H_2(M \times S^1) \rightarrow H_2(X),$$

we can use the exact sequence of the pair $(X, M \times S^1)$ to identify its kernel with

$$H_3(X, M \times S^1) \cong H_3(nE(1), nF) \cong H^1(nE(1) - nF) = 0.$$

Here we have used excision, Poincaré duality and the simple-connectivity of $E(1) - F$. Thus all three arrows are injective, and so i is injective. ■

Corollary 3.3 *For an n -component link, we have*

$$b_2^+(X(P, L)) \geq b_1(M) \geq n.$$

Here $b_2^+(X)$ denotes the rank of the maximal subspace of $H_2(X, \mathbb{R})$ on which the intersection form is positive-definite.

Proof. Since 1-cycles in general position on M are disjoint, the intersection form on $H^2(X, \mathbb{R})$ restricts to zero on $i(H_1(M, \mathbb{R}))$. But the intersection form is non-degenerate, so it must admit a positive (and negative) subspace of dimension at least $b_1(M) = \dim i(H_1(M, \mathbb{R}))$.

For the second inequality, just note that we have $b_1(M) \geq b_1(\partial M)/2 = n$. Indeed, by Lefschetz duality, the kernel of $H_1(\partial M) \rightarrow H_1(M)$ is Lagrangian, so the image has dimension n . ■

From fibrations to symplectic forms. A central point for us is that suitable fibrations α of P give rise to symplectic structures ω on $X(P, L)$.

Theorem 3.4 *For any fibration $\alpha \in H^1(P, \mathbb{Z})$ transverse to L , there is a symplectic form ω on $X(P, L)$ with*

$$c_1(\omega) = i(e(\alpha|M)).$$

Proof. Let $\alpha = d\pi$ be the closed 1-form representing a fibration $\pi : P \rightarrow S^1$ transverse to L .

Pick a closed 2-form β on M such that β restricts to an area form on each leaf of \mathcal{F} . (One can construct such a form by representing the monodromy

of the fibration by an area-preserving map.) As observed by Thurston, for $\epsilon > 0$ sufficiently small, the closed 2-form

$$\omega_0 = \alpha \wedge dt + \epsilon\beta$$

is a symplectic form on $P \times S^1$, nowhere vanishing on $L \times S^1$ [Th1]. (Here $[dt]$ is the standard 1-form on $S^1 = \mathbb{R}/\mathbb{Z}$, and α and β have been pulled back to the product).

By scaling the Kähler form, we can provide the i th copy of $E(1)$ with a symplectic form ω_i such that $\int_F \omega_i = \int_{L_i \times S^1} \omega$. Then as mentioned above, ω_0 and (ω_i) joined together under fiber-sum to yield a symplectic form ω on X .

Let $K \rightarrow X$ denote the canonical bundle of (X, ω) . We will compute $c_1(K)$ by constructing a section $\sigma : X \rightarrow K$.

Let $M = P - \mathcal{N}(L)$. As an oriented \mathbb{R}^2 -bundle, $K|(M \times S^1)$ is isomorphic to the pullback of $T\mathcal{F}$ from M . Let $s : M \rightarrow T\mathcal{F}$ be a section such that $s|\partial M$ is inward pointing and nowhere vanishing. Then the zero set of s is a 1-cycle γ representing the Euler class $e(\alpha|M) \in H_1(M, \mathbb{R})$. Pulling back s , we obtain a section $\sigma_0 : M \times S^1 \rightarrow K$ with zero set $\gamma \times S^1$.

Now consider the 4-manifold $E(1)' = E(1) - \mathcal{N}(F)$ attached to $M \times S^1$ along $T_i \times S^1$. If we have $\omega_i(F) > 0$, then $K|E(1)'$ is just the pullback of the canonical bundle of $E(1)$. Since $-F$ is a canonical divisor on $E(1)$, there is a nowhere vanishing section $\sigma_i : E(1)' \rightarrow K$, namely the restriction of a meromorphic 2-form on $E(1)$ with divisor $-F$.

We claim σ_0 and σ_i fit together under the gluing identification between $T_i \times S^1$ and $F \times S^1$. To check this, we use the framings to identify $K|T_i \times S^1$ and $K|F \times S^1$ with the trivial bundle over $T^2 \times S^1$. Under this identification,

$$\sigma_0 : T^2 \times S^1 \rightarrow \mathbb{C}^*$$

is homotopic to the projection $T^2 \times S^1 \rightarrow S^1 \subset \mathbb{C}^*$, since the vector field $s|T_i$ runs along the meridians of ∂M . Similarly,

$$\sigma_i : T^2 \times S^1 \rightarrow \mathbb{C}^*$$

is homotopic to $1/\sigma_0$, because of the simple pole along F . Since $T_i \times S^1$ is identified with $F \times S^1$ using the involution $(x, t) \sim (x, -t)$ on $T^2 \times S^1$, the two sections correspond under gluing.

In the case where we have $\omega_i(F) < 0$, both homotopy classes are reversed, so σ_0 and σ_i still agree under gluing. Thus σ_0 and (σ_i) join together to form a global section $\sigma : X \rightarrow K$ with no zeros outside $M \times S^1$. It follows that $c_1(X, \omega)$ is Poincaré dual to $\gamma \times S^1$; equivalently, that $c_1(\omega) = i(\alpha|M)$. ■

The Seiberg–Witten polynomial. A central feature of the fiber-sum $X = X(P, L)$ is that its Seiberg–Witten polynomial is directly computable.

Assume that X is simply-connected and $b_2^+(X) > 1$. Then the Seiberg–Witten invariant of X can be regarded as a map

$$\text{SW} : H^2(X, \mathbb{Z}) \rightarrow \mathbb{Z},$$

well-defined up to a sign and vanishing outside a finite set. This information is conveniently packaged as a Laurent polynomial

$$\mathcal{SW}_X = \sum_t \text{SW}(t) \cdot t \in \mathbb{Z}[H^2(X, \mathbb{Z})].$$

Theorem 3.5 *Suppose M is the complement of an n -component link $L \subset P$, and $\pi_1(\partial M)$ normally generates $\pi_1(M)$. Then $X = X(P, L)$ is simply-connected, we have $b_2^+(X) \geq n$, and*

$$\mathcal{SW}_X = \pm \sum a_t \cdot i(2t),$$

where $\Delta_M = \sum a_t \cdot t$ is the symmetrized Alexander polynomial of M .

Remarks. This Theorem was established by Fintushel and Stern in the special case where (P, L) is obtained by a certain surgery on a link in S^3 [FS2, Thm. 1.9].¹ To obtain the symmetrized Alexander polynomial, one multiplies $\Delta_K(t)$ by a monomial to arrange that its Newton polygon is centered at the origin. The exponents in the symmetrized polynomial may be half-integral.

Proof. To compute \mathcal{SW}_X , we regard X as the union of manifolds $X_0 = M \times S^1$ and $X_i = E(1) - \mathcal{N}(F)$, $i = 1, \dots, n$, glued together along their boundary. For such manifolds one can define a *relative* Seiberg–Witten polynomial $\mathcal{SW}_{X_i} \in \mathbb{Z}[H^2(X_i, \partial X_i; \mathbb{Z})]$, such that

$$\mathcal{SW}_X = \mathcal{SW}_{X_0} \cdot \mathcal{SW}_{X_1} \cdots \mathcal{SW}_{X_n},$$

using the natural map $H^2(X_i, \partial X_i) \rightarrow H^2(X)$ to compute the product. For this gluing formula, developed by Morgan, Mrowka, Szabo and Taubes, see [FS2, Thm. 2.2] and [Ta].

Now for each $X_i = E(1) - \mathcal{N}(F)$, the relative polynomial is simply 1. To see this, just apply the product formula above to the K3 surface

¹Note: contrary to [FS2, p. 371]: the cohomology classes $[T_j]$ in their formula for \mathcal{SW}_X are always linearly independent in $H^2(X, \mathbb{R})$, by Proposition 3.2 above.

$Z = E(1) \#_F E(1)$, which satisfies $\text{SW}_Z = 1$. (This well-known property of K3 surfaces follows, for example, from equations (4.17) and (4.20) in Witten's original paper [Wit].)

Thus we have $\mathcal{SW}_X = \mathcal{SW}_{X_0} = \mathcal{SW}_{M \times S^1}$. Finally the Seiberg-Witten polynomial for $M \times S^1$ is given in terms of Δ_M by the main result of [MeT], yielding the formula for \mathcal{SW}_X above.

To see $\pi_1(X) = \{1\}$ and $b_2^+(X) \geq n$, apply Proposition 3.1 and Corollary 3.3 above. \blacksquare

Proof of Theorem 1.1. Using the Seiberg–Witten invariants to control the action of $\text{Diff}(X)$, it is now easy to give an example of a simply-connected 4-manifold X with inequivalent symplectic forms.

For a concrete example, let $X = X(T^3, L)$ for the 4-component link $L \subset T^3$ studied in the preceding section, and choose any framing of L . As we have seen, the link-complement $M = T^3 - \mathcal{N}(L)$ is homeomorphic to the exterior $S^3 - \mathcal{N}(K)$ of the Borromean rings plus axis. In particular, $\pi_1(M)$ is the normal closure of $\pi_1(\partial M)$, so X is simply-connected and we have $b_2^+(X) \geq 4$.

Let $m_i, i = 1, \dots, 4$ be the basis for $H_1(M, \mathbb{Z})$ coming from the meridians of $K \subset S^3$. Then the classes $t_i = i(m_i)$ form a basis for $i(H_1(M, \mathbb{Z})) \subset H^2(X, \mathbb{Z})$. By Theorem 3.5, we have:

The Seiberg–Witten polynomial of X is given by

$$\mathcal{SW}_X = \Delta_M(t_1^2, t_2^2, t_3^3, t_4^2),$$

where $\Delta_M(x, y, z, t)$ is given by Lemma 2.4.

In particular, the Newton polygons satisfy $N(\mathcal{SW}_X) = 2i(N(\Delta_M))$.

Now identify $H_1(T^3, \mathbb{R})$ with the subspace of $H_1(M, \mathbb{R})$ spanned by (m_1, m_2, m_3) , and let

$$N_0 = N(\Delta_M) \cap H_1(T^3, \mathbb{R}).$$

As we have seen before, any vertex v of N_0 is dual to a fibered face F of the Thurston norm on $H^1(M, \mathbb{R})$; indeed, v is dual to a fibration pulled back from T^3 . All fibrations ϕ in the cone over F have the same Euler class e , which satisfies

$$\|\phi\|_T = 2\phi(v) = -\phi(e);$$

thus $e = -2v$.

By Theorem 3.4, the vertex

$$i(e) = i(-2v) \in 2i(N_0)$$

is the first Chern class of a symplectic structure on X . Since $v \in N_0$ was an arbitrary vertex, we have:

Every vertex of $2i(N_0) \subset N(\mathcal{SW}_X)$ is the first Chern class of a symplectic structure on X .

Now pick a pair combinatorially distinct vertices

$$v_0, v_1 \in 2i(N_0) \subset N(\mathcal{SW}_X).$$

More precisely, referring to Figure 3 (top), we see $2i(N_0)$ has vertices of degrees 3 and 4; choose one of each type. Then v_0 and v_1 have degrees 5 and 6 as vertices of $N(\mathcal{SW}_X)$, since

$$N(\mathcal{SW}_X) = 2i(N_0) + [-2, 2] \cdot t_4$$

is simply the suspension of $2i(N_0)$. As a consequence, no automorphism of $H^2(X, \mathbb{R})$ stabilizing $N(\mathcal{SW}_X)$ can transport v_0 to v_1 .

To complete the proof, choose symplectic forms on X with $c_1(\omega_0) = v_0$ and $c_1(\omega_1) = v_1$. Then the Chern classes of ω_0 and ω_1 lie in distinct orbits for the action of $\text{Diff}(X)$ on $H^2(X, \mathbb{R})$, since diffeomorphisms preserve the Newton polygon of the Seiberg–Witten polynomial. In particular, ω_0 and ω_1 are inequivalent symplectic forms on X . ■

Question. Could it be that $\text{Diff}(X)$ actually preserves the submanifold $M \times S^1 \subset X$ up to isotopy?

Further example: skirting gauge theory. To conclude, we sketch an *elementary* example of a 4-manifold X carrying a pair of inequivalent symplectic forms — but with $\pi_1(X) \neq 1$. By elementary, we mean the proof does not use the Seiberg–Witten invariants; instead, it uses the fundamental group.

To construct the example, simply let $X = N \times S^1$, where N is the closed 3-manifold discussed at the end of §2.

By considering N as a covering of T^3 with a $\mathbb{Z}/2$ -orbifold locus along L , one can show that $\pi_1(N)$ has trivial center. It follows that $\pi_1(S^1)$ is the center of $\pi_1(X)$, and thus the projection

$$\pi_1(X) \rightarrow \pi_1(N)$$

is canonical. In particular, every diffeomorphism of X induces an automorphism of $\pi_1(N)$.

Now let α_0, α_1 be fibrations of N whose Euler classes are in different orbits for the action of $\text{Aut}(\pi_1(N))$ on $H_1(N, \mathbb{Z})$. (These classes exist as before, because the Alexander polynomial is functorially determined by $\pi_1(N)$, and hence preserved by automorphisms.) Then the Euler classes $e(\alpha_0), e(\alpha_1)$ lie in disjoint orbits for the action of $\text{Diff}(X)$ on $H_1(N) = H_1(X)/H_1(S^1)$.

Now as we have seen above, each α_i gives a symplectic form ω_i on X with $c_1(\omega_i)$ dual to $e(\alpha_i) \times S^1$. Since the Euler classes lie in disjoint orbits for the action of $\text{Diff}(X)$, so do these Chern classes. In particular, ω_0 and ω_1 are inequivalent symplectic forms on X . ■

References

- [CF] R. H. Crowell and R. H. Fox. *Introduction to Knot Theory*. Springer-Verlag, 1977.
- [FS1] R. Fintushel and R. Stern. Constructions of smooth 4-manifolds. In *Proceedings of the International Congress of Mathematicians, Vol. II (Berlin, 1998)*, pages 443–452. Doc. Math., 1998.
- [FS2] R. Fintushel and R. Stern. Knots, links and 4-manifolds. *Invent. math.* **134**(1998), 363–400.
- [FS3] R. Fintushel and R. Stern. Symplectic structures in a fixed homology class. *In preparation*.
- [Fox] R. H. Fox. Free differential calculus I, II, III. *Annals of Math.* **57**, **59**, **64**(1953, 54, 56), 547–560, 196–210, 407–419.
- [FQ] M. Freedman and F. Quinn. *Topology of 4-manifolds*. Princeton University Press, 1990.
- [Fr] D. Fried. Fibrations over S^1 with pseudo-Anosov monodromy. In *Travaux de Thurston sur les surfaces*, pages 251–265. Astérisque, volume 66–67, 1979.
- [Go] R. Gompf. A new construction of symplectic manifolds. *Ann. Math.* **142**(1995), 527–595.
- [GS] R. Gompf and A. Stipsicz. *4-Manifolds and Kirby Calculus*. Amer. Math. Soc., 1999.

- [HKK] J. Harer, A. Kas, and R. Kirby. *Handlebody Decompositions of Complex Surfaces*. Mem. Amer. Math. Soc., No. 350, 1986.
- [Kr1] P. Kronheimer. Embedded surfaces and gauge theory in three and four dimensions. In *Surveys in differential geometry, Vol. III (Cambridge, MA, 1996)*, pages 243–298. Int. Press, 1998.
- [Kr2] P. Kronheimer. Minimal genus in $S^1 \times M^3$. *Invent. math.* **135**(1999), 45–61.
- [KM] P. Kronheimer and T. Mrowka. Scalar curvature and the Thurston norm. *Math. Res. Lett.* **4**(1997), 931–937.
- [MW] J. McCarthy and J. Wolfson. Symplectic normal connect sum. *Topology* **33**(1994), 729–764.
- [Mc1] C. McMullen. Polynomial invariants for fibered 3-manifolds and Teichmüller geodesics for foliations. *Ann. scient. Éc. Norm. Sup.* **33**(2000), 519–560.
- [Mc2] C. McMullen. The Alexander polynomial of a 3-manifold and the Thurston norm on cohomology. *Ann. scient. Éc. Norm. Sup.* **35**(2002), 153–172.
- [MeT] G. Meng and C. H. Taubes. \underline{SW} = Milnor torsion. *Math. Res. Lett.* **3**(1996), 661–674.
- [Ta] C. Taubes. The Seiberg-Witten invariants and 4-manifolds with essential tori. *In preparation*.
- [Th1] W. P. Thurston. Some simple examples of symplectic manifolds. *Proc. Amer. Math. Soc.* **55**(1976), 467–468.
- [Th2] W. P. Thurston. A norm for the homology of 3-manifolds. Mem. Amer. Math. Soc., No. 339, pages 99–130, 1986.
- [Vi] S. Vidussi. The Alexander norm is smaller than the Thurston norm: a Seiberg-Witten proof. *École Polytechnique Preprint 99-6*.
- [Wit] E. Witten. Monopoles and four-manifolds. *Math. Res. Letters* **1**(1994), 769–796.

MATHEMATICS DEPARTMENT
HARVARD UNIVERSITY
1 OXFORD ST
CAMBRIDGE, MA 02138-2901