

Some remarks on signs in functional equations

Benedict H. Gross

To Robert Rankin

Let k be a number field, and let M be a pure motive of weight n over k . Assume that there is a non-degenerate pairing $M \times M \rightarrow \mathbf{Q}(-n)$. For example, M could arise from the cohomology in degree n of a projective, smooth, geometrically connected variety X of dimension n over k , and the pairing from the cup product. In this case, the pairing will be symplectic (alternating) if n is odd, and orthogonal (symmetric) if n is even.

Then the conjectural functional equation for the complete L -function of M (including the local factors at infinity) takes the form (cf. [De, (1.2.3)])

$$(0.1) \quad L(M, s) = \epsilon(M, s)L(M, n + 1 - s).$$

Here $\epsilon(M, s) = \epsilon(M) \cdot f(M)^{(n+1)/2-s}$, with $f(M) \geq 1$ an integer, and $\epsilon(M) = \pm 1$ the sign of our title.

In the cases where $L(M, s)$ has an analytic interpretation, and its functional equation is known, the sign $\epsilon(M)$ is of considerable importance. We often get interesting formulae by computing $\epsilon(M)$ in distinct ways.

1. A famous example, illustrating the last point, is the calculation of the sign of the quadratic Gauss sum (cf. [Da, Ch 2]).

Let $k = \mathbf{Q}$, and assume M is orthogonal, of weight 0 and rank 1. Then M corresponds to a quadratic character of the Galois group, or equivalently, to a quadratic étale extension K . Let D be the discriminant of K .

Dirichlet reinterpreted the theorem of quadratic reciprocity to show that the factorization of a prime p in K depended only the class of $p \pmod{D}$. Hence

$$(1.1) \quad L(M, s) = L(\chi, s)$$

where χ is a quadratic character of $(\mathbf{Z}/D\mathbf{Z})^*$, and $L(\chi, s)$ is the (complete) Dirichlet L -function.

Let $f = \chi(-1) \cdot D = |D|$, which is equal to the conductor $f(M)$, and define the quadratic Gauss sum:

$$(1.2) \quad g(\chi) = \sum_{\substack{a=1 \\ (a,f)=1}}^f \chi(a) e^{2\pi i a/f}.$$

Let \sqrt{D} be the unique square-root of D in \mathbf{C} , which is positive when $D > 0$, and has positive imaginary part when $D < 0$. Then Dirichlet's functional equation gives the formula [Da, Ch 9]:

$$(1.3) \quad \epsilon(M) = g(\chi)/\sqrt{D}.$$

In particular, since $\epsilon(M)^2 = 1$, this yields the algebraic identity of Gauss:

$$(1.4) \quad g(\chi)^2 = D = \chi(-1) \cdot f.$$

On the other hand, $L(M, s)$ is the quotient of the (complete) zeta function of K by the (complete) zeta function of \mathbf{Q} . Since Hecke's functional equation for any zeta function has $\text{sign} = 1$, we have

$$(1.5) \quad \epsilon(M) = +1$$

$$(1.6) \quad g(\chi) = \sqrt{D}.$$

2. Another important result involving signs is the proof by Serre of Hecke's theorem on the class of the absolute different (cf. [W, pg. 291]).

The above argument shows that for any orthogonal motive M of rank 1 and weight 0 over k , $\epsilon(M) = +1$. Indeed, M corresponds to a quadratic, étale extension K , and $L(M, s)$ is the quotient of the zeta function of K by the zeta function of k .

Now assume further that K is everywhere unramified over k . By class field theory, M gives rise to a quadratic character χ of the ideal class group $C(k)$ of k , and $L(M, s)$ is equal to the Hecke L -series $L(\chi, s)$.

Hecke showed that the sign in the functional equation of $L(\chi, s)$ is $\chi(\mathcal{D})$, for any ideal class character χ , where \mathcal{D} is the absolute different of k (cf. [T, pg. 94]). Hence, for all *quadratic* χ , we have

$$(2.1) \quad \chi(\mathcal{D}) = +1.$$

This implies that the class $[\mathcal{D}]$ of \mathcal{D} is a *square* in $C(k)$, i.e. $[\mathcal{D}]$ lies in the subgroup $2C(k)$.

3. What can we say about the class of \mathcal{D} in the strict class group $C^+(k)$ —the quotient of the group of ideals by the subgroup of principal ideals with a totally positive generator? The following result was obtained in [G-M]; the proof given here follows ideas of Armitage and Fröhlich [A-F].

Proposition 3.1. *Let K/k be a quadratic extension which is unramified at all finite primes, and let χ be the corresponding quadratic character of $C(k)^+$. Let T be the set of real places of k which become complex in K .*

Then the integer $\#T$ is even, and $\chi(\mathcal{D}) = (-1)^{\#T/2}$. The class $[\mathcal{D}]$ of \mathcal{D} is a square in $C(k)^+$ if and only if for every unramified, quadratic extension K of k , the integer $\#T$ is divisible by 4.

Proof. The sign in the functional equation of $L(\chi, s) = L(M, s)$ is given by ([T, pg. 94]):

$$\epsilon(M) = (-1)^{\#T/2} \cdot \chi(\mathcal{D}).$$

Since $L(M, s)$ is the quotient of two zeta functions, $\epsilon(M) = +1$, and $\chi(\mathcal{D}) = (-1)^{\#T/2}$ as claimed. The class $[\mathcal{D}]$ lies in $2C(k)^+$ if and only if $\chi(\mathcal{D}) = 1$ for all quadratic characters χ of $C(k)^+$.

4. We end with some remarks on signs of functional equations for abelian varieties. Let A be an abelian variety of dimension g over k , and let $M_A = H_1(A)$, which is a motive of weight -1 and rank $2g$. The Weil pairing gives a non-degenerate, alternating bilinear form

$$(4.1) \quad M_A \times M_A \rightarrow \mathbf{Q}(1).$$

Let M_ρ be an orthogonal motive of weight 0 and rank 2 over k . Then M_ρ corresponds to a complex Galois representation of the form $\text{Ind}_K(\rho)$. Here K is the quadratic étale extension corresponding to the rank 1 orthogonal motive $\det(M_\rho)$, and ρ is a character of $\text{Gal}(\bar{k}/K)$ with trivial transfer to $\text{Gal}(\bar{k}/k)$. We view $\chi = \det(M_\rho)$ as a generalized ideal class character of k , and let T denote the set of real places of k which become complex in K .

Put $M = M_A \otimes M_\rho$, which is symplectic of weight -1 and rank $4g$. Local considerations (cf. [T, pg. 115]) lead one to predict that when the conductor ideal f_A of A is prime to the conductor of M_ρ ,

$$(4.2) \quad \epsilon(M) = \chi(f_A) \cdot (-1)^{g \cdot \#T}.$$

In fact, Rankin's method gives a proof of the functional equation of $L(M, s)$, with the above sign, when k is totally real and A appears as a new factor of the Jacobian of a Shimura curve of level N over k . Then $f_A = (N)^g$, and $L(M, s)$ is the product of g Rankin L -series, each with sign $\chi(N) \cdot (-1)^{\#T}$. The simplicity of this sign was the starting point for my work with Zagier on the conjecture of Birch and Swinnerton-Dyer.

5. Bibliography

- [A-F] J. Armitage and A. Fröhlich, Class numbers and unit signatures, *Mathematika* 14 (1967) 94–98.
- [Da] H. Davenport, *Multiplicative number theory*. Markham. 1967

- [De] P. Deligne, Valeurs de fonctions L et périodes d'intégrales. Proc. Sym. Pure Math. AMS 33 (1979), 313–346.
- [G-M] B.H. Gross and C.T. McMullen. Automorphisms of even unimodular lattices and unramified Salem numbers. Preprint.
- [T] J. Tate, Local constants. In: *Algebraic number fields*. Academic Press, 1979, 89–131.
- [W] A. Weil. *Basic number theory*. Springer-Verlag, 1967.