

CALCULUS ON WAVE FRONTS

OLIVER KNILL

ABSTRACT. We define a deformation of the exterior derivative that is a bounded operator and preserves the symmetries of the geometry. It satisfies a modified wave equation that honors the strong Huygens principle in all dimensions.

1. DEFORMED EXTERIOR DERIVATIVE

1.1. Denote by $\mathcal{H} = \bigoplus_{k=1}^q \overline{\Lambda}^k(M)$ the Hilbert space of **differential forms** on a Riemannian q -manifold (M, g) . It is the closure of the space of compactly supported differential forms $\Lambda(M)$ with respect to the norm defined by the inner product $\int_M \langle f, g \rangle_p dx$, where dx is the volume q -form on M . (See section 11.3 in [4]). With the **exterior derivative** d and its **adjoint** d^* , define the **Dirac operator** $D = d + d^*$ and the **Hodge Laplacian** $L = D^2 = d^*d + dd^*$. The wave equation $u_{tt} = -Lu$ has the explicit solution $u(t, x) = \cos(Dt)u(0, x) + t \operatorname{sinc}(Dt)u_t(u, x)$ in any dimension. It defines a global Hamiltonian flow in $\mathcal{H} \times \mathcal{H}$. The wave equation is equivalent to two Schrödinger equations $u_t = \pm iDu$ with a 2-dimensional solution space $c_+e^{iDt}u + c_-e^{-iDt}v$ in \mathcal{H} .¹

1.2. We work here on Euclidean space \mathbb{R}^q . But all notions that can be pushed over via coordinate changes hold also in general. In the case of a Riemannian manifold M , we would use the **exponential map** $\exp_p : T_pM \rightarrow M$ to transport the unit sphere $S_1(p) = \{v \in \mathbb{R}^q, |v| = 1\} \subset T_pM$ to the **wave front** $W_h(p) = \exp_p(hS_1(p)) \subset M$. For h smaller than the **radius of injectivity**, the wave front is diffeomorphic to a $(q - 1)$ -sphere. We can look at partial differential equations like the wave equation also on Riemannian manifolds. We work here for simplicity on flat $M = \mathbb{R}^q$.

1.3. The mathematics of wave fronts links differential geometry with partial differential equations. [3] outlines many of these relations. First investigations were done by Huygens who studied with waves and rays [2]. The principle that in odd dimensions, the solution $u(t, p)$ of the wave equation $u_{tt} = -Lu$ only involves the initial data on $W_h(p)$ is called the **strong Huygens principle**. It implies that there are **sharp wave fronts**. In even dimensions, also points closer to p do matter. Wave fronts are softer; there are **wakes**. The Kirchhoff solutions of the wave equation for scalar fields can be found for example in section 2.4 in [5].

1.4. Let $\phi_n(t)$ denote the unique solution of **Bessel equation**

$$(1) \quad f''(r) + (n - 1)\frac{f'(r)}{r} + f(r) = 0, f(0) = 1, f'(0) = 0.$$

Date: March 22, 2026.

¹When restricting to functions (0-forms), pseudo differential operators are involved. Extending the wave equation to all forms allows to avoid pseudo differential operators like $\sqrt{-\Delta}$. While pseudo differential operators are in general non-local, the expressions $\cos(Dt), \operatorname{sinc}(Dt)$ are both even and so functions of L . The solution formula is the same than for a wave equation on a discrete geometry like a finite abstract simplicial complex.

We have $\phi_1(r) = \cos(r)$, $\phi_2(t) = J_0(r)$, $\phi_3(t) = \text{sinc}(r) = \sin(r)/r$, $\phi_4(r) = 2J_1(r)/r$. Historically, Euler was one of the first to look in 1764 at the $q = 2$ wave equation $u_{tt} = u_{rr} + \frac{1}{r}u_r + \frac{1}{r^2}u_{\theta\theta}$ which with $u(t, r, \theta) = f(r) \cos(mt)$ leads to $f'' + f'/r + m^2f = 0$ and which after a variable change $r \rightarrow mr$ becomes $g'' + g'/r + g = 0$, the above Bessel equation in the case $q = 2$. (See [12] section 1.3). In an appendix, we say more about Bessel.

1.5. The following definition is new:

Definition 1. For $t > 0$ define the **deformed exterior derivative** using the ϕ_{q+2} Bessel function:

$$(2) \quad d_t = t\phi_{q+2}(tD)d .$$

1.6. Because $\phi_{q+2}(0) = 1$ and $d_0 = 0$, it satisfies $\lim_{t \rightarrow 0} d_t/t = \lim_{t \rightarrow 0} \frac{d}{dt}d_t = d$. The operator d_t is a bounded operator on \mathcal{H} because $r \rightarrow \phi_{q+2}(r)r$ is bounded. By the functional calculus for the self-adjoint operator tD (which is only densely defined), also $\phi_{q+2}(tD)tD$ is bounded so that $\phi_{q+2}(tD)d_t$ is bounded. A densely defined bounded operator extends to the entire Hilbert space.

1.7. For $q = 1$, where $\phi_3(r) = \text{sinc}(r) = \sin(r)/r$ and $D = d + d^* = \begin{bmatrix} 0 & -d/dx \\ d/dx & 0 \end{bmatrix} = id/dx$,

we have $t \text{sinc}(Dt)Du = \sin(Dt)u = [e^{iDt} - e^{-iDt}]u/(2i) = [e^{-\frac{d}{dx}t} - e^{\frac{d}{dx}t}]u/(2i) = i[u(x+t) - u(x-t)]/2$, using the **Taylor theorem** $e^{\frac{d}{dx}t}u(x) = \sum_{n=0}^{\infty} \frac{d^n}{dx^n} \frac{u(x)}{n!} = u(x+t)$. We see that on 0-forms u , its exterior derivative is the 1-form $d_t u(x) = [u(x+t) - u(x-t)]/2dx$. On a circle M , the only compact 1-manifold, all t rationally independent of the length of M have the same harmonic forms $d_t u = 0$. The Betti numbers b_k defined as the dimension of the kernel of $L_t = d_t^*d_t + d_t d_t^*$ restricted to \mathcal{H}_k are for almost all t the same as for the $L = dd^* + d^*d$.

1.8. Our original goal (pursued sporadically over the last two decades) had been to define an exterior derivative in arbitrary dimensions that preserves symmetries and does not require smoothness. It is the quest to pursue **multi-variable calculus without limits**. A concrete early question has been: is it true that if we take a 1-form f (vector field) on a 2-manifold and have all line integrals along all $p \in M$ and fixed $t > 0$ the wave fronts $W_t(p)$ are all zero for a fixed t , that f has to have zero classical curl? We would call the line integral $\int_{W_t(p)} f$ a **discretized curl** as it does not mind f to be only continuous or even piecewise continuous and allow M to be piecewise smooth only, like a **polyhedron** or have boundaries, where $W_t(p)$ is defined using the billiard flow. Green's theorem tells that the line integral along $W_t(p)$ is the integral of the curl df over the ball $B_t(p)$ (for small t at least). We can now answer this question and see that it depends on the spectrum of the form Laplacian on M as well as on t .

1.9. Any discretization in the form of a cell complex or triangulation would destroy rotational symmetry for example. Rotational symmetry is important in physics. The properties of the **hydrogen operator** for example explains much of the periodic elements in chemistry, the properties of Schwarzschild or Kerr metric explain measurable consequences in the general theory of relativity. In one dimensions, the scaled deformed derivative d_t/t is the standard symmetric difference gradient. Since $|t\phi_{q+2}(t)| \sim t^{(1-q)/2}$ for $t \rightarrow \infty$, we have $\|d_t\| \sim t^{(1-q)/2}$ for $t \rightarrow \infty$. Only in dimension 1, the exterior derivative does not go to zero and stays bounded for $t \rightarrow \infty$.

2. DEFORMED WAVE EQUATION

2.1.

Definition 2. *Define the Bessel acceleration*

$$(3) \quad D_{tt}u = u_{tt} + (q - 1)\left[\frac{u_t}{t} - \frac{u}{t^2}\right].$$

It needs that $u(0, x) = 0$ to have a limit at $t \rightarrow 0$.

2.2. It is a linear second order **Sturm-Liouville operator** that is singular at $t = 0$ and approaches u_{tt} for larger t . It factors as $D_{tt} = (\partial_t + \frac{q}{t})(\partial_t - \frac{1}{t})$. It is a Helmholtz Laplacian with zero energy and angular momentum 1.

2.3.

Definition 3. *Define also the Bessel acceleration*

$$(4) \quad d_{tt}u = u_{tt} + (q - 1)\frac{u_t}{t}.$$

It needs that $u_t(0, x) = 0$ to have a limit at $t \rightarrow 0$.

2.4. The differential equation $D_{tt} = a$ has for $u(0) = 0$ the solution family $u(t) = at^2/(q + 1) + tu'(0)$. This agrees with solutions of $\partial_t^2 u = 0$ except that the acceleration is for $q > 1$ smaller than the traditional acceleration u_{tt} . The equation $D_{tt}f = g$ is solved for analytic $g = \sum_{n=1}^{\infty} c_n t^{n-1}$ by noting that $D_{tt}f = t^{n-1}$ has the solution $-\frac{t(1-t^n)}{n(q+n)}$. If $g = \sum_{n=1}^{\infty} a_n t^{n-1}$ is bounded, there is exactly one bounded solution $f = \sum_{n=1}^{\infty} b_n t^n$. The condition $f(0) = 0$ is one of boundary condition, the boundedness forces the initial velocity and so fixes a second boundary condition.

2.5. Also this definition of a partial differential equation appears to be new:

Definition 4. *Define the deformed wave equation*

$$(5) \quad D_{tt}u + Lu = 0,$$

and

$$(6) \quad d_{tt}u + Lu = 0,$$

2.6. For $q = 1$, these are both the usual wave equation. The motivation to define these PDE's was that we will see that $u(t, p) = d_t f(p) = t\phi_{q+2}(tD)df$ is an explicit solution to $(D_{tt} + L)u = 0$ with $u(0, p) = 0$ and $\lim_{t \rightarrow 0} \frac{d}{dt} d_t f(p) = df(p)$. Also, $u(t, p) = \phi_q(tD)df(p)$ is an explicit solution of $(d_{tt} + L)u = 0$ with $u(0, p) = df(p)$ and $u_t(0, p) = 0$.

2.7. In order that the solution of the deformed wave equation $(D_{tt} + L)u = 0$ works we need in the case $q \neq 1$ that $u(t, x) = t^2 v(t, x)$ and especially $u(0, x) = 0$. In order that the solution of the deformed wave equation $(d_{tt} + L)u = 0$ makes sense, we need $u_t(0, x) = 0$. In all dimension q , the initial value problems with initial velocity or initial positions can be solved explicitly. The strong Huygens principle holds.

2.8. Asymptotically, for $t \rightarrow \infty$, for any q , the solutions of the deformed wave equation move like the wave equation $u_{tt} + Lu = 0$. While the later has the explicit solution $u(t, p) = \cos(Dt)u(0, p) + \text{sinc}(Dt)u'(0, p)$, this only honors the **weak Huygens property**: the classical wave equation features “wakes” in even dimensions as the explicit solution formulas of Kirchhoff show. By modifying the time acceleration from ∂_t^2 to D_{tt} slightly, the strong Huygens principle is achieved in arbitrary dimensions. This can be useful when trying to interpret a wave front in a space-time 4-manifold as “space”.²

3. RESULTS

3.1. Here are our main results about the bounded operator $d_t : \mathcal{H} \rightarrow \mathcal{H}$. Similarly as the classical exterior derivative d , also d_t maps k -forms \mathcal{H}_k to $(k + 1)$ -forms \mathcal{H}_{k+1} but it is a bounded operator. From $d^2 = 0$ follows immediately that also $d_h^2 = 0$. Harmonic forms stay harmonic.

Theorem 1 (Wave equation). *a) The form $u(t, p) = d_t f(p)$ solves $(D_{tt} + L)u = 0$ with initial position $u(0, p) = 0$ and initial velocity $\lim_{t \rightarrow 0} \frac{d}{dt} u_t(0, p) = df(p)$.*

b) The form $u(t, p) = \phi_q(tD)df(p)$ solves $(d_{tt} + L)u = 0$ with initial position $u(0, p) = df(p)$ and initial velocity $u_t(0, p) = 0$.

Proof. a) Just verify that $u(t, x) = t\phi_{q+2}(tD)Y(x)$ solves $u_{tt} + u_t(q-1)/t + u(q-1)/t^2 + D^2u = 0$ provided that $\phi''(r) + (q+1)\phi'(r)/r + \phi(r) = 0$ with $f(0) = 1, f'(0) = 0$. For $r = tD$ we get $\phi''(tD) + (q+1)\phi'(tD)/(tD) + \phi(tD) = 0$. Differentiate u twice using chain and product rule:

$$\begin{aligned} u &= t\phi(tD)Y \\ u_t &= t\phi'(tD)DY + \phi(tD)Y \\ u_{tt} &= t\phi''(tD)D^2Y + \phi'(tD)DY + \phi'(tD)DY. \end{aligned}$$

Multiply the first by D^2 , the second by $(q-1)/t$ and switch $q-1$ to $q+1$ using $t\phi'(tD)D^2Y(q-1)/(tD) = t\phi'(tD)D^2Y(q+1)/(tD) - 2\phi'(tD)DY$.

$$\begin{aligned} D^2u &= t\phi(tD)D^2Y \\ u_t(q-1)/t &= t\phi'(tD)D^2Y(q+1)/(tD) + \phi(tD)Y(q-1)/t - 2\phi'(tD)DY \\ u_{tt} &= t\phi''(tD)D^2Y + 2\phi'(tD)DY \\ -u(q-1)/t^2 &= -\phi(tD)Y(q-1)/t. \end{aligned}$$

The sum of the terms on the left hand side add up to 0 as ϕ satisfies the deformed wave equation. The right hand side add up also to zero because ϕ satisfies the Bessel equation $\phi'' + (q+1)\phi'/(tD) + \phi = 0$. b) By taking q rather than $q+2$, the forms $u(t, p) = \phi_q(Dt)f(p)$ have solve the **zero angular momentum PDE**

$$u_{tt} + (q-1)\frac{u_t}{t} = -D^2u$$

with $u(0, p) = df$ and $u_t(0, p) = 0$. The verification is the same. \square

3.2. Because the functions ϕ_k are all bounded, d_t is a bounded operator on the Hilbert space \mathcal{H} of forms, for all $t \geq 0$. (The case $t = 0$ holds trivially because $d_t = 0$ for $t = 0$.)

²As for references, see e.g. [?] 6.9.64 which uses the Fourier picture that works in arbitrary dimensions. Or [13] Chapter I, section 3, for wave equations on Riemannian manifolds.

3.3. On other original motivation was to search for modified wave equations that solve

$$\phi_n(Dt)u(0, p) + t\phi_{n+2}(Dt)u_t(0, p)$$

because $u(t, p) = \phi_1(Dt)u(0, p) + t\phi_3(Dt)u_t(0, p)$ satisfies the classical wave equation $u_{tt} + D^2u = 0$ in all dimensions $q \geq 1$, as $\phi_1(r) = \cos(r)$, $\phi_3(r) = \sin(r)/r = \text{sinc}(r)$. It turned out that we can indeed find such equations but that in the position and velocity case, different differential equations appeared.

3.4. As we have seen in the introduction, the 1-dimensional case $q = 1$ is a prototype case. It is historically important as d'Alembert was a pioneer in partial differential equations. Before him, wave phenomena had been discussed in a descriptive way. The deformed wave equation agrees in the case $q = 1$ with the wave equation $u_{tt} + Lu = 0$. In one dimensions, $u(t, p) = t\text{sinc}(Dt)f$ satisfies the wave equation with $u(0) = 0, u_t(0) = df$. It is the **d'Alembert solution** $u(t, p) = [u(0, p+t) - u(0, p-t)]/2$. The expression $u(t, p)/t$ is a **discrete numerical derivative** that converges for $t \rightarrow 0$ to the usual derivative df . In the limit $t \rightarrow 0$, we need the initial 0-form f to be differentiable. The calculus for d needs limits, the calculus for d_t does not need any limits for $t > 0$.

3.5. The second main result is about the modified partial differential equation:

Theorem 2 (Strong Huygens). *a) $u(t, p) = d_t f(p) = t\phi_{q+2}(tD)df$ only uses f on $W_t(p)$.
b) $u(t, p) = \phi_q(tD)df$ only uses f on $W_t(p)$.*

Proof. We focus on a) and give in the appendix the argument to reduce b) to a) using the identity $\phi_q(r) = r^{1-q}(\phi_{q+2}(r)r^q)'/q$.

(i) The statement is first first shown for $(q-1)$ -forms f , where it reduces to the **ball average formula** applied to the q -form $g = df$. There is a Taylor expansion [8] for the ball expectation $|B_t|^{-1} \int_{B_t} g = \sum_n c_n t^{2n} L^n g$. We noticed that this matches with $\phi_{q+2}(tD)g$. **Stokes theorem** then shows that $d_t f$ only invokes f on the wave front $W_t(p)$. Writing $\frac{1}{|S_t|} \int_{S_t(p)} f(x) dS$ as $\sum_{n=0}^{\infty} c_n t^{2n} L^n f(p)$ is a **Pizzetti type formula**, named after **Paolo Pizzetti** [9] who derived such a formula first in dimension $q = 3$.

(ii) To see the general case for k -forms with $k < q-1$, we use **inner derivatives** i_X and induction $k \rightarrow k-1$ starting with the induction assumption $k = q-1$, seen in (i). If X is a vector field, denote by $i_X : \Lambda_{k+1} \rightarrow \Lambda_k$ the **inner derivative** and by $L_X = i_X d + di_X$ the **Lie derivative** preserving k -forms. For background, see [1]. Expressing the Lie derivative as an anti-commutator of exterior and interior derivative is known as **Cartan's magic formula**. In order to extend the strong Huygens property from k -forms to $(k-1)$ -forms, use a k -form f and a constant vector field for which $L_X g = 0$ so that $i_X df = -di_X f$ and hence $i_X d_h f = -d_h i_X f$. The Huygens principle so holds for any $(k-1)$ -form $i_X f$ provided f was L_X invariant.

(iii) In order that (ii) works for all k -forms, we need to show for all $k \leq q-1$, any k -form f can be written as a linear combination of k -forms that are in the kernel of some L_X . This however follows from the **polarization identity**. It allows to write any monomial $\prod_i x_i^{n_i}$ of degree $n = \sum_j n_j$ as a linear combination of terms $(\sum_i s_i x_i)^n$, where $s_i \in \{-1, 1\}$. Every differential k -form adf can be written as a combination of forms $i_X g$, where g is a k -form such that $L_X g = 0$. □

3.6. Lets look at the case $q = 3$ and $k = 2$. We want to show that we can get any 1-form as a linear combination of forms $i_X g$, where g is a 2-form and $L_X g = 0$. For example, the 2-form $g = (x + y - z)^n d(x - y) \wedge dy$ satisfies $L_X g = 0$ for $X = (1, -1, 0)$ and $I_X g = (x + y - z)^n dy$.

Using polarization, any monomial 1-form $x^k y^l z^m dy$ can be realized in such a way. Again by linearity, every polynomial 1-form $Q(x, y, z)dy$ has this property. Using linear combinations of such forms, we can realize any 1-form $P(x, y, z)dx + Q(x, y, z)dy + R(x, y, z)dz$ as a linear combination of inner derivatives of 1-forms $i_X g$, where g is a 2-form satisfying $L_X g = 0$.

3.7. In the proof, we used the ball average formula in Euclidean space and that in flat Euclidean space we can use vector fields X that are parallel. In a curved space, there are no parallel vector fields in general. Also ball averages are different. But once we have established the strong Huygens principle for $D_{tt}u = -D^2u$ in Euclidean space \mathbb{R}^n we have, (like in the usual wave equation $u_{tt} = -D^2u$), that the strong Huygens property goes over to the Riemannian manifold case, using that the exponential map near a point is just a coordinate change and that $D = d + d^*$ is a coordinate independent notion.

4. HARMONIC FORMS

4.1. Define $D_h = d_h + d_h^*$ and $L_h = D_h = d_h^* d_h + d_h d_h^*$ so that with $\psi_n(r) = \phi_n(r)r$, we have $D_h = \psi_{q+2}(hD)$. Since $\psi_{q+2}(0) = 0$, all classically harmonic forms f are also **h -harmonic**: they satisfy $L_h f = 0$. For any compact Riemannian manifold M there is a Hilbert space \mathcal{H} and a Dirac operator $D = d + d^*$. The set-up does not require coordinates and locally, we can parametrize a coordinate patch and so have the above results for h smaller than the **radius of injectivity** $\rho(M)$ of M . In the compact space D has discrete spectrum so that $\psi_h(D)$

Theorem 3 (Harmonic forms). *If M is a compact Riemannian manifold, then there are no new h -harmonic forms for almost all $h < \rho(M)$.*

4.2. For example, if $M = \mathbb{T}^1$ with a metric giving it length is 1, then for all irrational $0 < h < 1/2$, no new h -harmonic forms are obtained for d_h . For rational $h > 0$, all $2h$ -periodic function f or 1-form is h -harmonic because $2d_h f(p) = f(p+h) - f(p-h) = 0$.

4.3. If cohomology is defined via Hodge, the **Betti numbers** b_k can be defined as the dimension of the space of h -harmonic k -forms.

Corollary 1. *The Hodge cohomology does not change for most h .*

5. PHYSICS

5.1. Our goal had been to generalize the 1-dimensional discrete derivative to higher dimensions without introducing any symmetry breaking in M . The derivative d_h honors all Euclidean symmetries on \mathbb{R}^q . Symmetries are important because they are related to **conservation laws**. This **Nöther relation** links momentum or angular momentum conservation with translation and rotational symmetry for example. Any triangulation or lattice approach would break symmetries.

5.2. If G is a symmetry group on M , it has a representation as unitary operators in \mathcal{H} . If T is a rotation for example, then $f \rightarrow f(T)f$ defines a unitary U_T in \mathcal{H} that commutes with d . It still commutes with the new derivative d_h .

Theorem 4 (Symmetry). *If a unitary operator on \mathcal{H} commutes with d , it continues to commute with d_h .*

5.3. The fact that symmetry is preserved and allows to transition easily from a continuum to a quantized situation. The hydrogen operator $L = -h^2\Delta/2m - \frac{e^2}{4\pi\epsilon_0}/r$ for example on 0-forms has eigenvalues $\lambda_n = -C/n^2$ with $C = me^4/(2(4\pi\epsilon_0)^2h^2)$. Now $d_h \sim dh$ and $L_h \sim Lh^2 = -\Delta h^2$. The Hydrogen operator in spherical coordinates is now $O(h^4)$ close to the deformed operator $L_h/m - \frac{e^2}{4\pi\epsilon_0}/r$. The eigenvalues in the deformed situation are h^2 close. In the case of the Hydrogen atom, one has $C = mc^2\alpha^2/2 = 13.605...eV = 2.1798... \cdot 10^{-18}J$, with the **fine-structure constant** $\alpha \sim 1/137$. This C is the **ground state energy**. When replacing L with L_h , the spectrum is indistinguishable. Modeling the Hydrogen atom using finite geometries would be hard because spherical coordinates are not available in such a frame work. If we replace $-h^2\Delta$ with the zero Laplacian in L_h and take h to be of Planck scale, all the physics would look the same.

5.4. The **Maxwell equations** can be played on any Riemannian manifold ³ M with exterior derivative d . Given a 1-form j , it is the system of partial differential equation $dF = 0, d^*F = j$ for an unknown 2-form F . If $F = dA$ (which is possible if M is simply connected or even if $b_1 = 0$), then Maxwell means $d^*dA = j$. Under a **Coulomb gauge** $A \rightarrow A + df$, we can assume $d^*A = 0$ which means that the Maxwell equations are reduced to the **Poisson equation** $LA = (dd^* + d^*d)A = j$. If j is in the orthogonal complement of the harmonic forms $\ker(L)$, we have a unique solution $A = L^{-1}j$, (where L^{-1} is the pseudo inverse). The electro-magnetic field then is given as $F = dA$. In vacuum and in a space time manifold with signature $-+++$, the equation $Lu = 0$ is a wave equation. In our case, where time is separated from space, we have the d'Alembert operator $\square = \partial_t^2 + L$ so that the wave equation in space dimension q is $Au = 0$. The deformed d'Alembert operator is $\square = D_{tt} + L$. We have seen that $\square u = 0$ with $u = 0, \lim_{t \rightarrow 0} u_t(t, x) = du(x)$ is solved by $u(t, x) = d_t f(x) = t\phi_{q+2}(tD)df$.

6. QUESTIONS

6.1. A discrete time wave evolution $T : (u, v) \rightarrow (D_h u - v, u)$ is defined, if h is such that D_h has norm smaller than 1. It is conjugated to $(e^{i\tilde{D}_h}, e^{-i\tilde{D}_h})$ for a slightly changed \tilde{D} . See [6]. For small h , this is almost indistinguishable from the actual unitary evolution. What are the properties of this wave evolution? Do we still have the strong Huygens principle? We liked the discrete cellular automaton type time evolution for numerical purposes so that the wave or Schrödinger evolution in the discrete has finite speed of propagation. On a discrete geometry with continuous time t , there is no locality for the wave equation $u_{tt} = -Lu$; we need also time to be discrete. The above evolution T has a finite speed of propagation because d_h only taps into parts of space in distance h .

6.2. The spectrum of the deformed Laplacian $L_h = d_h^*d_h + d_h d_h^* = D_h^2$ is determined by the spectrum of L if $h > 0$. (We write here h rather than t because one of the applications we have in mind is to use d_h for small h , where h is on the order of the **Planck constant**. As $d_h^* = h\phi_{q+2}(hD)d^*$, we have $D_h = d_h + d_h^* = \phi_{q+2}(hD)hD$. This implies that D_h has the same eigenfunctions than D and $\lambda_j(D_h) = \psi(\lambda_j(D))$, where $\psi(r) = \phi_{q+2}(r)r$, a bounded function. Similarly, also the eigenfunctions of L_h are the same than the eigenfunctions of L and $\lambda_j(D_h) = \psi(\lambda_j(L))$. It would be nice to use L_h as a numerical tool. Is there a natural modification $V_h(r)$ of the potential $V(r) = 1/r$ so that we have a bounded analog of the Hydrogen operator in the form of $L_h + V_h$?

³pseudo Riemannian manifold

6.3. The geometry of wave fronts can become very complicated, even in integrable situations like the ellipsoid. Can we relate the spectrum of L and so the spectrum of d_t with the wave front? It would be nice for example to see spectrally whether wave front become dense. In the case of a round sphere, the wave front oscillates and stays bounded.

6.4. The classical wave equation $u_{tt} + Lu = 0$ has **Klein-Gordon generalizations** $u_{tt} + (L + m)u = 0$, which does not satisfy the strong Huygens principle. What happens in the modified case $u_{tt} + (q - 1)u_t/t + (L + m)u$ or $u_{tt} + (q - 1)u_t/t + (q - 1)u/t^2 + (L + m)u$? Are there cases, where we have sharp wave fronts?

6.5. If we look at a 4-manifold and consider the front $W_t(p)$ to be space, we have a **space-time model** which locally shows Lorentz symmetry. Moving p around in M is a space time change. Every starting point p defines a different space-time structure. On a Riemannian manifold, the radius of injectivity in general depends on p . So, there are points p_1 , where the **space 3-manifold** $W_t(p_1)$ is regular while for p_2 the **space 3-manifold** $W_t(p_2)$ has developed a singularity as t is larger than the radius of injectivity. The space-time structure depends on the starting point p , where the space-3-manifold is a point. On a 4-dimensional ellipsoid as space time 4-manifold, there are initial points for which there is a “big crunch”, a time where $W_t(p)$ is again a point. For most other points, the space manifold expands indefinitely. This is highly non-trivial even in toy models like circular billiard, a 2-manifold with boundary where the length of the wave front of a ”a drop of water in a cup” asymptotically behaves like $|W_P(t)| = \arcsin(|P|)t + o(t)$ [11].

6.6. For a finite abstract simplicial complex G with n elements, there is an exterior derivative d on the finite dimensional Hilbert space $l^2(G)$. Functions on $G_k = \{x \in G, |x| = k - 1\}$ are k -forms. The exterior derivative maps $l^2(G_k)$ to $l^2(G_{k+1})$. If x is a simplex, then $df(x)$ is the flux of f through the boundary δx . We do not need to deform d because it is already a bounded operator. If G is a q -manifold and $D = d + d^*$ is the Dirac matrix, a $n \times n$ matrix, then we can still do a deformation $D_t = \phi_{q+2}(Dt)tD$. We can also look at the now ordinary differential equation $u_{tt} + (q - 1)[u_t/t - u/t^2] = -D^2u$. We can still deform exterior derivatives using the geodesic flow, but what happens there is still unexplored.

7. CONCLUSION

7.1. We have deformed the exterior derivative d so that d_h has become a bounded operator on \mathcal{H} . Boundedness was important. Operators like D or $L = D^2$ were only densely defined self-adjoint operators on the Hilbert space \mathcal{H} . Having a bounded operator implies that the operator naturally extends to all elements of the Hilbert space \mathcal{H} . The new Laplacian $L_h = (d_h + d_h^*)^2$ behaves like h^2L for small h .

7.2. The **Stokes theorem** $\int_G dF = \int_{\delta G} F$ for smooth k -forms in a compact $(k + 1)$ manifold G with k -manifold δG as boundary now reads as

$$\langle G, d_t F \rangle = \langle d_t^* G, F \rangle .$$

Fields F and smooth geometries G (which are examples of **de Rham currents**) are now on the same footing as both are elements in the same Hilbert space. There is a symmetry between fields and geometries.

7.3. It follows for example that we can relax about the regularity of geometries. We have a bounded exterior derivative on manifolds which are singular like polyhedra or varieties on which there is a natural geodesic metric. Geometries can also be fractal. If G for example is the **Koch fractal**, a region G in \mathbb{R}^2 with the fractal boundary δG called Koch curve. If F is a 1-form, we can look at the curl $d_t F$ for some small t and integrate this over G . It is just an inner product $\langle G, d_t F \rangle$ as we can see also write $G = 1_G(x, y) dx dy$ as a 2-form. Now this is $\langle d_t^* G, F \rangle$ and $d_t^* G$ is an L^2 1-form a relatively regular de-Rham current, compared with the traditional boundary.

7.4. This derivative **quantizes distances**: only $f(q)$ for q in distance h from p matter when evaluating $d_h f(p)$. We have a **calculus without limits** because we do not need to take the limit $h \rightarrow 0$. In physical circumstances, taking a Planck distance size h produces the same result in traditional physics. The derivative in principle works for any Riemannian manifold for which singularities in the metric are of codimension 2. On a solid polyedron for example, a 3-manifold with boundary, the calculus works as the singularities that affect the wave fronts are locate on vertices only.

7.5. The calculus resembles the discrete case if M is an abstract simplicial complex, where $df(p)$ for a k -simplex p only needs to know q , where q are the **boundary simplices** of p of codimension 1. We can in discrete manifolds for example define a new exterior derivative that instead of summing up over all boundary simplices sums up over all boundary simplices of all adjacent simplices.

APPENDIX: BESSEL

7.6. The Bessel type differential equation $u'' + (q - 1)u'/r + u = 0$ for a function $u(r)$ is the **radial Helmholtz equation** for $k = 1$ in dimension q , with energy 1 and angular momentum 0. We wrote $\phi_q(r)$ for its solution given the initial condition $u(0) = 1, u'(0) = 0$. The eigenvalue problem $\Delta u + c^2 u = 0$ becomes in spherical coordinates

$$u_{rr} + \frac{(q-1)}{r}u_r + \frac{1}{r^2}\Delta_{S^{q-1}}u + c^2u = 0.$$

If r is replaced by r/c , the energy can be scaled to become 1. With a substitution $\nu = (q - 2)/2$ and $y(r) = r^\nu u(r)$ it becomes the **standard Bessel equation**

$$y'' + \frac{y'}{r} - \nu^2 \frac{y}{r^2} + y = 0$$

The solutions are Bessel functions of order ν given by $r^\nu J_\nu(r)$ and $r^\nu Y_\nu(r)$, which are known as **Bessel functions of the first kind and second kind**.

7.7. Related to the deformed wave equation, we had introduced the **modified acceleration operator** $u'' + (q - 1)[\frac{u'}{r} - \frac{u}{r^2}]$. It is the **radial Helmholtz operator**

$$u'' + (q-1)\frac{u'}{r} - l(l+q-2)\frac{u}{r^2} + c^2u = 0$$

with **angular momentum** $l = 1$ and **energy** $c = 0$ in **dimension** q .

7.8. For angular momentum $l = 1$ and energy $c = 0$, this radial Helmholtz operator becomes

$$u'' + (q - 1)\frac{u'}{r} - (q - 1)\frac{u}{r^2}.$$

For $l = 0$ and $c = 0$, it is the operator

$$u'' + (q - 1)\frac{u'}{r}.$$

These are the time acceleration operators we have used in the modified wave equations.

7.9. The equation

$$u'' + (q - 1)\frac{u'}{r} - (q - 1)\frac{u}{r^2} + c^2u = 0$$

becomes after a substitution $y = r^{((q-2)/2)}u$ the Bessel equation

$$y'' + \frac{y'}{r} - \left(\frac{q}{2}\right)^2\frac{y}{r^2} + c^2y = 0.$$

7.10. The solutions ϕ_q to the Bessel equation

$$u''(r) + (n - 1)\frac{u'(r)}{r} + u(r) = 0, u(0) = 1, u'(0) = 0$$

are also **hypergeometric functions**. One can see this from the Taylor expansion $\phi_q(r) = \sum_{k=0}^{\infty} b_k r^{2k}$ with $1/b_k = B(q, k) = \prod_{j=1}^k (-2j)(q - 2 + 2j)$.

7.11. To summarize, we have now seen four different ways to look at $\phi_q(r)$. It is a solution of a radial Helmholtz equation, expressible using Bessel functions of the first kind, given as a confluent hypergeometric functions or given in a series expansion which appeared, when looking at sphere averages.

Lemma 1. $\phi_q(r) = F_1\left(\frac{q}{2}, -\frac{r^2}{4}\right) = J_{\frac{q}{2}-1}\Gamma\left(\frac{q}{2}\right)\left(\frac{r}{2}\right)^{\frac{q}{2}-1} = \sum_{k=0}^{\infty} b_k r^{2k}$.

7.12. While already explained by changes of variables, it is best illustrated in the language of a computer algebra system which has these functions already baked in:

```

B[q-, k-] := Product[(-2 j) (q-2+2 j), {j, 1, k}];
CheckBesselIdentities[q-]:=Module[{phi, hyper, bessel, series},
phi = f[r] /. First[DSolve[{f''[r]+(q-1)*f'[r]/r+f[r]==0, f[0]==1, f'[0]==0}, f[r], r]];
hyper = Hypergeometric0F1[q/2, -r^2/4];
bessel = BesselJ[q/2-1, r] Gamma[q/2]/(r/2)^(q/2-1);
series = Sum[r^(2 k)/B[q, k], {k, 0, Infinity}];
Print[{phi, hyper, bessel, series}];
FullSimplify[phi==hyper==bessel==series];
Table[CheckBesselIdentities[q], {q, 1, 10}]

```

7.13. From the classical Bessel identity $(r^n * J_n(r))' = r^n * J_{n-1}(r)$, one gets that the functions ϕ_q satisfy the following recursion:

Lemma 2. $(\phi_{q+2}(r)r^q)' = q\phi_q(r)r^{q-1}$.

7.14. This shows that if $d_t f = t\phi_{q+2}(tD)df$ satisfies the strong Huygens property, also $\phi_q(tD)df$ satisfies the strong Huygens property. Since $\phi_{q+2}(tD)df$ only depends on $f \in W_t(p)$, we have for $h < t$ that $\phi_{q+2}(tD)df$ does not involve $f(x)$ with $x \in B_h(p)$. So, also the traditional time derivative $D\phi'_{q+2}(tD)df$ does not depend on $x \in B_h(p)$ and also $D\phi'_{q+2}(tD)(tD)^{1-q}df$ not, where D^{-1} is understood as the pseudo inverse. But the above identity sees this as $Dq\phi_q(tD)df$. So, also $\phi_q(tD)df$ does not depend on $x \in B_h(p)$.

APPENDIX: KIRCHHOFF EQUATIONS

7.15. Reusing the constants $B(q, k) = \prod_{j=1}^k (-2j)(q + 2j)$, [8] gives the following **Taylor formula** for the mean value over a ball

$$E_{B_t}[g] = \sum_{k=0}^{\infty} \frac{t^{2k}}{B(q+2, k)} L^k g$$

and over a wave front W_t and

$$E_{W_t}[g] = \sum_{k=0}^{\infty} \frac{t^{2k}}{B(q, k)} L^k g .$$

4

7.16. If $g = df$, where f is a $(q-1)$ -form, then $E_{B_t}[g]$ is by **Stokes theorem** equal to the normalized flux $(1/|B_t|) \int_{W_t} f$ through the sphere W_t . Hence, the sphere and ball volumes relate as $|W_t|t = q|B_t|$ we have $(t/q)/|B_t| = 1/|W_t|$ and so $E_{W_t}[f] = (t/q)E_{B_t}[g]$.

7.17. We observed that $E_{B_t}[g] = \phi_{q+2}(Dt)g$, and that $E_{W_t}[g] = \phi_q(Dt)g$, where ϕ_q solves the **Bessel differential equation** $f_{rr} + (q-1)f_r/r + f = 0$ with $f(0) = 1, f_r(0) = 0$.

7.18. In the following, we could assume f to be a $(q-1)$ form in \mathcal{H}_{q-1} but then it would only hold for almost all p . If f is assumed to be continuous, then it holds for all center points p .

Corollary 2. *If f is a continuous $(q-1)$ -form, then $d_t f(p) = t\phi_{q+2}(Dt)df(p)$ is q times the average flux of f through $W_t(p)$.*

7.19. This illustrates the strong Huygens principle for the deformed wave equation. But this interpretation as an average flux only holds in Euclidean space: we have used a relation between ball and sphere volumes which do not hold in a general Riemannian manifold. In a general Riemannian manifold M , it is still a multiple of the flux and we still have a strong Huygens principle.

7.20. The **Kirchhoff solutions** of the classical wave equations generalizes to all dimensions (see [5] page 73 or [10] Theorem 6.9.8, Theorem 6.9.10). One in general assumes the wave equation to evolve scalar fields g . It also applies to q -forms = volume forms gdx , which is the case we use.

7.21. Define $E_{W_t}[g]$ and $E_{B_t}[g]$ as averages over the sphere or ball. Define the pseudo differential operators $T_t = [(\frac{1}{t}\partial_t)]^{1/2}$ and $\gamma_q = 1, 3, 5 \dots (q-2)$. Assume the initial condition g and initial velocity h are given. In odd dimensions:

$$u = \frac{1}{\gamma_q} \partial_t T_t^{(q-3)/2} (t^{q-2} E_{W_t}[g]) + \frac{1}{\gamma_q} T_t^{(q-3)/2} (t^{q-2} E_{W_t}[h]) .$$

Define in $B_t(x)$ the function $\tilde{g}(y) = g(y)/\sqrt{t^2 - |x-y|^2}$. In even dimensions,

$$u = \frac{1}{\gamma_q} \partial_t T_t^{(q-2)/2} (t^q E_{B_t}[\tilde{g}]) + \frac{1}{\gamma_q} T_t^{(q-2)/2} (t^q E_{B_t}[\tilde{h}]) .$$

⁴[8] works with scalar Laplacians $\Delta = -L_0$ and scalar functions $g \in C^{2p}$ and finite Taylor sums. We only need the formula for polynomials, as they are dense in L^2 . In general, polynomial differential forms are dense in the Hilbert space \mathcal{H} .

The **strong Huygens principle** is that the solution $u(x, t)$ in odd dimensions only depends on information of g, h on the wave front W_t and not the entire ball. The **weak Huygens principle** is that the solution $u(x, t)$ in even dimensions only depends on information of g, h in the wave ball B_t , not outside. There is **finite speed of propagation**.

APPENDIX: POLARIZATION

7.22. Given n variables x_1, \dots, x_n in an arbitrary commutative ring \mathcal{R} with 1. Define $S = \{-1, 1\}^n$, the set of all ± 1 strings of length n . The **polarization identity** is

Lemma 3. $n! \prod_{i=1}^n x_i = |S|^{-1} \sum_{s \in S} (\prod_j s_j) (\sum_i s_i x_i)^n$.

7.23. It shows that every element in the polynomial ring $\mathcal{R}[x_1, \dots, x_n]$ can be written as a linear combination of powers of linear functions. This can then be applied also to cases, where some x_i are the same, allowing so to write any monomial $x_1^{m_1} x_2^{m_2} \dots x_k^{m_k}$ as a linear combination of powers $(\sum_{i=1}^k a_i x_i)^m$ of linear functions, where $m = m_1 + m_2 + \dots + m_k$ is the degree of the monomial.

7.24. The best known case of the polarization identity is the **parallelogram law**. It writes xy as $[(x+y)^2 - (x-y)^2 - (-y+x)^2 + (-x-y)^2]/8$ where $8 = 2!*2^2$. The parallelogram law is used to show that in any Hilbert space, we only need to know **lengths** to get **angles** as the **dot product** in a Hilbert space is recoverable from **norms**. It can be used in statistics as $\text{Cov}[X, Y] = (\text{Var}[X + Y] - \text{Var}[X - Y])/4$. An other example is $xyz = \frac{1}{3!2^3} \sum_s s_1 s_2 s_3 (s_1 x + s_2 y + s_3 z)^3$. It can also cover cases like $x^2 y$ and see it as a linear combination of cubic powers $(ax + by)^3$ of linear functions in x and y .

7.25. The proof of the polarization property is done by averaging over the set S of all possible ± 1 vectors s . By symmetry, it is a multiple of a monomial. The symmetrization renders all signs the same, so that it must be $C \prod_{i=1}^n x_i$ for some constant C . To find the constant, plug in $x_i = 1$ and compute $\sum_{s \in S} (\prod_j s_j) (n - 2k)^n$. It reduces to show that $R = \frac{1}{n!} \sum_{k=0}^n \binom{n}{k} (-1)^k (n - 2k)^n = 2^n$.

7.26. Plugging in $(n - 2k)^n = \sum_{j=0}^n \binom{n}{j} n^{n-j} (-2k)^j$ gives

$$R = \frac{1}{n!} \sum_{j=0}^n \binom{n}{j} n^{n-j} (-2)^j \sum_{k=0}^n (-1)^k \binom{n}{k} k^j .$$

The finite-difference identity show that $\sum_{k=0}^n (-1)^k \binom{n}{k} k^j = 0$ is only nonzero for $j = n$, where it is $(-1)^n n!$.

APPENDIX: COMPUTER ALGEBRA RELATED TO THEOREM 1

7.27. We list now some computer algebra that we had used to find the partial differential equation. ⁵ Since Mathematica has built in D as the derivative operator, we use d for the Dirac matrix D . Note that all the verification of Theorem 1 works in the Abelian operator algebra generated by the operator D , so that D can just be treated as a variable. The two computations are almost identical. In the first case, we look at $u(t, x) = t\phi_{q+2}(tD)Y$ with $Y = df$, and verify, that it satisfies $u_{tt} + (q - 1)[u_t/t - u_{tt}/t^2] = -D^2u$.

⁵It was pretty much trial and error, first mostly done by hand, going through a few failures at first.

```
q=5; g=First[f[r]/.DSolve[{f'[r]+(q+1)f'[r]/r+f[r]=0,f[0]==1,f'[0]==0},f[r],r]];
phi=g/.r->d t; u=t phi d f; {Limit[u,t->0], Limit[D[u,t],t->0]}
FullSimplify[D[u,{t,2}]+(q-1)D[u,t]/t-(q-1)u/t^2==d^2*u]
```

7.28. In the second case, we look at $u(t, x) = \phi_q(tD)Y$ with $Y = df$. Note that now, there is no t in front of the solution $u(t, x)$ and that instead of ϕ_{q+2} we have now ϕ_q . The curve $u(t, x)$ satisfies $u_{tt} + (q - 1)[u_t/t] = -D^2u$ with initial conditions $u(0, x) = df$ and $u_t(0, x) = 0$.

```
q=7; g=First[f[r]/.DSolve[{f'[r]+(q-1)f'[r]/r+f[r]=0,f[0]==1,f'[0]==0},f[r],r]];
phi=g/.r->d t; u= phi d f; {Limit[u,t->0], Limit[D[u,t],t->0]}
FullSimplify[D[u,{t,2}]+(q-1)D[u,t]/t==d^2*u]
```

APPENDIX: COMPUTER ALGEBRA FOR SPHERE AVERAGING

7.29. As for the sphere average formulas, this also started first by experimenting by trial and error during the winter of 2010/2011. We noticed then experimentally that Taylor coefficients of flux are related to Bessel functions but could not prove this yet. The sphere and ball average formulas due to Ovall [8] (which in the case $d = 3$ are known as Pizzetti type formulas) made things clear in the winter of 2025/2026. What the code does in dimension $q=1,2,3$ is to take a random polynomial $(q - 1)$ -form and integrate the divergence over a ball of radius r , then make the Taylor expansion in r , and then notice that this is related to coefficients appearing in Bessel functions.

```
q=1; R:=Random[Integer,5]; L[f_.]:=D[f,{x,2}]; F=Sum[R x^n,{n,0,9}];
div=D[F,x]; X=Integrate[ div /. x->r,{r,-s,s}];
c=Simplify[CoefficientList[Series[X/(2s),{s,0,12}],s]];
c0=c[[1]]; c1=c[[3]]; c2=c[[5]]; c3=c[[7]]; c4=c[[9]];
a0=div /. x->0 /. y->0; a1=L[div] /. x->0 /. y->0;
a2=L[L[div]] /. x->0 /. y->0; a3=L[L[L[div]]] /. x->0 /. y->0;
a4=L[L[L[L[div]]]] /. x->0 /. y->0;
B[q_,m_.]:=1/Product[-(-2j)(q+2j),{j,1,m}];
{c0,c1,c2,c3,c4}-{a0 B[1,0],a1 B[1,1],a2 B[1,2],a3 B[1,3],a4 B[1,4]}
```

```
q=2; R:=Random[Integer,5]; L[f_.]:=D[f,{x,2}]+D[f,{y,2}];
F={P,Q}={Sum[R x^n y^m,{n,0,9},{m,0,9}],Sum[R x^n y^m,{n,0,9},{m,0,9}]}];
div=D[P,x]+D[Q,y];
X=Integrate[r div /. x->r Cos[t] /. y->r Sin[t],{t,0,2Pi},{r,0,s}];
c=Simplify[CoefficientList[Series[X/(s^2 Pi),{s,0,12}],s]];
c0=c[[1]]; c1=c[[3]]; c2=c[[5]]; c3=c[[7]]; c4=c[[9]];
a0=div /. x->0 /. y->0; a1=L[div] /. x->0 /. y->0;
a2=L[L[div]] /. x->0 /. y->0; a3=L[L[L[div]]] /. x->0 /. y->0;
a4=L[L[L[L[div]]]] /. x->0 /. y->0;
B[q_,m_.]:=1/Product[-(-2j)(q+2j),{j,1,m}];
{c0,c1,c2,c3,c4}-{a0 B[2,0],a1 B[2,1],a2 B[2,2],a3 B[2,3],a4 B[2,4]}
```

```
q=3; RR:=Random[Integer,4]; RRR:=Sum[RR x^n y^m z^k,{n,0,6},{m,0,6},{k,0,6}];
L[f_.]:=D[f,{x,2}]+D[f,{y,2}]+D[f,{z,2}]; F={P,Q,R}={RRR,RRR,RRR};
div=D[P,x]+D[Q,y]+D[R,z];
X=Integrate[r^2*Sin[s] div /. x->r*Sin[s]*Cos[t] /. y->r Sin[s] Sin[t] /. z->r Cos[s],
{t,0,2Pi},{s,0,Pi},{r,0,w}];
c=Simplify[CoefficientList[Series[X*3/(4Pi w^3),{w,0,12}],w]];
c0=c[[1]]; c1=c[[3]]; c2=c[[5]]; c3=c[[7]]; c4=c[[9]];
a0=div /. x->0 /. y->0 /. z->0; a1=L[div] /. x->0 /. y->0 /. z->0;
a2=L[L[div]] /. x->0 /. y->0 /. z->0; a3=L[L[L[div]]] /. x->0 /. y->0 /. z->0;
a4=L[L[L[L[div]]]] /. x->0 /. y->0 /. z->0;
B[q_,m_.]:=1/Product[-(-2j)(q+2j),{j,1,m}];
{c0,c1,c2,c3,c4}-{a0 B[3,0],a1 B[3,1],a2 B[3,2],a3 B[3,3],a4 B[3,4]}
```

APPENDIX: ILLUSTRATIONS

7.30. We made our first experiments in this project by computing line integrals on circles on round 2-spheres. This was in the context of teaching multi-variable calculus and exploring new problems for exams. We switch to multi-variable calculus language: take an arbitrary vector field $F = [P, Q, R]$ and compute the line integrals $\int_C F dr$ along geodesic spheres C of geodesic radius r on the 2-sphere. Then integrate the result over M , which for polynomial F is an exercise in double integrals, alas a bit too complicated in general to do by hand. We noticed first experimentally that the result was zero, then explained it using cancellation.




FIGURE 1. A non-local exterior derivative of a 1-form f on a 2-manifold evaluates a line integral of f along a wave front. By Green's theorem applied to a coordinate patch, it is an integral of curl.

7.31. In the Riemannian geometry case the metric for which the metric lacks symmetry, the computation of the wave fronts requires to solve the geodesic differential equations $\ddot{x} + \Gamma_{ij}^k \dot{x}^i \dot{x}^j = 0$. The length of the wave front $W_t(p)$ starting at $p \in M$ is

$$|W_t(p)| = \int_0^{2\pi} |J(t, \theta)| d\theta ,$$

where $J(t, \theta)$ is the **Jacobi field** along the geodesic $x(t)$ starting at p in direction θ . Each $J(t, \theta)$ satisfies the Jacobi differential equations

$$J_{tt}(t, \theta) + K(x(t))J(t, \theta) = 0, J(0, \theta) = 0, J_t(0, \theta) = 1 ,$$

where $K(x(t))$ is the curvature. See [3]. In regions of negative curvature, the wave front length expands fast.

7.32. Let M be a compact 2-manifold with boundary. Given a 1-form f , define the line integral along $C = W_t(p)$ with p at the boundary to be $\int_C d_r f$, a line integral along a half circle. The line integral part is supported in a neighborhood of the boundary.

7.33. Symmetry shows that the integral $\int_M d_t f = 0$ for any $(q - 1)$ form f : every point p and for almost all initial direction θ has for a given t a wave front point $x = W_t(p, \theta)$ a unique geodesic. Continue the geodesic for an other time interval t to get a dual point p' and new direction θ' in the coordinate system at p' that reaches x in time t . The points p and p' both have distance t to x . The line integral contributions of $W_t(p, \theta)$ and $W_t(p', \theta')$ cancel.




FIGURE 2. On a general Riemannian 2-manifold we can evaluate a line integral along a wave front $W_t(p)$ and get so a notion of a **discrete derivative** for 1-forms. By Green's theorem, it is a double integral over the ball $B_t(p) = \{x \in M, d(x, p) = t\}$ at least if t is smaller than the radius of injectivity.




FIGURE 3. In a manifold with boundary, the wave front are reflected at the boundary. We still can show that the total integral of df is zero.

APPENDIX: CURVATURE VIA WAVE FRONTS

7.34. We originally slithered into this topic in the context of **Puiseux type formulas** for curvatures in the discrete [7]. The point there had been **not to take limits** but to use second order derivative notions.

7.35. The starting point was the **Bertrand-Diquet-Puiseux formula** $K = \lim_{r \rightarrow 0} \frac{3(2\pi r - |W_r|)}{r^3 \pi}$. It motivates to define

$$K_h(p) = \frac{2|W_h| - |W_{2h}|}{2\pi h^3}.$$

without taking limits on any smooth surface M . We called this the **R2-D2 formula** because the expression compares the length of wave fronts for two radii in dimension two.

7.36. For example, for a sphere of radius R , where $r = R\phi$ and $|S_r| = 2\pi R \sin(r/R)$ the classical Puiseux formula gives $2(2\pi R \sin(r/R) - 2\pi r)$. The R2-D2 formula gives $4\pi R \sin(r/R) - 2\pi R \sin(2r/R) = 2\pi r^3/R^2 - 2\pi r^5/(4R^4) + \dots$ so that $K = 1/R^2$. Similarly, for a hyperbolic plane, where $|W_r| = 2\pi R \sinh(r/R)$, we get $K = -1/R^2$.

7.37. At the boundary of a region, the R2-D2 formula is

$$K = \lim_{r \rightarrow 0} \frac{2|W_r| - |W_{2r}|}{2\pi r^2}.$$

It is enough to verify this for a circular curve C bounding a disc of radius R , where $|W_r| = 2\pi r \arccos(r/(2R)) = 2\pi K r^2 + 7\pi K^3 r^4/12 + \dots$

7.38. While the Puiseux formula refers to a flat situation, where the circle has circumference $2\pi r$, the R2-D2 formula does refer to the "flat" case. Flatness in this space emerged as the property that the length of a circular wave front grows linearly in time. Again, like here, the motivation for such definitions had been to have bounded curvature notions in situations which are not smooth, like polyhedra or fractal regions or even to push it to more general metric spaces. In general, it is necessary to adjust the constants to get Gauss-Bonnet.




FIGURE 4. R2-D2 formula for curvature for 2 manifolds in interior or boundary.




FIGURE 5. R2-D2 formula at boundary of polygon or Koch region.

REFERENCES

- [1] R. Abraham, J.E. Marsden, and T. Ratiu. *Manifolds, Tensor Analysis and Applications*. Applied Mathematical Sciences, 75. Springer Verlag, New York etc., second edition, 1988.
- [2] V.I. Arnold. *Huygens and Barrow, Newton and Hooke*. Birkhäuser, 1990.
- [3] M. Berger. *A Panoramic View of Riemannian Geometry*. Springer, 2003.
- [4] H.L. Cycon, R.G.Froese, W.Kirsch, and B.Simon. *Schrödinger Operators—with Application to Quantum Mechanics and Global Geometry*. Springer-Verlag, 1987.
- [5] L.C. Evans. *Partial Differential equations*, volume 19 of *Graduate Studies in Mathematics*. AMS, second edition, 2010.
- [6] O. Knill. A remark on quantum dynamics. *Helvetica Physica Acta*, 71:233–241, 1998.
- [7] O. Knill. A discrete Gauss-Bonnet type theorem. *Elemente der Mathematik*, 67:1–17, 2012.
- [8] J.S. Ovall. The Laplacian and Mean and Extreme Values. *American Mathematical Monthly*, 121, 2014.
- [9] P. Pizzetti. Sulla media dei valori che una funzione dei punti dello spazio assume alla superficie di una sfera, rendiconti lincci. *Rendiconti Lincci*, 18:309–316, 1909.
- [10] B. Simon. *A comprehensive course in Analysis*. AMS, 2017.
- [11] D. Vicente. Une goutte d’eau dans un bol. *Quadrature*, 117:13–22, 2020.
- [12] G.N. Watson. *A treatise on the theory of Bessel functions*. Cambridge at the University Press, second edition, 1962.
- [13] S. Zeldich. Wave equation on a Riemannian manifold. Mini course notes Northwestern Analysis SNAP, 2017.