

## Midterm, Mathematics 101

### *Solutions*

**Problem 1.** Give an example of set  $A$  such that  $\cap A = \emptyset$  but  $\cap B \neq \emptyset$  for every finite nonempty subset  $B \subset A$

**Solution.** Let  $A = \{C_0, C_1, \dots\}$ , where  $C_n = \{i \in \mathbb{N} : i \neq n\}$ . Then, since  $n \notin C_n$ ,  $\cap A = \emptyset$ . On the other hand, if  $B$  is a finite subset of  $A$ , and  $m = \max\{n : C_n \in B\} + 1$ , then  $m \in \cap B$ , so  $\cap B \neq \emptyset$

**Problem 2.** Give an example of an injective map  $f : \mathbb{R} \rightarrow \mathbb{R}$  which is not surjective.

**Solution.** You can take  $f(x) = e^x$ .

**Problem 3.** Prove that if  $|A| = |B|$  then  $|P(A)| = |P(B)|$ .

**Solution.** Let  $f : A \rightarrow B$  be a bijection. Define  $F : P(A) \rightarrow P(B)$  by  $F(C) = \{f(c), c \in C\}$  for every  $C \in P(A)$ .

If  $F(C_1) = F(C_2)$ , then  $f(C_1) = f(C_2)$ , so, since  $f$  is an injection,  $C_1 = f^{-1}(f(C_1)) = f^{-1}(f(C_2)) = C_2$ . Thus  $F$  is injective.

Now pick any  $D \in P(B)$ . Take  $C = f^{-1}(D)$ . Since  $f$  is surjective,  $f(C) = D$ . Thus  $F$  is surjective.

**Problem 4.** Let  $G$  be a group for which  $g^2 = 1$  for all  $g \in G$ . Prove that  $G$  is abelian.

**Solution.** Let  $g_1, g_2 \in G$ . Then  $(g_1 * g_2)^2 = 1$ . Multiply this equality by  $g_2$  from the right and by  $g_1$  from the left and use associativity and the fact that  $g_1^2 = g_2^2 = 1$ . Then we get  $g_1 * g_2 = g_2 * g_1$ .

**Problem 5.** Give an example of a group  $G$  with subgroups  $H$  and  $K$  such that  $H \cup K$  is not a subgroup of  $G$ .

**Solution.** Take  $G = (\mathbb{C}, +)$ ,  $H = (\mathbb{R}, +)$ ,  $K = (i\mathbb{R}, +)$ . Then  $H$  and  $K$  are subgroups of  $G$ , but  $H \cup K$  is not closed under addition ( $1 \in H \subset H \cup K$ ,  $i \in K \subset H \cup K$ , but  $1 + i \notin H \cup K$ ).