

MATHEMATICS 191, FALL 2004-2005
MATHEMATICAL PROBABILITY
Check List for the Final Exam

1 You can influence the final exam

Provide your input by Monday, Jan. 17!! Send email to bamberg@tiac.net. For each section list (by number) up to 8 topics that you would like to see left off the final exam and up to six topics that you would especially like to see included. Topics with a preponderance of negative votes will be included only in multiple-choice items or dropped entirely, and those preponderance of positive votes will be favored.

Results of last year's tally were as follows:

Positive votes exceed negative votes

Definitions and Simple Theorems 4,5,7,8,11,21,22,24,33,34,38,42,46

Applications 3,5,9,10,14,16,24,29,31

Proofs 4,8,9,10,11,18,27

Negative votes exceed positive votes

Definitions and Simple Theorems

2,3,10,14,15,17,18,19,26,27,28,29,31,37,40,41,47,48

Applications 6,13,15,21,26,28,33,34,35

Proofs 1,2,3,6,7,13,14,15,17,19,21,25,30

2 Definitions and simple theorems

These are appropriate subject matter for multiple-choice or true-false items.

1. Countability - finite Cartesian products, countable unions and arbitrary intersections of countable sets are countable. The collection of all subsets of a countably infinite subset is uncountable.
2. Definition of a σ -field - closure under complement and countable union implies closure under difference and intersection. Example of two σ -fields whose union is not a σ -field.
3. Inclusion-exclusion formula for two events, and related inequalities.
4. Probabilities for increasing and decreasing infinite sequences of events.
5. Definition of conditional probability.
6. Definition of independent events; example that three events can be pairwise independent but not independent.
7. Probability of k successes in n independent identical experiments.

8. Geometric distribution - probability that the first success in a sequence of identical independent experiments occurs on the k th attempt.
9. Definition of random variables - discrete, continuous, and those that are neither.
10. General properties of distribution functions for random variables.
11. Definition of indicator functions.
12. Definition of independence for discrete random variables.
13. Probability mass functions for binomial, Poisson, geometric and negative binomial distributions.
14. Definition of expectation and variance for discrete random variables; examples where these are not defined because the series is not absolutely convergent.
15. Definition of “uncorrelated” for two random variables; example of two random variables that are uncorrelated but not independent.
16. Statement and proof of the “law of the unconscious statistician” for one discrete random variable.
17. Expectation of the sum of random variables.
18. Variance of the sum of independent random variables and of a multiple of a random variable.
19. Definition of covariance and correlation for two random variables.
20. Basic properties of random walks - homogeneity in time and space. Markov property.
21. Random walks - formula for number of paths in n steps from level a to level b .
22. Reflection principle for random walks - use to obtain a formula for the number of paths from level a to level b that do not cross level 0.
23. Ballot theorem, including the equivalent case of the “hitting time theorem:” the fraction of paths from level 0 to level b in N steps that reach level b for the first time at the N th step.
24. Key lemma for arc sine laws. For a symmetric random walk ($p = \frac{1}{2}$) that starts in level 0, the following are all equal:
 - u_{2m} = the probability of being in level 0 after $2m$ steps.
 - u_{2m} = the probability of never revisiting level 0 during $2m$ steps.

- u_{2m} = the probability of never visiting a negative level during $2m$ steps.
- u_{2m} = the probability of never visiting a positive level during $2m$ steps.

25. By Sterling's approximation, for a symmetric random walk,

$$u_{2k} = 2^{-2k} \frac{(2k)!}{(k!)^2}$$

is approximately $\frac{1}{\pi k}$

26. Arc sine law for last return (mass function): after $2n$ steps of a symmetric random walk, the probability that the last visit to level 0 was at step $2k$ is exactly $u_{2k}u_{2n-2k}$ and approximately

$$\frac{1}{\pi \sqrt{k(n-k)}}$$

27. Arc sine law for last return (distribution function): after $2n$ steps of a symmetric random walk, the probability that the last visit to level 0 was before step $2x$ is approximately $\frac{2}{\pi} \arcsin \sqrt{x}$.

28. Arc sine law for sojourn times (mass function): after $2n$ steps of a symmetric random walk, the probability that $2k$ segments were in "positive territory" is exactly $u_{2k}u_{2n-2k}$ and approximately

$$\frac{1}{\pi \sqrt{k(n-k)}}$$

29. Arc sine law for sojourn times (distribution function): after $2n$ steps of a symmetric random walk, the probability that fraction of time spent in "positive territory" is approximately $\frac{2}{\pi} \arcsin \sqrt{x}$.

30. General properties of density functions for continuous random variables.

31. In terms of the density function $f_X(x)$, integrals for expectations $\mathbb{E}(X)$ and $\mathbb{E}(X^2)$

32. "Law of the unconscious statistician" for the expectation of a function $Y = g(X)$ of a continuous random variable.

33. Exponential distribution and its expectation.

34. Density function for normal distribution $N(0, 1)$ - the variance is 1.

35. Density function for normal distribution $N(\mu, \sigma)$ - the mean is μ and the variance is σ .

36. For a pair of random variables, express the distribution function $F_{X,Y}(x, y)$ in terms of the density function $f_{X,Y}(x, y)$ and vice versa.
37. “Law of the unconscious statistician” for calculating the expectation of a random variable $Z = g(X, Y)$.
38. Definition of marginal density and conditional expectation.
39. Definition of independence for two random variables; what this means in terms of the density function.
40. Density function for the sum of two independent random variables X and Y as a convolution of the density functions for X and Y .
41. Bertrand’s paradox: state the condition for an equilateral triangle based on a random chord to fit inside a circle. Describe three reasonable but different interpretations of “random chord,” and for each of them determine the probability that the triangle fits inside the circle.
42. Crofton’s method: the general strategy for turning a geometrical probability problem into a differential equation.
43. Generating functions: definition of $G(s)$ in terms of the probability mass function. Determining a value of the probability mass function by evaluating an appropriate derivative of $G(s)$. Significance of $G(1)$.
44. Generating function for the sum of two or more independent random variables or for the sum of n independent copies of a random variable.
45. Statement of the “weak law of large numbers.”
46. Statement of the central limit theorem, and the recipe that it implies for generating random numbers with a normal distribution.
47. Statement of Markov’s inequality and Chebyshev’s inequality, and use to obtain crude bounds on how fast the probability must fall to zero for large values of a random variable with zero mean.
48. Statement of the “strong law of large numbers” and what makes it “stronger.” than the weak law.

3 Applications

These can generate a wide variety of problems, some of which are simple enough to cast as multiple-choice items.

1. Equations and inequalities that follow by using induction and the inclusion-exclusion formula.

2. Calculating probabilities by using the inclusion-exclusion formula for two or three events.
3. Probabilities for poker hands.
4. Probabilities for bridge - distribution of suits, or distribution of missing cards in a suit between two opposing hands.
5. Conditional probability problems involving two events (the "bearded man" problem).
6. Generalizations of the Monty Hall problem.
7. Problems based on "Bernoulli trials" - a certain number of successes in n independent identical experiments.
8. Conditional probability problems based on the veracity of witnesses (Edgington's controversy, Queen of Sheba)
9. Analysis of Simpson's paradox in terms of conditional probability.
10. Problems based on geometric and negative binomial distributions - what is the probability that it takes exactly n trials to get k successes?
11. "Problem of the points" - what is the probability that m successes occur before n failures?
12. Problems that involve drawing (without replacement) from an urn containing r red balls and b blue balls.
13. Analysis of union, intersection, etc. in terms of indicator functions.
14. Problems involving the "probabilistic method" and the pigeonhole principle.
15. Problems based on the Poisson distribution.
16. Random walk problems, solved by using conditional probability to write down a linear difference equation and then solving the difference equation with appropriate boundary conditions.
17. Apply the key lemma for arc sine theorems to get various results about random walks (involving maximum or minimum level, for example)
18. For random walks with a small even number of steps, get exact results for the probability that the last visit to the origin was at some specific time $2r$.
19. Given the density or distribution function for a random variable X , determine its expectation and variance.

20. Given the density or distribution function for a random variable X and a new random variable $Y = g(X)$, determine the density function and expectation of Y .
21. Given a random variable X with a uniform distribution on $[0,1]$, invent a random variable $Y = g(X)$ that has some specified distribution function or density function.
22. Given a joint density function $f_{X,Y}(x, y)$, determine whether or not X and Y are independent.
23. Given a joint density function $f_{X,Y}(x, y)$, calculate the conditional density function $f_{Y|X}(y, x)$.
24. Given a joint density function $f_{X,Y}(x, y)$ and new random variables U and V defined by formulas $u(x, y)$ and $v(x, y)$, determine the density function $f_{U,V}(u, v)$.
25. Given the density functions for two independent random variables, calculate the density function for their sum.
26. Given a specification of a way to select the “random chord” for Bertrand’s paradox, calculate the joint density function for the polar coordinates of the midpoint of the chord.
27. Solve geometric probability problems involving two points by dividing the region of interest in half and conditioning on whether the two points lie in the same half or in different halves.
28. Solve geometric probability problems where one of the points of interest lies at a randomly chosen point on a line or circular arc by conditioning on the position of that point.
29. Solve geometric probability problems by Crofton’s method, considering first the simpler case where a point of interest lies on the boundary, then setting up a differential equation.
30. Write down the generating function for a discrete random variable that can take a small number of integer values and for the sum of several such variables.
31. Given the generating function for a random variable X , determine the expectation and variance of X .
32. Given a generating function for the sum S of k independent copies of a random variable, expand in a power series to find the probability that $S = n$. For example, what is the probability to throw a total of n with k fair dice?

33. Use generating functions to solve problems involving the sum of random variables with the geometric distribution.
34. Given a random variable X (discrete or continuous), determine what the law of large numbers has to say about the average of many such random variables.
35. Given a random variable X (discrete or continuous), calculate its mean and variance, and construct a rescaled sum of n independent copies of X whose distribution, according to the central limit theorem, will be approximately normal.
36. “Munchkin problems” (the hypergeometric distribution): given n objects selected from N of which b are special, find the probability that k of the n are special.
37. Use Waring’s theorem to relate probabilities of various events that involve the occurrence of subsets of n events (like finding a bust of Larry Summers in your cereal)

4 Proofs

These generally involve multiple steps and a strategy that is not obvious unless you have already seen the proof. However, there are only finitely many of them.

1. Bernstein’s inequality and the law of averages: Suppose that the probability of a head when a coin is tossed is p . Define the random variable S_n to be the number of heads in n tosses. Prove that the probability

$$\mathbb{P}\left(\frac{S_n}{n} \geq p + \epsilon\right) \leq e^{-\frac{1}{4}n\epsilon^2}$$

2. Given the joint distribution function for two discrete random variables X and Y , derive a formula for $\mathbb{P}(X = x, Y = y)$.
3. Given the joint distribution function for two continuous random variables X and Y , derive a formula for $\mathbb{P}(a < x < leqb, c < Y \leq d)$.
4. Calculate the mean and variance for the binomial distribution.
5. Obtain the Poisson distribution as a limiting case of the binomial distribution, and calculate its mean and variance.
6. Use indicator functions or inclusion-exclusion to prove that if n letters are placed randomly into n matching envelopes, the probability that exactly r letters are placed in the correct envelope is

$$\mathbb{P}(X = r) = \frac{1}{r!} \sum_{s=0}^{n-r} \frac{(-1)^s}{s!}$$

7. Prove the Cauchy-Schwarz inequality for the case of two random variables with zero mean, and use it to show that correlation cannot exceed 1 in magnitude.
8. Gambler's ruin: prove that for a symmetric simple random walk, starting in level k with an absorbing barrier at level 0, the probability of reaching level 0 is 1 but the expected time to do so is infinite.
9. Prove the ballot theorem from the reflection principle.
10. Either by using a combinatorial argument based on the ballot theorem or by a geometrical argument establishing a bijection between paths, prove that the number of paths that start in level 0 and are again in level 0 after $2m$ steps is equal to the number of paths that do not return to level 0 during $2m$ steps.
11. By a geometrical argument establishing a bijection between paths, prove that the number of paths that start in level 0 and do not return to level 0 during $2m$ steps is equal to the number of paths that start in level 0 and do not visit any negative level during $2m$ steps.
12. Prove that after $2n$ steps of a symmetric random walk, the probability that the last visit to level 0 was at step $2k$ is exactly $u_{2k}u_{2n-2k}$.
13. Prove that after $2n$ steps of a symmetric random walk, the probability that the last visit to level 0 was before step xn is approximately $\frac{2}{\pi} \arcsin \sqrt{x}$.
14. Prove that the probability f_{2r} that a symmetric random walk that started in level 0 is again in level 0 for the first time after $2r$ steps is $f_{2r} = \frac{1}{2r-1} \frac{(2r)!}{(r!)^2}$. (This was homework problem 3 on set 6 and is also problem 3.10.1 in the text).
15. By conditioning on the time $2r$ of first return to the origin, prove that after $2n$ steps of a symmetric random walk, the probability that $2k$ segments were in "positive territory" is exactly $u_{2k}u_{2n-2k}$.
16. Prove that for a random variable with a density function $f_X(x)$ that is 0 for negative x ,

$$\int_0^\infty \mathbb{P}(X > x) dx = \mathbb{E}(X)$$
 .
17. For the special case of a random variable $Y = g(X)$ where $g(x) > 0$ for all x , prove the law of the unconscious statistician,

$$\mathbb{E}(Y) = \int_0^\infty g(x) f_X(x) dx$$
 .

18. Given the density function for normal distribution

$$N(\mu, \sigma),$$

$$f(x) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(x-\mu)^2}{2\sigma^2}},$$

make a change of variable and invoke properties of the distribution $N(0, 1)$ to show that the mean is μ and the variance is σ^2 .

19. Prove that for the bivariate normal distribution

$$f(x, y) = \frac{1}{2\pi\sqrt{1-\rho^2}} e^{-\frac{(x^2-2\rho xy+y^2)}{2(1-\rho^2)}},$$

$$\text{var}(X) = \text{var}(Y) = 1 \text{ and } \text{cov}(X, Y) = \rho.$$

20. Prove the convolution formula for the density function of the sum of two continuous random variables $Z = X + Y$.
21. Buffon's needle: Consider a fixed needle of length L whose midpoint is at the center of a circle of diameter d . Choose a random diameter of the circle, then construct a chord whose midpoint lies at a point chose with uniform density on the diameter. Prove that the probability that the needle intersects the chord is $\frac{2L}{\pi d}$.
22. By interchanging the order of summation over two indices, prove that the product of the generating functions for two discrete random variables is the generating function for the convolution of their mass functions and is therefore the generating function for their sum.
23. Derive the formula for the variance of a random variable in terms of its generating function.
24. Use generating functions to calculate mean and variance for the Poisson distribution, show that the sum of two independent Poisson random variables is also Poisson.
25. Let p_n be the probability that when n letters are stuffed randomly into n matching envelopes, no letter ends up in its correct envelope. Starting from the difference equation, valid for $n \geq 3$,

$$np_n = p_{n-2} + (n-1)p_{n-1},$$

derive and solve a differential equation to show that

$$G(s) = \sum_{n=1}^{\infty} p_n s^n = \frac{e^{-s}}{1-s} - 1$$

26. Use generating functions to prove Waring's theorem: Given events A_1, A_2, \dots, A_n with

$$S_m = \sum_{i_1 < i_2 < \dots < i_m} \mathbb{P}(A_{i_1} \cap A_{i_2} \dots \cap A_{i_m}),$$

the probability that precisely X of the events occur is

$$\mathbb{P}(X = i) = \sum_{j=i}^n (-1)^{j-i} \frac{j!}{(j-i)!i!} S_j$$

27. (Not covered in 2004-5) Let $f_r(n)$, for $r > 0$, be the probability that a simple random walk (up one level with probability p , down one level with probability $q = 1 - p$) reaches level r for the first time at the n th step. Define the generating function

$$F_r(s) = \sum_{n=r}^{\infty} f_r(n) s^n.$$

Explain why $F_r(s) = [F_1(s)]^r$, and show that

$$F_1(s) = \frac{1 - \sqrt{1 - 4pqs^2}}{2qs}$$

28. Suppose that independent random variables X_i all have the same distribution function, and expectation $\mathbb{E}(X_i) = \mu$. Let $S_n = \sum_{i=1}^n X_i$. Prove that as $n \rightarrow \infty$ the generating function for $\frac{S_n}{n}$ approaches the constant μ .
29. Suppose that independent random variables X_i all have the same distribution function, with expectation 0 and variance $\mathbb{E}(X_i^2) = \sigma^2$. Let $S_n = \sum_{i=1}^n X_i$. Prove that as $n \rightarrow \infty$ the generating function for $\frac{S_n}{n}$ approaches the generating function for the normal distribution $N(0, 1)$.
30. State and prove the strong law of large numbers for the case of a random variable with zero mean for which $\mathbb{E}(X^4)$ is finite.
31. Prove Waring's theorem from inclusion-exclusion.
32. Prove the generalized inclusion-exclusion formula by induction.