

1) We are given  $xy^3 = 80$  and that  $\frac{dy}{dt} = -0.8$  ft/sec when  $y = 10$  ft

And we want  $\frac{dx}{dt}$  when  $y = 10$  ft

Use implicit differentiation:

$$xy^3 = 80$$


$$\Rightarrow \frac{dx}{dt} y^3 + 3y^2 \frac{dy}{dt} x = 0$$

$$\Rightarrow \frac{dx}{dt} = \frac{-3y^2 x \frac{dy}{dt}}{y^3}$$

when  $y = 10$ ,  $x = \frac{80}{10^3} = .08$

$$\Rightarrow \frac{dx}{dt} = \frac{-3(.08)(-.8)}{10} = .019 \text{ ft/s}$$

2)



$$2x + y = p \Rightarrow x = \frac{p-y}{2}$$

$$\text{Area} = A = \frac{1}{2}bh$$

see that by Pythagoras

$$h = \sqrt{x^2 - \left(\frac{y}{2}\right)^2}$$

$$\Rightarrow \text{Area} = \frac{1}{2}y\sqrt{x^2 - \left(\frac{y}{2}\right)^2}$$

now substitute in for  $x$ :

$$\begin{aligned} A(y) &= \frac{1}{2}y\sqrt{\left(\frac{p-y}{2}\right)^2 - \left(\frac{y}{2}\right)^2} \\ &= \frac{1}{2}y\sqrt{\frac{p^2 - 2py + y^2 - y^2}{4}} \end{aligned}$$

$$A(y) = \frac{1}{4}y\sqrt{p^2 - 2py}$$

$$A'(y) = \frac{1}{4}\sqrt{p^2 - 2py} + \frac{1}{4}y \cdot (-2p) \cdot \frac{1}{2\sqrt{p^2 - 2py}}$$

now set this equal to zero and multiply by  $4\sqrt{p^2 - 2py}$ :

$$\Rightarrow p^2 - 2py - py = 0$$

$$\Rightarrow p - 3y = 0$$

$$\Rightarrow y = \frac{1}{3}p$$

Since the endpoints of the interval give  $A = 0$  ft<sup>2</sup> (minima), we see that  $y = x = \frac{1}{3}p$  gives the maximal area.

$$3) f(x) = \frac{x^2 + 5x + 4}{x}$$

First, examine limits:  $\lim_{x \rightarrow \infty} f(x) = +\infty$ ,  $\lim_{x \rightarrow -\infty} f(x) = -\infty$   
 also, we see that we have a vertical asymptote at  $x = 0$ , and  $\lim_{x \rightarrow 0^+} = +\infty$ ,  $\lim_{x \rightarrow 0^-} = -\infty$

Also, see that  $f(x) = \frac{(x+4)(x+1)}{x}$

$\Rightarrow f(x) = 0$  for  $x = -1, -4$ , so we have the only roots at these points.

$$f'(x) = \frac{x(2x+5) - (x^2 + 5x + 4)}{x^2} = \frac{2x^2 + 5x - x^2 - 5x - 4}{x^2} = \frac{x^2 - 4}{x^2}$$

And we see  $f'(x) = 0$  for  $x = \pm 2$

and since  $f'(x) = \frac{x^2 - 4}{x^2} = 1 - \frac{4}{x^2}$

$$\Rightarrow f''(x) = -2 \cdot (-4) \cdot x^{-3} = \frac{8}{x^3}$$


and so  $f''(x) < 0$  for  $x < 0$

and  $f''(x) > 0$  for  $x > 0$

so  $f$  is concave down on  $(-\infty, 0)$

and  $f$  is concave up on  $(0, \infty)$

So now we can graph it!



$$f(x) = x^3 - 3x^2 + 2x$$

$$f'(x) = 3x^2 - 6x + 2 = 0 \quad \text{for } x = \pm 1 \pm \frac{\sqrt{5}}{3}$$

$$f''(x) = 6x - 6 = 1 - \frac{5}{3} = -\frac{2}{3}$$

$$f''(x) = -5 \cdot (-1) \cdot \frac{2}{3} = \frac{10}{3}$$

$f''(x) > 0$  for  $x < 0$   
 $f''(x) < 0$  for  $x > 0$

concave up on  $(-\infty, 0)$   
 concave down on  $(0, \infty)$

is now we can graph it!

$$\textcircled{4} \quad f(x) = \frac{1}{1 - e^x \sin 5x} \quad ; \quad f'(x) = \frac{-(-e^x \cos 5x \cdot 5)}{(1 - e^x \sin 5x)^2} \cdot \frac{5e^x \cos 5x}{(1 - e^x \sin 5x)}$$

a) linear approximation around zero:

$$f(x) \approx f(0) + x \cdot f'(0)$$

$$f(0) = \frac{1}{1 - 1 \cdot 0} = 1$$

$$f'(0) = \frac{5 \cdot 1 \cdot \cos 0}{(1 - 1 \cdot \sin 0)^2} = 5$$

$$\therefore f(x) \approx 1 + 5x$$

$$\text{b) } f(0.03) \approx 1 + 5 \times 0.03 \\ \approx 1.15$$

$$\textcircled{5} \text{ a) } v = \int_0^t a \, dt + v_0 \quad ; \quad v_0 = 0$$

$$= \int_0^t \frac{t}{10} \, dt = \frac{t^2}{20} \quad \text{for } 0 \leq t \leq 20$$

$$v_{20} = 20$$

$$v = \int_{20}^t a \, dt + v_{20} \quad ; \quad v_{20} = 20$$

$$= \int_{20}^t 2 \, dt + v_{20} = [2t]_{20}^t + 20$$

$$= 2t - 20 \quad \text{for } 20 \leq t \leq 40$$

$$\therefore v(30) = 60 - 20 = 40 \text{ ft/sec}^2$$

$$\text{b) } s = \text{distance}$$

$$s = \int_0^{30} v \, dt = \int_0^{20} \frac{t^2}{20} \, dt + \int_{20}^{30} 2t - 20 \, dt$$

$$= \left[ \frac{t^3}{60} \right]_0^{20} + \left[ t^2 - 20t \right]_{20}^{30}$$

$$= \frac{8000}{60} + 900 - 600 + 400 - 400$$

$$= 433 \text{ m.}$$

$$\begin{aligned}
 \textcircled{6} \text{ a)} \quad M &= \int_0^{40} D(t) dt \\
 &= \int_0^{20} D(t) dt + \int_{20}^{40} D(t) dt \\
 &= \int_0^{20} \frac{t}{10} dt + \int_{20}^{40} 2 dt \\
 &= \left[ \frac{t^2}{20} \right]_0^{20} + \left[ 2t \right]_{20}^{40} \\
 &= \frac{400}{20} + 2 \cdot 40 + (-2) \cdot 20
 \end{aligned}$$

$$M = 60 \text{ kg}$$

b)  $d$  = average density ;  $T$  = total length

$$d = \frac{M}{T} = \frac{60}{40} = 1.5 \text{ kg/m}$$

7.) (A) May 2000  
 $y = (\tan^{-1} x)^x \rightarrow (e^{\ln(\tan^{-1} x)})^x = e^{x \ln(\tan^{-1} x)}$

derivative =  $e^{x \ln(\tan^{-1} x)} \cdot \left( \ln(\tan^{-1} x) + (x) \frac{1}{(\tan^{-1} x)} \cdot \frac{1}{1+x^2} \right)$

=  $(\tan^{-1} x)^x \left( \ln(\tan^{-1} x) + \frac{x}{\tan^{-1} x (1+x^2)} \right)$

(B)

$y = e^{6x} + 6x^e + 6^x$

$\frac{\partial y}{\partial x} = 6e^{6x} + 6e x^{e-1} + 6^x \ln 6$

(C)

$y = \int_{x^3+x}^{100} \sqrt{p^2-p}$

$f(p) = \sqrt{p^2-p}$

=  $F(p) \Big|_{x^3+x}^{100}$

want derivative

derivative

=  $f(p) \Big|_{x^3+x}^{100} = (10000 - 100)^{1/2} - ((x^3+x)^{1/2} - (x^3+x))^{1/2}$

(D)

$x^3 y^3 + xy = 10$

$\partial_y (3y^2 x^3 + x) = -\partial_x (3x^2 + y^3)$   
 $\frac{\partial y}{\partial x} = \frac{-(3x^2 + y^3 + y)}{3y^2 x^3 + x}$

$3x^2 y^3 \partial x + 3y^2 x^3 \partial y + y \partial x + x \partial y = 0$  at (2,1)

$\partial_x (3x^2 y^3 + y) + \partial_y (3y^2 x^3 + x) = 0$   $\frac{\partial y}{\partial x} = \frac{-14}{26}$

May 2000

$$8.) \textcircled{A} \int \frac{4x}{(9x^2+25)^{1/2}} dx$$

$$u = 9x^2 + 25 \quad \rightarrow \quad dx = \frac{du}{18x}$$

$$du = 18x dx$$

$$\int \frac{4x}{u^{1/2}} \frac{du}{18x} = \frac{4}{18} \int u^{1/2} du = \frac{4}{18} (2(9x^2+25)^{1/2})$$

$$\textcircled{B} \int \frac{4}{(9x^2+25)^{1/2}} dx$$

$$u = 3x \quad du = 3dx \quad dx = \frac{du}{3}$$

$$\frac{4}{3} \int \frac{du}{(u^2+5^2)^{1/2}} = \frac{4}{3} \ln(u + (5^2+u^2)^{1/2}) + C$$

$$= \frac{4}{3} \ln(3x + (25+9x^2)^{1/2}) + C$$

$$\textcircled{C} \int_5^{21} \frac{x+3}{(3x+1)^{1/2}} dx$$

$$u = 3x+1 \quad x = \frac{u-1}{3}$$

$$du = 3dx \quad dx = \frac{du}{3}$$

$$\int \frac{(\frac{u-1}{3} + 3) \frac{du}{3}}{u^{1/2}} = \int \left( \frac{u-1}{9} + 1 \right) \frac{du}{u^{1/2}}$$

$$\int \frac{u-1}{9u^{1/2}} + \int \frac{1}{u^{1/2}} = \frac{1}{9} \int u^{1/2} - \int \frac{1}{u^{1/2}} + \int \frac{1}{u^{1/2}}$$

$$= \frac{1}{9} \int u^{1/2} \rightarrow \frac{1}{9} \left( \frac{2}{3} \right) (3x+1)^{3/2} \Big|_5^{21} = 3.5$$

May 2000

$$\textcircled{D} \int_3^{\infty} \frac{\partial x}{(5x+1)^{3/2}} \quad u = 5x+1 \quad \partial x = \frac{\partial u}{5}$$

$$\partial u = 5 \partial x$$

$$\frac{1}{5} \int \frac{\partial u}{u^{3/2}} \rightarrow \frac{1}{5} (-2) u^{-1/2} \Big| = -\frac{2}{5} \left( \frac{1}{(5x+1)^{1/2}} \right) \Big|_3^{\infty}$$

$$= 0 + \frac{2}{5\sqrt{16}} = \frac{2}{5\sqrt{16}}$$

$$9.) \quad x_{n+1} = x_n - \frac{f(x)}{f'(x)}$$

$$= x_n - \frac{x^3 - 30x^2 + 200x + 1}{3x^2 - 60x + 200}$$

when  $x_n = 10$   
and is a root of  
 $x^3 - 30x^2 + 200x$

$$x_{n+1} = 10 - \frac{1}{300 - 600 + 200}$$

$$x_{n+1} = 10 - \frac{1}{100} = \boxed{9.99}$$

10.  $\lim_{n \rightarrow \infty} P_0 \left(1 + \frac{.04}{n}\right)^n = 1$  so use l'Hopital's

$$\lim_{n \rightarrow \infty} P_0 \left(1 + \frac{.04}{n}\right)^n = (1+y)^t P_0 \quad t=1$$

$$\lim_{n \rightarrow \infty} \left(1 + \frac{.04}{n}\right)^n = (1+y)$$

$$\lim_{n \rightarrow \infty} n \ln \left(1 + \frac{.04}{n}\right) = \ln(1+y)$$

$$\lim_{n \rightarrow \infty} \ln \left(1 + \frac{.04}{n}\right) = \ln(1+y)$$

$$\lim_{n \rightarrow \infty} \frac{1}{n}$$

$$\lim_{n \rightarrow \infty} \frac{\left(\frac{-1}{n}\right) \frac{1}{1 + \frac{.04}{n}}}{-\frac{1}{n^2}} = \ln(1+y)$$

$$\lim_{n \rightarrow \infty} \frac{.04}{1 + \frac{.04}{n}} = \ln(1+y)$$

$$.04 = \ln(1+y)$$

$$1+y = e^{.04}$$

$$y = e^{.04} - 1$$

$$\text{SO APR} = 1 + e^{.04} - 1 = e^{.04}$$

$$P = W(C - .5) \quad W = W_0 - \frac{\Delta C}{.1} (10000) \quad \text{let } \Delta C = X \quad C = 1 + X$$

$$P = (100,000 - 100,000X)(1 + X - .5) \quad W = 100,000 - X(100,000)$$

Find  $P' = 0$

$$\begin{aligned} \frac{dP}{dx} &= (-100,000)(.5 + X) + (100,000 - 100,000X)(1) \quad \text{product rule} \\ &= -50,000 - 100,000X + 100,000 - 100,000X \\ &= 50,000 - 200,000X = 0 \end{aligned}$$

$$X = \frac{50,000}{200,000} = \frac{1}{4} \Rightarrow \Delta C = \frac{1}{4} = \$1.25 \Rightarrow C = 1 + .25 = \boxed{\$1.25}$$

Check! profit for  $c = 1.15 = \$50,000$ ,

$$\text{profit for } c = 1.25 = .75(75000) = \$56,250$$

12. At some con  $[1, 5]$   $f(c)$  will equal  $\frac{f(5) + f(1)}{2} = \frac{54 + 10}{2} = \frac{104}{2} = \frac{104}{10}$

mean value

$$x^2 + 5x + 4 = \frac{104}{10}x$$

$$10x^2 - 54x + 40 = 0$$

$$15x^2 - 27x + 20 = 0$$

$$x = \frac{27 \pm \sqrt{27^2 - 4(15)(20)}}{2 \cdot 15} = \frac{27 \pm \sqrt{329}}{30}$$

$$54(x-5) = 2(x)$$