

3. (a)  $g(x) = \int_0^x f(t) dt.$

$$g(0) = \int_0^0 f(t) dt = 0$$

$$g(1) = \int_0^1 f(t) dt = 1 \cdot 2 = 2 \text{ [rectangle]},$$

$$g(2) = \int_0^2 f(t) dt = \int_0^1 f(t) dt + \int_1^2 f(t) dt = g(1) + \int_1^2 f(t) dt \\ = 2 + 1 \cdot 2 + \frac{1}{2} \cdot 1 \cdot 2 = 5 \text{ [rectangle plus triangle]},$$


$$g(3) = \int_0^3 f(t) dt = g(2) + \int_2^3 f(t) dt = 5 + \frac{1}{2} \cdot 1 \cdot 4 = 7,$$

$$g(6) = g(3) + \int_3^6 f(t) dt \text{ [the integral is negative since } f \text{ lies under the } x\text{-axis]} \\ = 7 + \left[ -\left(\frac{1}{2} \cdot 2 \cdot 2 + 1 \cdot 2\right) \right] = 7 - 4 = 3$$

(b)  $g$  is increasing on  $(0, 3)$  because as  $x$  increases from 0 to 3, we keep adding more area.

(c)  $g$  has a maximum value when we start subtracting area; that is, at  $x = 3$ .

(d)




4. (a)  $g(-3) = \int_{-3}^{-3} f(t) dt = 0$ ,  $g(3) = \int_{-3}^3 f(t) dt = \int_{-3}^0 f(t) dt + \int_0^3 f(t) dt = 0$  by symmetry, since the area above the  $x$ -axis is the same as the area below the axis.

(b) From the graph, it appears that to the nearest  $\frac{1}{2}$ ,  $g(-2) = \int_{-3}^{-2} f(t) dt \approx 1$ ,  $g(-1) = \int_{-3}^{-1} f(t) dt \approx 3\frac{1}{2}$ , and  $g(0) = \int_{-3}^0 f(t) dt \approx 5\frac{1}{2}$ .

(c)  $g$  is increasing on  $(-3, 0)$  because as  $x$  increases from  $-3$  to  $0$ , we keep adding more area.

(d)  $g$  has a maximum value when we start subtracting area; that is, at  $x = 0$ .

(e)



(f) The graph of  $g'(x)$  is the same as that of  $f(x)$ , as indicated by FTC1.

9.  $f(t) = t^2 \sin t$  and  $g(y) = \int_2^y t^2 \sin t dt$ , so by FTC1,  $g'(y) = f(y) = y^2 \sin y$ .

12. Let  $u = x^2$ . Then  $\frac{du}{dx} = 2x$ . Also,  $\frac{dh}{dx} = \frac{dh}{du} \frac{du}{dx}$ , so

$$h'(x) = \frac{d}{dx} \int_0^{x^2} \sqrt{1+r^3} dr = \frac{d}{du} \int_0^u \sqrt{1+r^3} dr \cdot \frac{du}{dx} = \sqrt{1+u^3}(2x) = 2x \sqrt{1+(x^2)^3} = 2x \sqrt{1+x^6}.$$

15.  $g(x) = \int_{2x}^{3x} \frac{u^2-1}{u^2+1} du = \int_{2x}^0 \frac{u^2-1}{u^2+1} du + \int_0^{3x} \frac{u^2-1}{u^2+1} du = -\int_0^{2x} \frac{u^2-1}{u^2+1} du + \int_0^{3x} \frac{u^2-1}{u^2+1} du \Rightarrow$

$$g'(x) = -\frac{(2x)^2-1}{(2x)^2+1} \cdot \frac{d}{dx}(2x) + \frac{(3x)^2-1}{(3x)^2+1} \cdot \frac{d}{dx}(3x) = -2 \cdot \frac{4x^2-1}{4x^2+1} + 3 \cdot \frac{9x^2-1}{9x^2+1}$$

18. For the curve to be concave upward, we must have  $y'' > 0$ .  $y = \int_0^x \frac{1}{1+t+t^2} dt \Rightarrow y' = \frac{1}{1+x+x^2} \Rightarrow$

$$y'' = \frac{-(1+2x)}{(1+x+x^2)^2}. \text{ For this expression to be positive, we must have } (1+2x) < 0, \text{ since } (1+x+x^2)^2 > 0 \text{ for all } x.$$

$$(1+2x) < 0 \Leftrightarrow x < -\frac{1}{2}. \text{ Thus, the curve is concave upward on } (-\infty, -\frac{1}{2}).$$


19. (a) By FTC1,  $g'(x) = f(x)$ . So  $g'(x) = f(x) = 0$  at  $x = 1, 3, 5, 7$ , and  $9$ .  $g$  has local maxima at  $x = 1$  and  $5$  (since  $f = g'$  changes from positive to negative there) and local minima at  $x = 3$  and  $7$ . There is no local maximum or minimum at  $x = 9$ , since  $f$  is not defined for  $x > 9$ .

- (b) We can see from the graph that  $|\int_0^1 f dt| < |\int_1^3 f dt| < |\int_3^5 f dt| < |\int_5^7 f dt| < |\int_7^9 f dt|$ . So  $g(1) = |\int_0^1 f dt|$ ,

$$g(5) = \int_0^5 f dt = g(1) - |\int_1^3 f dt| + |\int_3^5 f dt|, \text{ and } g(9) = \int_0^9 f dt = g(5) - |\int_5^7 f dt| + |\int_7^9 f dt|. \text{ Thus,}$$

$$g(1) < g(5) < g(9), \text{ and so the absolute maximum of } g(x) \text{ occurs at } x = 9.$$

- (c)  $g$  is concave downward on those intervals where  $g'' < 0$ . But  $g'(x) = f(x)$ , so  $g''(x) = f'(x)$ , which is negative on (approximately)  $(\frac{1}{2}, 2)$ ,  $(4, 6)$  and  $(8, 9)$ . So  $g$  is concave downward on these intervals.



26. (a) If  $x < 0$ , then  $g(x) = \int_0^x f(t) dt = \int_0^x 0 dt = 0$ .


If  $0 \leq x \leq 1$ , then  $g(x) = \int_0^x f(t) dt = \int_0^x t dt = \left[\frac{1}{2}t^2\right]_0^x = \frac{1}{2}x^2$ .

If  $1 < x \leq 2$ , then

$$\begin{aligned} g(x) &= \int_0^x f(t) dt = \int_0^1 f(t) dt + \int_1^x f(t) dt \\ &= g(1) + \int_1^x (2-t) dt = \frac{1}{2}(1)^2 + \left[2t - \frac{1}{2}t^2\right]_1^x \\ &= \frac{1}{2} + \left(2x - \frac{1}{2}x^2\right) - \left(2 - \frac{1}{2}\right) = 2x - \frac{1}{2}x^2 - 1. \end{aligned}$$

If  $x > 2$ , then  $g(x) = \int_0^x f(t) dt = g(2) + \int_2^x 0 dt = 1 + 0 = 1$ . So

$$g(x) = \begin{cases} 0 & \text{if } x < 0 \\ \frac{1}{2}x^2 & \text{if } 0 \leq x \leq 1 \\ 2x - \frac{1}{2}x^2 - 1 & \text{if } 1 < x \leq 2 \\ 1 & \text{if } x > 2 \end{cases}$$



(c)  $f$  is not differentiable at its corners at  $x = 0, 1$ , and  $2$ .  $f$  is differentiable on  $(-\infty, 0)$ ,  $(0, 1)$ ,  $(1, 2)$  and  $(2, \infty)$ .  $g$  is differentiable on  $(-\infty, \infty)$ .

27. Using FTC1, we differentiate both sides of  $6 + \int_a^x \frac{f(t)}{t^2} dt = 2\sqrt{x}$  to get  $\frac{f(x)}{x^2} = 2 \frac{1}{2\sqrt{x}} \Rightarrow f(x) = x^{3/2}$ . To find  $a$ ,

we substitute  $x = a$  in the original equation to obtain  $6 + \int_a^a \frac{f(t)}{t^2} dt = 2\sqrt{a} \Rightarrow 6 + 0 = 2\sqrt{a} \Rightarrow$

$$3 = \sqrt{a} \Rightarrow a = 9.$$