

## 5.4


3. (a)  $g(x) = \int_0^x f(t) dt$ .

$$g(0) = \int_0^0 f(t) dt = 0, g(1) = \int_0^1 f(t) dt = 1 \cdot 2 = 2 \text{ [rectangle],}$$

$$g(2) = \int_0^2 f(t) dt = \int_0^1 f(t) dt + \int_1^2 f(t) dt = g(1) + \int_1^2 f(t) dt \\ = 2 + 1 \cdot 2 + \frac{1}{2} \cdot 1 \cdot 2 = 5 \text{ [rectangle plus triangle],}$$

$$g(3) = \int_0^3 f(t) dt = g(2) + \int_2^3 f(t) dt = 5 + \frac{1}{2} \cdot 1 \cdot 4 = 7,$$

$$g(6) = g(3) + \int_3^6 f(t) dt \text{ [the integral is negative since } f \text{ lies under the } x\text{-axis]} \\ = 7 + \left[ -\left(\frac{1}{2} \cdot 2 \cdot 2 + 1 \cdot 2\right) \right] = 7 - 4 = 3$$



(b)  $g$  is increasing on  $(0, 3)$  because as  $x$  increases from 0 to 3, we keep adding more area.

(c)  $g$  has a maximum value when we start subtracting area; that is, at  $x = 3$ .

4. (a)  $g(-3) = \int_{-3}^{-3} f(t) dt = 0, g(3) = \int_{-3}^3 f(t) dt = \int_{-3}^0 f(t) dt + \int_0^3 f(t) dt = 0$  by symmetry, since the area above the  $x$ -axis is the same as the area below the axis.

(b) From the graph, it appears that to the nearest  $\frac{1}{2}$ ,


$$g(-2) = \int_{-3}^{-2} f(t) dt \approx 1, g(-1) = \int_{-3}^{-1} f(t) dt \approx 3\frac{1}{2}, \text{ and}$$

$$g(0) = \int_{-3}^0 f(t) dt \approx 5\frac{1}{2}.$$

(c)  $g$  is increasing on  $(-3, 0)$  because as  $x$  increases from  $-3$  to 0, we keep adding more area.

(d)  $g$  has a maximum value when we start subtracting area; that is, at  $x = 0$ .

(f) The graph of  $g'(x)$  is the same as that of  $f(x)$ , as indicated by FTC1.



9.  $f(t) = t^2 \sin t$  and  $g(y) = \int_2^y t^2 \sin t dt$ , so by FTC1,  $g'(y) = f(y) = y^2 \sin y$ .

12. Let  $u = x^2$ . Then  $\frac{du}{dx} = 2x$ . Also,  $\frac{dh}{dx} = \frac{dh}{du} \frac{du}{dx}$ , so

$$h'(x) = \frac{d}{dx} \int_0^{x^2} \sqrt{1+r^3} dr = \frac{d}{du} \int_0^u \sqrt{1+r^3} dr \cdot \frac{du}{dx} = \sqrt{1+u^3}(2x) = 2x \sqrt{1+(x^2)^3} = 2x \sqrt{1+x^6}.$$

15.  $g(x) = \int_{2x}^{3x} \frac{u^2-1}{u^2+1} du = \int_{2x}^0 \frac{u^2-1}{u^2+1} du + \int_0^{3x} \frac{u^2-1}{u^2+1} du = -\int_0^{2x} \frac{u^2-1}{u^2+1} du + \int_0^{3x} \frac{u^2-1}{u^2+1} du \Rightarrow$

$$g'(x) = -\frac{(2x)^2-1}{(2x)^2+1} \cdot \frac{d}{dx}(2x) + \frac{(3x)^2-1}{(3x)^2+1} \cdot \frac{d}{dx}(3x) = -2 \cdot \frac{4x^2-1}{4x^2+1} + 3 \cdot \frac{9x^2-1}{9x^2+1}$$


19. (a) By FTC1,  $g'(x) = f(x)$ . So  $g'(x) = f(x) = 0$  at  $x = 1, 3, 5, 7$ , and  $9$ .  $g$  has local maxima at  $x = 1$  and  $5$  (since  $f = g'$  changes from positive to negative there) and local minima at  $x = 3$  and  $7$ . There is no local maximum or minimum at  $x = 9$ , since  $f$  is not defined for  $x > 9$ .

(b) We can see from the graph that  $\left| \int_0^1 f dt \right| < \left| \int_1^3 f dt \right| < \left| \int_3^5 f dt \right| < \left| \int_5^7 f dt \right| < \left| \int_7^9 f dt \right|$ .

So  $g(1) = \int_0^1 f dt$ ,  $g(5) = \int_0^5 f dt = g(1) - \left| \int_1^3 f dt \right| + \left| \int_3^5 f dt \right|$ , and


$g(9) = \int_0^9 f dt = g(5) - \left| \int_5^7 f dt \right| + \left| \int_7^9 f dt \right|$ . Thus,  $g(1) < g(5) < g(9)$ , and so the absolute maximum of  $g(x)$  occurs at  $x = 9$ .

- (c)  $g$  is concave downward on those intervals where  $g'' < 0$ . But  $g'(x) = f(x)$ , so  $g''(x) = f'(x)$ , which is negative on (approximately)  $(\frac{1}{2}, 2)$ ,  $(4, 6)$  and  $(8, 9)$ . So  $g$  is concave downward on these intervals.



22. (a) In Maple, we should start by setting

`si:=int(sin(t)/t,t=0..x);` In Mathematica, the command is `si=Integrate[Sin[t]/t,{t,0,x}]`. Note that both systems recognize this function; Maple calls it `Si(x)` and Mathematica calls it `SinIntegral[x]`. In Maple, the command to generate the graph is `plot(si,x=-4*Pi..4*Pi);` In Mathematica, it is `Plot[si,{x,-4*Pi,4*Pi}]`. In Derive, we load the utility file `EXP_INT` and plot `SI(x)`.



- (b)  $Si(x)$  has local maximum values where  $Si'(x)$  changes from positive to negative, passing through 0. From the Fundamental Theorem we know that  $Si'(x) = \frac{d}{dx} \int_0^x \frac{\sin t}{t} dt = \frac{\sin x}{x}$ , so we must have  $\sin x = 0$  for a maximum, and for  $x > 0$  we must have  $x = (2n - 1)\pi$ ,  $n$  any positive integer, for  $Si'$  to be changing from positive to negative at  $x$ . For  $x < 0$ , we must have  $x = 2n\pi$ ,  $n$  any positive integer, for a maximum, since the denominator of  $Si'(x)$  is negative for  $x < 0$ . Thus, the local maxima occur at  $x = \pi, -2\pi, 3\pi, -4\pi, 5\pi, -6\pi, \dots$

- (c) To find the first inflection point, we solve  $Si''(x) = \frac{\cos x}{x} - \frac{\sin x}{x^2} = 0$ . We can see from the graph that the first inflection point lies somewhere between  $x = 3$  and  $x = 5$ . Using a root finder gives the value  $x \approx 4.4934$ . To find the  $y$ -coordinate of the inflection point, we evaluate  $Si(4.4934) \approx 1.6556$ . So the coordinates of the first inflection point to the right of the origin are about  $(4.4934, 1.6556)$ . Alternatively, we could graph  $Si''(x)$  and estimate the first positive  $x$ -value at which it changes sign.

- (d) It seems from the graph that the function has horizontal asymptotes at  $y \approx 1.5$ , with  $\lim_{x \rightarrow \pm\infty} Si(x) \approx \pm 1.5$  respectively. Using the limit command, we get  $\lim_{x \rightarrow \infty} Si(x) = \frac{\pi}{2}$ . Since  $Si(x)$  is an odd function,  $\lim_{x \rightarrow -\infty} Si(x) = -\frac{\pi}{2}$ . So  $Si(x)$  has the horizontal asymptotes  $y = \pm \frac{\pi}{2}$ .

- (e) We use the `fsolve` command in Maple (or `FindRoot` in Mathematica) to find that the solution is  $x \approx 1.1$ . Or, as in Exercise 21(c), we graph  $y = Si(x)$  and  $y = 1$  on the same screen to see where they intersect.

24. (a) If  $x < 0$ , then  $g(x) = \int_0^x f(t) dt = \int_0^x 0 dt = 0$ .

If  $0 \leq x \leq 1$ , then  $g(x) = \int_0^x f(t) dt = \int_0^x t dt = \left[\frac{1}{2}t^2\right]_0^x = \frac{1}{2}x^2$ .



If  $1 < x \leq 2$ , then

$$\begin{aligned} g(x) &= \int_0^x f(t) dt = \int_0^1 f(t) dt + \int_1^x f(t) dt \\ &= g(1) + \int_1^x (2-t) dt = \frac{1}{2}(1)^2 + \left[2t - \frac{1}{2}t^2\right]_1^x \\ &= \frac{1}{2} + \left(2x - \frac{1}{2}x^2\right) - \left(2 - \frac{1}{2}\right) = 2x - \frac{1}{2}x^2 - 1. \end{aligned}$$

If  $x > 2$ , then  $g(x) = \int_0^x f(t) dt = g(2) + \int_2^x 0 dt = 1 + 0 = 1$ . So

$$g(x) = \begin{cases} 0 & \text{if } x < 0 \\ \frac{1}{2}x^2 & \text{if } 0 \leq x \leq 1 \\ 2x - \frac{1}{2}x^2 - 1 & \text{if } 1 < x \leq 2 \\ 1 & \text{if } x > 2 \end{cases}$$

(b)



(c)  $f$  is not differentiable at its corners at  $x = 0, 1$ , and  $2$ .  $f$  is differentiable on  $(-\infty, 0)$ ,  $(0, 1)$ ,  $(1, 2)$  and  $(2, \infty)$ .  
 $g$  is differentiable on  $(-\infty, \infty)$ .