

1. Show that  $\int_0^3 \sqrt{9-x^2} dx = \frac{1}{4}9\pi$  - in other words, show analytically that the area of a circle of radius 3 is  $9\pi$  by doing the following:

We'd like to eliminate  $\sqrt{9-x^2}$  by making a substitution that makes the integrand a perfect square. We will exploit the trig identity  $\sin^2 t + \cos^2 t = 1$ , or, equivalently,  $9\sin^2 t + 9\cos^2 t = 9$ . We know that  $9 - 9\sin^2 t$  is a perfect square, so we'll use the substitution  $x = 3\sin t$ . Now we need to write the entire integral in terms of  $t$ .

- a) If  $x = 3\sin t$  then what is  $dx$  in terms of  $t$ ?
- b) If  $x = 3\sin t$  then what is  $\sqrt{9-x^2}$  in terms of  $t$ ?
- c) If  $x = 3\sin t$  then what are the new endpoints of integration in terms of  $t$ ?
- d) Write the integral in terms of  $t$ .
- e) Evaluate the integral in (d).
- f) Conclude that the area of a circle of radius 3 is  $9\pi$ .

For the next three problems, evaluate the integrals. Not *all* require trigonometric substitution. First see if straightforward substitution works.

2.  $\int \frac{x}{\sqrt{4+x^2}} dx$
3.  $\int_0^1 x^3 \sqrt{4-x^2} dx$
4.  $\int_0^1 \frac{x^3}{\sqrt{9-x^2}} dx$
5. The integrands in the following integrals have no simple antiderivative. Use Taylor Series to evaluate (a) and to approximate (b) with error less than  $10^{-4}$ .  
 (a)  $\int e^{-x^2} dx$     (b)  $\int_0^5 \sqrt{1+x^3} dx$

6. Show that the area enclosed by the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ , where  $a$  and  $b$  are positive, is given by  $\pi ab$ . (Recall that in a previous homework you've already shown that the area is between  $2ab$  and  $4ab$ .) (This uses a technique learned in another class; it is not a question using the techniques of §5.9.)

(For extra credit try p. 441 #72 or the following problem:

7. a) Consider Simpson's Rule for  $n = 3$  so that  $a_0 = a$ ,  $a_1 = (a+b)/2$ ,  $a_2 = b$  and Simpson's Rule says that

$$\int_a^b f(x) dx \approx \frac{(b-a)}{6} \left[ f(a) + 4f\left(\frac{a+b}{2}\right) + f(b) \right].$$

Use this formula for  $f(x) = 1$ ,  $f(x) = x$ ,  $f(x) = x^2$ , and  $f(x) = x^3$  and check that in each case it gives the exact value of the integral.

- b) Let  $f(x) = e + fx + gx^2 + hx^3$ , where  $e, f, g, h$  are numbers. Without computation, only basing on a), find out whether for such a function Simpson's Rule gives an exact value of the integral.