

Homework for Monday, March 19, Nr. 3,4,6,16*,36,40*

REMINDER. The **determinant** of a **square matrix** A was defined as the sum over all possible products $(-1)^\pi a_{1\pi(1)} \cdots a_{n\pi(n)}$, where $(-1)^\pi$ is the sign of the permutation (pattern).

SIMPLE CASES. The determinant of a diagonal or tridiagonal matrix is the product of its diagonal elements. The determinant of a block matrix

$$\begin{bmatrix} A & 0 \\ 0 & B \end{bmatrix}$$

is the product $\det(A)\det(B)$.

LINEARITY OF THE DETERMINANT. If the columns of A and B are the same except for the i 'th column, then

$$\det([v_1, \dots, v, \dots, v_n]) + \det([v_1, \dots, w, \dots, v_n]) = \det([v_1, \dots, v+w, \dots, v_n])$$

$$\det([v_1, \dots, kv, \dots, v_n]) = k \det([v_1, \dots, v, \dots, v_n]) .$$

The same holds for rows.

PROPERTIES OF DETERMINANTS.

$$\det(AB) = \det(A)\det(B) .$$

$$\det(A^{-1}) = \det(A)^{-1}$$

$$\det(SAS^{-1}) = \det(A) .$$

$$\det(A^T) = \det(A)$$

If B is obtained from A by switching two rows, then $\det(B) = -\det(A)$. If B is obtained by adding an other row to a given row, then this does not change the value of the determinant.

PROBLEM. Determine $\det(A^{100})$, where A is the matrix $\begin{vmatrix} 1 & 2 \\ 3 & 16 \end{vmatrix}$.

SOLUTION. $\det(A) = 10$, $\det(A^{100}) = (\det(A))^{100} = 10^{100} = 1 \cdot \text{gogool}$. This name as well as the gogoolplex = $10^{10^{100}}$ are official. They are huge numbers: the mass of the universe for example is 10^{52}kg and $1/10^{10^{51}}$ is the chance to find yourself on Mars by quantum fluctuations. (R.E. Crandall, Scient. Amer., Feb. 1997).

ROW REDUCED ECHELON FORM. Determining $\text{rref}(A)$ also determines $\det(A)$.

If A is a matrix and α_i are the factors which are used to scale different rows and s is the number of times, two rows are switched, then $\det(A) = (-1)^s \alpha_1 \cdots \alpha_n \det(\text{rref}(A))$.

PROOF OF $\det(AB) = \det(A)\det(B)$, one brings the $n \times n$ matrix $[A|AB]$ into row reduced echelon form. Similar than the augmented matrix $[A|b]$ was brought into the form $[1|A^{-1}b]$, we end up with $[1|A^{-1}AB] = [1|B]$. By looking at the $n \times n$ matrix to the right during the Gauss elimination, the determinant has changed by $\det(A)$. We end up with a matrix B which has determinant $\det(B)$. Therefore, $\det(AB) = \det(A)\det(B)$.

APPLICATION. If a matrix has a kernel (which is equivalent to be noninvertible), then the determinant is zero.

The reason is that $\det(\text{rref}(A))$ is zero in that case.

THE LAGRANGE EXPANSION.

In order to calculate by hand the determinant of $n \times n$ matrices for $n > 3$, the following expansion is useful. Choose a column i . For each entry a_{ji} in that column, take the $(n-1) \times (n-1)$ matrix A_{ij} called **minor** which does not contain the i 'th column and j 'th row. Then

$$\det(A) = (-1)^{i+1} a_{i1} \det(A_{i1}) + \cdots + (-1)^{i+n} a_{in} \det(A_{in}) = \sum_{j=1}^n (-1)^{i+j} a_{ij} \det(A_{ij}) .$$

VAN DER MONDE DETERMINANT.

If we define for a scalar a the vector $\vec{a} = [1, a, a^2, \dots, a^n]$. For $n + 1$ scalars a_0, \dots, a_n we can consider the matrix with rows \vec{a}_i .

CLAIM: $\det(A) = \prod_{i>j} (a_i - a_j)$. To prove this, we make a Lagrange expansion with respect to the last column. To do so, we call $a_n = x$ and see that the determinant is a polynomial $f(x)$ of degree n in x . It satisfies $f(a_0) = f(a_1) = \dots = f(a_{n-1}) = 0$ because in those cases, the determinant is zero. Therefore $f(x) = k(t - a_1) \dots (t - a_{n-1})$ for some constant k . The number k is the coefficient in front of x^n : it is the van der Monde determinant in the case $n - 1$.

DETERMINANTS IN PHYSICS. Physicists are excited about determinants because summing over all possible "paths" is used as a quantisation method. The Feynman path integral is a "summation" over a suitable set of paths and leads to quantum mechanics. What does it have to do with determinants? Each summand of the determinant can be interpreted like a contribution of a path in a finite graph with n nodes.



The article of Hawking deals with a determinant functional in physics.

This Ray-Singer determinant is a number attached to a geometry resp. an infinite dimensional matrix (the Laplacian) attached to the geometry. Physicists trying to glue quantum mechanics with general relativity hope to make sense of expressions like $\int_g e^{i \det(A(g))}$, where A is an operator attached to some geometry g and where the integral "sums" over all possible geometries.

Commun. math. Phys. 55, 133–148 (1977)

Communications in
Mathematical
Physics
© by Springer-Verlag 1977

Zeta Function Regularization of Path Integrals in Curved Spacetime

S. W. Hawking
Department of Applied Mathematics and Theoretical Physics, University of Cambridge,
Cambridge CB3 9EW, England

Abstract. This paper describes a technique for regularizing quadratic path integrals on a curved background spacetime. One forms a generalized zeta function from the eigenvalues of the differential operator that appears in the action integral. The zeta function is a meromorphic function and its gradient at the origin is defined to be the determinant of the operator. This technique agrees with dimensional regularization where one generalises to n dimensions by adding extra flat dimensions. The generalized zeta function can be expressed as a Mellin transform of the kernel of the heat equation which describes diffusion over the four dimensional spacetime manifold in a fifth dimension of parameter time. Using the asymptotic expansion for the heat kernel, one can deduce the behaviour of the path integral under scale transformations of the background metric. This suggests that there may be a natural cut off in the integral over all black hole background metrics. By functionally differentiating the path integral one obtains an energy momentum tensor which is finite even on the horizon of a black hole. This energy momentum tensor has an anomalous trace.

BRAIN TEASERS.

1) What is the determinant of

$$\begin{bmatrix} 1 & 1 & 0 & 0 & 0 & 0 \\ \pi & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & \pi & 0 & 0 & 0 \\ 0 & 0 & 1 & \pi & 0 & 0 \\ 0 & 0 & 0 & 0 & \pi & 1 \\ 0 & 0 & 0 & 0 & 1 & 1 \end{bmatrix}$$


2) Assume $A = -A^T$. What is $\det(A)$?

3) Assume a A has integer entries and A^{-1} has also integer entries, then $\det(A) = 1$ or $\det(A) = -1$.

4) What values can the determinant of an orthogonal matrix have? (Remember that it satisfies $AA^T = 1$.)

QR DECOMPOSITION. Also the QR decomposition allows to compute the determinant. The determinant of Q is ± 1 , the determinant of R is the product of the diagonal elements of that tridiagonal matrix.

HOW FAST CAN ONE COMPUTE THE DETERMINANT?. Doing the Gauss elimination needs about n^3 steps. The cost to compute the determinant is therefore also of the order n^3 .



The graph to the left shows the time Mathematica needs to calculate the determinant in dependence on the size of the $n \times n$ matrix. The matrix size goes from $n=1$ to $n=300$. The best cubic fit of these data has been obtained by the least square method from chapter IV.