

7/25/2019 SECOND HOURLY PRACTICE V Maths 21a, O.Knill, Summer 2019

Name:

- Start by printing your name in the above box.
- Try to answer each question on the same page as the question is asked. If needed, use the back or the next empty page for work. If you need additional paper, write your name on it.
- Do not detach pages from this exam packet or unstaple the packet.
- Provide details to all computations except for problems 1-3.
- Please write neatly. Answers which are illegible for the grader can not be given credit.
- No notes, books, calculators, computers, or other electronic aids can be allowed.
- You have 90 minutes time to complete your work.

1		20
2		10
3		10
4		10
5		10
6		10
7		10
8		10
9		10
10		10
Total:		110







Problem 1) True/False questions (20 points). No justifications needed.

Mark for each of the 20 questions the correct letter. No justifications are needed.

- 1)  T  F The chain rule tells that  $\frac{d}{dt}f(\vec{r}(t)) = \nabla f(\vec{r}(t)) \cdot \vec{r}'(t)$ .
- 2)  T  F The point  $(1, -1)$  is a critical point of  $f(x, y) = x + y$ .
- 3)  T  F The point  $(0, 1)$  is a critical point of  $f(x, y) = x$  under the constraint  $g(x, y) = x^2 + y^2 = 1$ .
- 4)  T  F The equation  $u_y(x, y) = u_{yy}(x, y)$  is an example of an ordinary differential equation.
- 5)  T  F A point  $(x_0, y_0)$  at which the  $D_{[1,1]^T/2^{1/2}}(x, y)$  is zero, is called a critical point.
- 6)  T  F The relation  $f_{xxyyxx} = f_{xyxyxy}$  holds everywhere for  $f(x, y) = \sin(x^{10} + \cos(xy))$ .
- 7)  T  F The tangent plane to a surface  $z = x^2 + y^2$  at the point  $(1, 1, 2)$  is given by  $2x + 2y = 2$ .
- 8)  T  F Fubini's theorem and Clairot's theorem together imply  $\int_0^1 \int_0^2 f_{xy}(x, y) dydx = \int_0^1 \int_0^2 f_{yx}(x, y) dx dy$ .
- 9)  T  F A Monkey saddle point  $(x_0, y_0)$  of a function  $f(x, y)$  has a negative discriminant  $D$ .
- 10)  T  F  $\int \int_R x^2 + y^2 dx dy$  is the surface area of the paraboloid  $z = x^2 + y^2$  located over the region  $R$  in the  $xy$ -plane.
- 11)  T  F If  $f(0, 0) = 0$  and the discriminant  $D = 0$ , then the linearization  $L(x, y)$  of  $f$  at a point  $(0, 0)$  is constant zero.
- 12)  T  F The directional derivative in the direction of the gradient is negative as it is the direction of steepest decent).
- 13)  T  F If  $\vec{r}(t)$  is a curve on the circle  $g(x, y) = x^2 + y^2 = 1$ , then  $\nabla g(\vec{r}(t)) \cdot \vec{r}'(t) = 0$ .
- 14)  T  F A point  $(0, 0)$ , where the discriminant  $D$  of  $f$  is maximal is a local minimum of  $f$ .
- 15)  T  F If  $f_{xx} > 0$ , then the discriminant is always positive.
- 16)  T  F To maximize  $f(x, y, z)$  under the constraint  $g(x, y, z) = 1$ , we have to solve the Lagrange equations  $\nabla f(x, y, z) = \lambda \nabla g(x, y, z), g(x, y, z) = 1$ .
- 17)  T  F If  $f(x, y) = \sin(x - y)$  then the discriminant  $D$  is zero at every critical point of  $f$ .
- 18)  T  F The gradient vector  $\nabla f(x_0, y_0)$  is a vector which is perpendicular to the normal vector of the surface  $z = f(x, y)$ .
- 19)  T  F If  $|\nabla f(0, 0)| = 10$ , then there is a unit vector  $\vec{v}$  such that  $D_{\vec{v}}f(0, 0) = -11$ .
- 20)  T  F Assume  $f(x, y) = x^2 + y^4$  and a curve  $\vec{r}(t)$  satisfies  $\vec{r}'(t) = \nabla f(\vec{r}(t))$ , then  $\frac{d}{dt}f(\vec{r}(t)) \geq 0$ .

Problem 2) (10 points) No justifications are needed

a) (6 points) Match the regions with the integrals. If no region matches, enter  $O$ .



Enter A-F or O	Integral
	$\int_{-1}^1 \int_{ y }^1 f(x, y) dx dy$
	$\int_{-1}^1 \int_{x^3}^x f(x, y) dy dx$
	$\int_{-1}^1 \int_{-1}^x f(x, y) dy dx$
	$\int_{-1}^1 \int_{-x^4}^{x^4} f(x, y) dy dx$
	$\int_{-1}^1 \int_{\sqrt{ x }}^1 f(x, y) dy dx$
	$\int_{-1}^1 \int_{x^2}^1 f(x, y) dy dx$
	$\int_{-1}^1 \int_{-1}^{x^2} f(x, y) dy dx$
	$\int_{-1}^1 \int_{- x }^{ x } f(x, y) dy dx$

b) (4 points) Name the partial differential equations correctly. Each equation matches one name.

Fill in 1-4	Name
	Wave
	Laplace
	Burgers
	Heat


Equation Number	PDE
1	$f_{\xi\xi} + f_{\eta\eta} = 0$
2	$f_{\eta} + f f_{\eta} - f_{\xi\xi} = 0$
3	$f_{\xi\xi} - f_{\eta\eta} = 0$
4	$f_{\xi} - f_{\eta\eta} = 0$

Problem 3) (10 points) (No justifications are needed.)

a) (5 points) You see a contour map of a function  $f(x, y)$ . Draw the gradient at each of the 5 points A-E. If the gradient should be zero, just mark the point with a bubble.


b) (5 points) Check the boxes which apply. It is in principle possible that more than one box has to be checked in a row or column or that no box needs to be checked in a row or column.

	A	B	C	D	E	F	G	H	I	J	K	L
Local maxima												
Local minima												
Saddle points												
Maximal steepness among A-L												
$f_x = 0, f_y \neq 0$												
$f_y = 0, f_x \neq 0$												
$D_{[1,-1]^T/2^{1/2}}f = 0$												
$D_{[1,1]^T/2^{1/2}}f = 0$												



Problem 4) (10 points)

**Claypit pond** near **Belmont high school** is a nice pond to run around. It has the shape  $g(x, y) = (x - 2)^2 + (y - 3)^2 \leq 1$ . Find the minimal and maximal distance of the “Orchard center” at  $(0, 0)$  to the pond. To do so, we find the maxima and minima of  $f(x, y) = x^2 + y^2$  using Lagrange.



Problem 5) (10 points)


Graph theorists look at the **Tutte polynomial**  $f(x, y)$  of a network. We work with the Tutte polynomial

$$f(x, y) = x + 2x^2 + x^3 + y + 2xy + y^2$$

of the **Kite network**.

a) (4 points) Find the equations for the critical points and check that  $(-2/3, 1/6), (0, -1/2)$  are solutions.

b) (6 points) Classify the two critical using the second derivative test.




**Remark.** The polynomial is useful:  $xf(1 - x, 0)$  tells in how many ways one can color the nodes of the network with  $x$  colors and  $f(1, 1)$  tells how many spanning trees there are. This picture illustrates that the number of spanning trees of the kite graph is  $f(1, 1) = 8$  as you see the 8 possible trees.

Problem 6) (10 points)

At the **Harvard graduate school of design**, a student constructs a wall parametrized by

$$\vec{r}(t, s) = [\sin(t^3), \cos(t^3), ts^2]^T$$

with  $0 \leq t \leq 3$  and  $0 \leq s \leq 1$ . Find its surface area.




**Remark:** The upper figure shows the wall from the problem. Below you see an actual wall design from GSD photographed by Oliver in 2009 near GSD. By the way, the school is close to Memorial Hall. In the garden behind the school near the church, you can find still this interesting wall design.



Problem 7) (10 points)


a) (6 points) Find the linearization  $L(x, y)$  of  $f(x, y) = (x^2 + y)^{1/5}$  at  $(x_0, y_0) = (32, 0)$ .

b) (4 points) Use this to estimate  $(33^2 + 1)^{1/5}$ .



Problem 8) (10 points)

In the science fiction movie **Elysium** of 2013, humans have built a “paradise” for the rich in space. Inspired by this, we imagine an exact copy of the earth glued onto the surface  $x^2 - y^4 + z^2 = 7$ . Find the tangent plane at the point  $(2, 1, 2)$  called “New-Boston”. Our new hyperbolic world is for everybody. And imagine to see the North and South pole on the horizon! Great place to go climbing and skiing.




Problem 9) (10 points)

a) (5 points) We become typographer and design new mathematically defined **typeface** of the alphabet. The new letter "e" in this "21a" design is given by a polar region  $r(t) \leq t^{1/7}$ , with  $0 \leq t \leq 2\pi$ . Find the area of this region.

b) (5 points) Integrate

$$\int_0^1 \int_0^{\arccos(y)} \frac{1}{\cos(x)} dx dy .$$

**Remark:** Computer scientist **Donald Knuth** once wrote an entire article about "The Letter S".



114  
The Letter S  
Donald E. Knuth

SEVERAL YEARS AGO when I began to look at the problem of designing suitable alphabets for use with modern printing equipment, I found that 25 of the letters were comparatively easy to deal with. The other letter was "S". For three days and nights I had a terrible time trying to understand how a proper "S" could really be defined. The solution I finally came up with turned out to involve some interesting mathematics, and I believe that students of calculus and analytic geometry may enjoy looking into the question as I did. The purpose of this paper is to explain what I now consider to be the "right" mathematics underlying printed "S", and also to give an example of the METAFONT language I have recently been developing. (A complete description of METAFONT, which is a computer system and language intended to aid in the design of letter shapes, appears in [1], and [2].)

Before getting into a technical discussion, I should probably mention why I spent so much time on such things in the first place. The central reason is that today's printing technology is essentially based on discrete mathematics and computer science, not on properties of metals or of movable type. The task of making a plate for a printed page is now essentially that of constructing a suitable matrix of 0's and 1's, where the 0's specify white space and the 1's specify ink. I started the second edition of one of my books to look like the first edition, although the first edition had been typeset with the old hot-lead technology, and when I realized that this problem could be solved by using appropriate techniques of discrete mathematics and computer science, I couldn't resist trying to find my own solution.

Reference [2] explains more of the background of my work, and it also discusses the early history of mathematical approaches to type design. In particular, it illustrates how several people proposed to construct "S"

„... con quella tavola quale ho io non potrei de macerare del quadro lungo da la inferiore linea del quadro punto zero. Poi laigo la stessa tavola a... provando una punta dove è stato la inferiore parte del S, qua lo face a dita linea, cioè lungo da la linea del spazio da parte dritta punto 1, a dita

punto 1 da la linea inferiore del quadro. L'altro punto lungo da quella del spazio da parte destra punto 1, a disotto della linea superiore della tavola del quadro sopra la linea da linea. Poi con dita lunghezza de cifra percola l'una punta dove è il primo punto. L'altro punto lungo da la linea del spazio da parte sinistra punto 2, reverso del detto ultimo punto del S, punto che si forma da la inferiore linea del quadro punto 1. Poi da questo ultimo parte in modo comparo a dita linea a comparare con la inferiore tavola lungo da la linea da parte sinistra del quadro punto 1, a parte destra, e qua forma la lettera S, come esperienza a vede.

Fig. 1. Francesco Torricelli's method of "squaring the S" in 1644. (This is page 41 of [2], reproduced by kind permission of Officina Italiana in Vicenza, Italy.)

Problem 10) (10 points)

Compute the weighted surface area

$$\int \int_R (u^2 + v^2) |\vec{r}_u \times \vec{r}_v| du dv$$

of the monkey saddle parametrized by  $\vec{r}(u, v) = [u, v, u^3 - 3uv^2]^T$  over the domain  $R : u^2 + v^2 \leq 1$ . This quantity is also known as the moment of inertia of the surface. Spin that monkey!

