

DERIVED INVARIANCE OF THE NUMBER OF HOLOMORPHIC 1-FORMS AND VECTOR FIELDS

MIHNEA POPA AND CHRISTIAN SCHNELL

1. INTRODUCTION

Given a smooth projective variety X , we denote by $\mathbf{D}(X)$ the bounded derived category of coherent sheaves $\mathbf{D}^b(\mathrm{Coh}(X))$. All varieties we consider below are over the complex numbers. A result of Rouquier, [Ro] Théorème 4.18, asserts that if X and Y are smooth projective varieties with $\mathbf{D}(X) \simeq \mathbf{D}(Y)$ (as linear triangulated categories), then there is an isomorphism of algebraic groups

$$\mathrm{Aut}^0(X) \times \mathrm{Pic}^0(X) \simeq \mathrm{Aut}^0(Y) \times \mathrm{Pic}^0(Y).$$

We refine this by showing that each of the two factors is almost invariant under derived equivalence. According to Chevalley's theorem $\mathrm{Aut}^0(X)$, the connected component of the identity in $\mathrm{Aut}(X)$, has a unique maximal connected affine subgroup $\mathrm{Aff}(\mathrm{Aut}^0(X))$, and the quotient $\mathrm{Alb}(\mathrm{Aut}^0(X))$ by this subgroup is an abelian variety, the Albanese variety of $\mathrm{Aut}^0(X)$. The affine parts $\mathrm{Aff}(\mathrm{Aut}^0(X))$ and $\mathrm{Aff}(\mathrm{Aut}^0(Y))$, being also the affine parts of the two sides in the isomorphism above, are isomorphic. The main result of the paper is

Theorem A. *Let X and Y be smooth projective varieties such that $\mathbf{D}(X) \simeq \mathbf{D}(Y)$. Then*

(1) *$\mathrm{Pic}^0(X)$ and $\mathrm{Pic}^0(Y)$ are isogenous; equivalently, $\mathrm{Alb}(\mathrm{Aut}^0(X))$ and $\mathrm{Alb}(\mathrm{Aut}^0(Y))$ are isogenous.*

(2) *$\mathrm{Pic}^0(X) \simeq \mathrm{Pic}^0(Y)$ unless X and Y are étale locally trivial fibrations over isogenous positive dimensional abelian varieties (hence $\chi(\mathcal{O}_X) = \chi(\mathcal{O}_Y) = 0$).*

The key content is part (1), while (2) simply says that $\mathrm{Aut}^0(X)$ and $\mathrm{Aut}^0(Y)$ are affine unless the geometric condition stated there holds (hence the presence of abelian varieties is essentially the only reason for the failure of the derived invariance of the Picard variety).

Corollary B. *If $\mathbf{D}(X) \simeq \mathbf{D}(Y)$, then*

$$h^0(X, \Omega_X^1) = h^0(Y, \Omega_Y^1) \quad \text{and} \quad h^0(X, T_X) = h^0(Y, T_Y).$$

The Hodge number $h^{1,0}(X) = h^0(X, \Omega_X^1)$ is also called the *irregularity* $q(X)$, the dimension of the Picard and Albanese varieties of X . The invariance of the sum $h^0(X, \Omega_X^1) + h^0(X, T_X)$ was already known, and is a special case of the derived invariance of the Hochschild cohomology of X ([Or], [Ca]; cf. also [Hu] §6.1). Alternatively, it follows from Rouquier's result above. Corollary B, together with the derived invariance of

2000 *Mathematics Subject Classification.* 14F05, 14K30.

Key words and phrases. Derived categories, Picard variety, Hodge numbers.

First author partially supported by NSF grant DMS-0758253 and a Sloan Fellowship.

Hochschild homology (cf. *loc. cit.*), implies the invariance of all Hodge numbers for all derived equivalent threefolds. This was expected to hold as suggested by work of Kontsevich [Ko] (cf. also [BK]).

Corollary C. *Let X and Y be smooth projective threefolds with $\mathbf{D}(X) \simeq \mathbf{D}(Y)$. Then*

$$h^{p,q}(X) = h^{p,q}(Y)$$

for all p and q .

Proof. The fact that the Hochschild homology of X and Y is the same gives

$$(1.1) \quad \sum_{p-q=i} h^{p,q}(X) = \sum_{p-q=i} h^{p,q}(Y).$$

for all i . A straightforward calculation shows that this implies the invariance of all Hodge numbers except for $h^{1,0}$ and $h^{2,1}$, about which we only get that $h^{1,0} + h^{2,1}$ is invariant. We then apply Corollary B. \square

Corollary C is already known (in arbitrary dimension) for varieties of general type: for these derived equivalence implies K -equivalence by a result of Kawamata [Ka], while K -equivalent varieties have the same Hodge numbers according to Batyrev [Ba] and Kontsevich, Denef-Loeser [DL]. It is also well known for Calabi-Yau threefolds; more generally it follows easily for threefolds with numerically trivial canonical bundle (condition which is preserved by derived equivalence, see [Ka] Theorem 1.4). Indeed, since for threefolds Hirzebruch-Riemann-Roch gives $\chi(\omega_X) = \frac{1}{24}c_1(X)c_2(X)$, in this case $\chi(\omega_X) = 0$, hence $h^{1,0}(X)$ can be expressed in terms of Hodge numbers that are known to be derived invariant as above. Finally, in general the invariance of $h^{1,0}$ would follow automatically if X and Y were birational, but derived equivalence does not necessarily imply birationality.

The proof of Theorem A in §3 uses a number of standard facts in the study of derived equivalences: invariance results and techniques due to Orlov and Rouquier, Mukai's description of semi-homogeneous vector bundles, and Orlov's fundamental characterization of derived equivalences. The main new ingredients are results of Nishi-Matsumura and Brion on actions of non-affine algebraic groups (see §2). Further numerical applications of Corollary B to fourfolds or abelian varieties are provided in Remark 3.3.

Finally, the case of abelian varieties shows the existence of Fourier-Mukai partners with non-isomorphic Picard varieties. We expect however the following stronger form of Theorem A(1).

Conjecture. *If $\mathbf{D}(X) \simeq \mathbf{D}(Y)$, then $\mathbf{D}(\text{Pic}^0(X)) \simeq \mathbf{D}(\text{Pic}^0(Y))$.*

Derived equivalent curves must be isomorphic (see e.g. [Hu], Corollary 5.46), while in the case of surfaces the conjecture is checked in the upcoming thesis of Pham [Ph] using the present methods and the classification of Fourier-Mukai equivalences in [BM] and [Ka].

Acknowledgements. We thank A. Căldăraru, L. Ein, D. Huybrechts and M. Mustață for useful comments, and a referee for suggesting improvements to the exposition.

2. ACTIONS OF NON-AFFINE ALGEBRAIC GROUPS

Most of the results in this section can be found in Brion [Br1], [Br2], or are at least implicit there. Let G be a connected algebraic group. According to Chevalley's theorem

(see e.g. [Br1] p.1), G has a unique maximal connected affine subgroup $\text{Aff}(G)$, and the quotient $G/\text{Aff}(G)$ is an abelian variety. We denote this abelian variety by $\text{Alb}(G)$, since the map $G \rightarrow \text{Alb}(G)$ is the Albanese map of G , i.e. the universal morphism to an abelian variety (see [Se2]). Thus $G \rightarrow \text{Alb}(G)$ is a homogeneous fiber bundle with fiber $\text{Aff}(G)$.

Lemma 2.1 ([Br2], Lemma 2.2). *The map $G \rightarrow \text{Alb}(G)$ is locally trivial in the Zariski topology.*

Now let X be a smooth projective variety. We abbreviate $G_X := \text{Aut}^0(X)$, and let $a(X)$ be the dimension of the abelian variety $\text{Alb}(G_X)$. The group G_X naturally acts on the Albanese variety $\text{Alb}(X)$ as well (see [Br1] §3).

Lemma 2.2. *The action of G_X on $\text{Alb}(X)$ induces a map of abelian varieties*

$$\text{Alb}(G_X) \rightarrow \text{Alb}(X),$$

whose image is contained in the Albanese image $\text{alb}_X(X)$. More precisely, the composition $G_X \rightarrow \text{Alb}(X)$ is given by the formula $g \mapsto \text{alb}_X(gx_0 - x_0)$, where $x_0 \in X$ is an arbitrary point.

Proof. From $G_X \times X \rightarrow X$, we obtain a map of abelian varieties

$$\text{Alb}(G_X) \times \text{Alb}(X) \simeq \text{Alb}(G_X \times X) \rightarrow \text{Alb}(X).$$

It is clearly the identity on $\text{Alb}(X)$, and therefore given by a map of abelian varieties $\text{Alb}(G_X) \rightarrow \text{Alb}(X)$. To see what it is, fix a base-point $x_0 \in X$, and write the Albanese map of X in the form $X \rightarrow \text{Alb}(X)$, $x \mapsto \text{alb}_X(x - x_0)$. Let $g \in G_X$ be an automorphism of X . By the universal property of $\text{Alb}(X)$, it induces an automorphism $\tilde{g} \in \text{Aut}^0(\text{Alb}(X))$, making the diagram

$$\begin{array}{ccc} X & \xrightarrow{g} & X \\ \downarrow & & \downarrow \\ \text{Alb}(X) & \xrightarrow{\tilde{g}} & \text{Alb}(X) \end{array}$$

commute; in other words, $\tilde{g}(\text{alb}_X(x - x_0)) = \text{alb}_X(gx - x_0)$. Any such automorphism is translation by an element of $\text{Alb}(X)$, and the formula shows that this element has to be $\text{alb}_X(gx_0 - x_0)$. It follows that the map $G_X \rightarrow \text{Alb}(X)$ is given by $g \mapsto \text{alb}_X(gx_0 - x_0)$. By Chevalley's theorem, it factors through $\text{Alb}(G_X)$. \square

A crucial fact is the following theorem of Nishi and Matsumura (cf. also [Br1]).

Theorem 2.3 ([Ma], Theorem 2). *The map $\text{Alb}(G_X) \rightarrow \text{Alb}(X)$ has finite kernel. More generally, any connected algebraic group G of automorphisms of X acts on $\text{Alb}(X)$ by translations, and the kernel of the induced homomorphism $G \rightarrow \text{Alb}(X)$ is affine.*

Consequently, the image of $\text{Alb}(G_X)$ is an abelian subvariety of $\text{Alb}(X)$ of dimension $a(X)$. This implies the inequality $a(X) \leq q(X)$. Brion observed that X can always be fibered over an abelian variety which is a quotient of $\text{Alb}(G_X)$ of the same dimension $a(X)$; the following proof is taken from [Br1], p.2 and §3, and is included for later use of its ingredients.

Lemma 2.4. *There is an affine subgroup $\text{Aff}(G_X) \subseteq H \subseteq G_X$ with $H/\text{Aff}(G_X)$ finite, such that X admits a G_X -equivariant map $\psi: X \rightarrow G_X/H$. Consequently, X is isomorphic to the equivariant fiber bundle $G_X \times^H Z$ with fiber $Z = \psi^{-1}(0)$.*

Proof. By the Poincaré complete reducibility theorem, the map $\text{Alb}(G_X) \rightarrow \text{Alb}(X)$ splits up to isogeny. This means that we can find a subgroup H containing $\text{Aff}(G_X)$, such that there is a surjective map $\text{Alb}(X) \rightarrow G_X/H$ with $\text{Alb}(G_X) \rightarrow G_X/H$ an isogeny. It follows that $H/\text{Aff}(G_X)$ is finite, and hence that H is an affine subgroup of G_X whose identity component is $\text{Aff}(G_X)$. Let $\psi: X \rightarrow G_X/H$ be the resulting map; it is equivariant by construction. Since G_X acts transitively on G_X/H , we conclude that ψ is an equivariant fiber bundle over G_X/H with fiber $Z = \psi^{-1}(0)$, and therefore isomorphic to

$$G_X \times^H Z = (G_X \times Z)/H,$$

where H acts on the product by $(g, z) \cdot h = (g \cdot h, h^{-1} \cdot z)$. \square

Note that the group H naturally acts on Z ; the proof shows that we obtain X from the principal H -bundle $G_X \rightarrow G_X/H$ by replacing the fiber H by Z (see [Se1], §3.2). While $X \rightarrow G_X/H$ is not necessarily locally trivial, it is so in the étale topology.

Lemma 2.5. *Both $G_X \rightarrow G_X/H$ and $X \rightarrow G_X/H$ are étale locally trivial.*

Proof. Consider the pullback of X along the étale map $\text{Alb}(G_X) \rightarrow G_X/H$,

$$\begin{array}{ccc} X' & \longrightarrow & X \\ \downarrow & & \downarrow \\ \text{Alb}(G_X) & \longrightarrow & G_X/H. \end{array}$$

One notes that $X' \rightarrow \text{Alb}(G_X)$ is associated to the principal bundle $G_X \rightarrow \text{Alb}(G_X)$. The latter is locally trivial in the Zariski topology by Lemma 2.1. \square

Corollary 2.6. *If $a(X) > 0$ (i.e. G_X is not affine), then $\chi(\mathcal{O}_X) = 0$.*

Proof. Clearly $\chi(\mathcal{O}_{X'}) = 0$ since X' is locally isomorphic to the product of Z and $\text{Alb}(G_X)$. But $\chi(\mathcal{O}_{X'}) = \deg(X'/X) \cdot \chi(\mathcal{O}_X)$. \square

3. PROOF OF THE MAIN RESULT

Let $\Phi: \mathbf{D}(X) \rightarrow \mathbf{D}(Y)$ be an exact equivalence between the derived categories of two smooth projective varieties X and Y . By Orlov's criterion, F is uniquely up to isomorphism a Fourier-Mukai functor, i.e. $\Phi \simeq \Phi_{\mathcal{E}}$ with $\mathcal{E} \in \mathbf{D}(X \times Y)$, where $\Phi_{\mathcal{E}}(\cdot) = p_{Y*}(p_X^*(\cdot) \otimes \mathcal{E})$. (Here and in what follows all functors are derived.) A result of Rouquier, [Ro] Théorème 4.18 (see also [Hu], Proposition 9.45), says that Φ induces an isomorphism of algebraic groups¹

$$(3.1) \quad F: \text{Aut}^0(X) \times \text{Pic}^0(X) \simeq \text{Aut}^0(Y) \times \text{Pic}^0(Y)$$

¹Note that in the quoted references the result is stated for the semidirect product of $\text{Pic}^0(X)$ and $\text{Aut}^0(X)$. One can however check that the action of $\text{Aut}^0(X)$ on $\text{Pic}^0(X)$ is trivial. Indeed, $\text{Aut}^0(X)$ acts on $\text{Pic}^0(X)$ by elements in $\text{Aut}^0(\text{Pic}^0(X))$, which are translations. Since the origin is fixed, these must be trivial. This shows in particular that $\text{Aut}^0(X)$ and $\text{Pic}^0(X)$ commute as subgroups of $\text{Aut}(\mathbf{D}(X))$.

in the following manner: A pair of $\varphi \in \text{Aut}(X)$ and $L \in \text{Pic}(X)$ defines an auto-equivalence of $\mathbf{D}(X)$ by the formula $\varphi_*(L \otimes (\cdot))$; its kernel is $(\text{id}, \varphi)_*L \in \mathbf{D}(X \times X)$. When $(\varphi, L) \in \text{Aut}^0(X) \times \text{Pic}^0(X)$, Rouquier proves that the composition $\Phi_{\mathcal{E}} \circ \Phi_{(\text{id}, \varphi)_*L} \circ \Phi_{\mathcal{E}}^{-1}$ is again of the form $\Phi_{(\text{id}, \psi)_*M}$ for a unique pair $(\psi, M) \in \text{Aut}^0(Y) \times \text{Pic}^0(Y)$. We then have $F(\varphi, L) = (\psi, M)$. The following interpretation in terms of the kernel \mathcal{E} was proved by Orlov (see [Or], Corollary 5.1.10) for abelian varieties; the general case is similar, and we include it for the reader's convenience.

Lemma 3.1. *One has $F(\varphi, L) = (\psi, M)$ if and only if*

$$p_1^*L \otimes (\varphi \times \text{id})^*\mathcal{E} \simeq p_2^*M \otimes (\text{id} \times \psi)_*\mathcal{E}.$$

Proof. By construction, $F(\varphi, L) = (\psi, M)$ is equivalent to the relation

$$\Phi_{\mathcal{E}} \circ \Phi_{(\text{id}, \varphi)_*L} = \Phi_{(\text{id}, \psi)_*M} \circ \Phi_{\mathcal{E}}.$$

Since both sides are equivalences, their kernels have to be isomorphic. Mukai's formula for the kernel of the composition of two integral functors (see [Hu], Proposition 5.10) gives

$$(3.2) \quad p_{13*}(p_{12}^*(\text{id}, \varphi)_*L \otimes p_{23}^*\mathcal{E}) \simeq p_{13*}(p_{12}^*\mathcal{E} \otimes p_{23}^*(\text{id}, \psi)_*M).$$

To compute the left-hand side of (3.2), let $\lambda: X \times Y \rightarrow X \times X \times Y$ be given by $\lambda(x, y) = (x, \varphi(x), y)$, making the following diagram commutative:

$$\begin{array}{ccccc} X \times Y & \xrightarrow{\lambda} & X \times X \times Y & \xrightarrow{p_{13}} & X \times Y \\ p_1 \downarrow & & p_{12} \downarrow & & \\ X & \xrightarrow{(\text{id}, \varphi)} & X \times X & & \end{array}$$

By the base-change formula, $p_{12}^*(\text{id}, \varphi)_*L \simeq \lambda_*p_1^*L$; using the projection formula and the identities $p_{13} \circ \lambda = \text{id}$ and $p_{23} \circ \lambda = \varphi \times \text{id}$, we then have

$$p_{13*}(p_{12}^*(\text{id}, \varphi)_*L \otimes p_{23}^*\mathcal{E}) \simeq p_1^*L \otimes \lambda^*p_{23}^*\mathcal{E} \simeq p_1^*L \otimes (\varphi \times \text{id})^*\mathcal{E}.$$

To compute the right-hand side of (3.2), we similarly define $\mu: X \times Y \rightarrow X \times Y \times Y$ by the formula $\mu(x, y) = (x, y, \psi(y))$, to fit into the diagram

$$\begin{array}{ccccc} X \times Y & \xrightarrow{\mu} & X \times Y \times Y & \xrightarrow{p_{13}} & X \times Y \\ p_2 \downarrow & & p_{23} \downarrow & & \\ Y & \xrightarrow{(\text{id}, \psi)} & Y \times Y & & \end{array}$$

Since $p_{13} \circ \mu = (\text{id} \times \psi)$ and $p_{12} \circ \mu = \text{id}$, the same calculation as above shows that

$$p_{13*}(p_{12}^*\mathcal{E} \otimes p_{23}^*(\text{id}, \psi)_*M) \simeq (\text{id} \times \psi)_*(\mathcal{E} \otimes p_2^*M) \simeq (\text{id} \times \psi)_*\mathcal{E} \otimes p_2^*M,$$

where the last step uses that the action of $\text{Aut}^0(Y)$ on $\text{Pic}^0(Y)$ is trivial, so $(\text{id} \times \psi)^*p_2^*M \simeq p_2^*M$. \square

We now give the proof of Theorem A. It is in fact more convenient to start directly with the numerical Corollary B. Note that Rouquier's result (or the invariance of the first Hochschild cohomology) implies the derived invariance of the quantity $h^0(X, \Omega_X^1) + h^0(X, T_X)$. Hence it suffices to show that $q(X) = q(Y)$, where we set $q(X) = h^0(X, \Omega_X^1)$, and similarly for Y .

We continue to write $G_X = \text{Aut}^0(X)$ and $G_Y = \text{Aut}^0(Y)$. Let \mathcal{E} be the kernel defining the equivalence, and let $F: G_X \times \text{Pic}^0(X) \rightarrow G_Y \times \text{Pic}^0(Y)$ be the isomorphism of

algebraic groups from Rouquier's theorem, as above. To prove the assertion, we consider the map

$$\beta: \text{Pic}^0(X) \rightarrow G_Y, \quad \beta(L) = p_1(F(\text{id}, L)),$$

and let $B = \text{Im } \beta$. Similarly, we define

$$\alpha: \text{Pic}^0(Y) \rightarrow G_X, \quad \alpha(M) = p_1(F^{-1}(\text{id}, M)),$$

and let $A = \text{Im } \alpha$. One easily verifies that F induces an isomorphism

$$F: A \times \text{Pic}^0(X) \rightarrow B \times \text{Pic}^0(Y).$$

If both A and B are trivial, we immediately obtain $\text{Pic}^0(X) \simeq \text{Pic}^0(Y)$. Excluding this case from now on, we let the abelian variety $A \times B$ act on $X \times Y$ by automorphisms. Take a point (x, y) in the support of the kernel \mathcal{E} , and consider the orbit map

$$f: A \times B \rightarrow X \times Y, \quad (\varphi, \psi) \mapsto (\varphi(x), \psi(y)).$$

By Lemma 2.2 and the Nishi-Matsumura Theorem 2.3, the induced map $A \times B \rightarrow \text{Alb}(X) \times \text{Alb}(Y)$ has finite kernel. Consequently, the dual map $f^*: \text{Pic}^0(X) \times \text{Pic}^0(Y) \rightarrow \widehat{A} \times \widehat{B}$ is surjective.

Now let $\mathcal{F} := f^*\mathcal{E} \in \mathbf{D}(A \times B)$; it is nontrivial by our choice of (x, y) . For $F(\varphi, L) = (\psi, M)$, the formula in Lemma 3.1 can be rewritten in the more symmetric form (again using the fact that $\psi^*M \simeq M$):

$$(3.3) \quad (\varphi \times \psi)^*\mathcal{E} \simeq (L^{-1} \boxtimes M) \otimes \mathcal{E}.$$

For $(\varphi, \psi) \in A \times B$, let $t_{(\varphi, \psi)} \in \text{Aut}^0(A \times B)$ denote translation by (φ, ψ) . The identity in (3.3) implies that $t_{(\varphi, \psi)}^*\mathcal{F} \simeq f^*(L^{-1} \boxtimes M) \otimes \mathcal{F}$, whenever $F(\varphi, L) = (\psi, M)$. We introduce the map

$$\pi = (\pi_1, \pi_2): A \times \text{Pic}^0(X) \rightarrow (A \times B) \times (\widehat{A} \times \widehat{B}), \quad \pi(\varphi, L) = (\varphi, \psi, L^{-1}|_A, M|_B),$$

where we write $L^{-1}|_A$ for the pull-back from $\text{Alb}(X)$ to A , and same for M . We can then write the identity above as

$$(3.4) \quad t_{\pi_1(\varphi, L)}^*\mathcal{F} \simeq \pi_2(\varphi, L) \otimes \mathcal{F}.$$

Since $\pi_1: A \times \text{Pic}^0(X) \rightarrow A \times B$ is surjective, it follows that each cohomology object $H^i(\mathcal{F})$ is a semi-homogeneous vector bundle on $A \times B$, and that $\dim(\text{Im } \pi) \geq \dim A + \dim B$. On the other hand Mukai [Mu], Proposition 5.1, shows that the semi-homogeneity of $H^i(\mathcal{F})$ is equivalent to the fact that the closed subset

$$\Phi(H^i(\mathcal{F})) := \{(x, \alpha) \in (A \times B) \times (\widehat{A} \times \widehat{B}) \mid t_x^*H^i(\mathcal{F}) \simeq H^i(\mathcal{F}) \otimes \alpha\}$$

has dimension precisely $\dim A + \dim B$. This implies that $\dim(\text{Im } \pi) = \dim A + \dim B$ (and in fact that $\text{Im } \pi = \Phi^0(H^i(\mathcal{F}))$, the neutral component, for any i , though we will not use this; note that Φ is denoted Φ^0 , and Φ^0 is denoted Φ^{00} in [Mu]). Furthermore, we have

$$\begin{aligned} \text{Ker}(\pi) &= \{(\text{id}, L) \in A \times \text{Pic}^0(X) \mid F(\text{id}, L) = (\text{id}, M) \text{ and } L|_A \simeq \mathcal{O}_A \text{ and } M|_B \simeq \mathcal{O}_B\} \\ &\subseteq \{L \in \text{Pic}^0(X) \mid L|_A \simeq \mathcal{O}_A\} = \text{Ker}(\text{Pic}^0(X) \rightarrow \widehat{A}). \end{aligned}$$

Now the surjectivity of f^* implies in particular that the restriction map $\text{Pic}^0(X) \rightarrow \widehat{A}$ is surjective, so we get $\dim(\text{Ker } \pi) \leq q(X) - \dim A$, and therefore

$$\dim A + \dim B = \dim A + q(X) - \dim(\text{Ker } \pi) \geq 2 \dim A.$$

Thus $\dim A \leq \dim B$; by symmetry, $\dim A = \dim B$, and finally, $q(X) = q(Y)$. This concludes the proof of the fact that $\text{Pic}^0(X)$ and $\text{Pic}^0(Y)$ have the same dimension.

We now use this to show that they are in fact isogenous. Let $d = \dim A = \dim B$. The reasoning above proves that $\text{Im } \pi$ is an abelian subvariety of $(A \times B) \times (\widehat{A} \times \widehat{B})$, with $\dim(\text{Im } \pi) = 2d$. For dimension reasons, we also have

$$(3.5) \quad (\text{Ker } \pi)^0 \simeq (\text{Ker}(\text{Pic}^0(X) \rightarrow \widehat{A}))^0 \simeq (\text{Ker}(\text{Pic}^0(Y) \rightarrow \widehat{B}))^0,$$

where the superscripts indicate neutral components. We claim that the projection $p: \text{Im } \pi \rightarrow A \times \widehat{A}$ is an isogeny (likewise for $B \times \widehat{B}$). Indeed, a point in $p^{-1}(\text{id}, \mathcal{O}_A)$ is of the form $(\text{id}, \psi, \mathcal{O}_A, M|_B)$, where $F(\text{id}, L) = (\psi, M)$ and $L|_A \simeq \mathcal{O}_A$. By (3.5), a fixed multiple of (id, L) belongs to $\text{Ker } \pi$, and so $\text{Ker } p$ is a finite set. It follows that $\text{Im } \pi$ is isogenous to both $A \times \widehat{A}$ and $B \times \widehat{B}$; consequently, A and B are themselves isogenous.

To conclude the proof of part (1), note that we have extensions

$$0 \rightarrow \text{Ker } \beta \rightarrow \text{Pic}^0(X) \rightarrow B \rightarrow 0 \quad \text{and} \quad 0 \rightarrow \text{Ker } \alpha \rightarrow \text{Pic}^0(Y) \rightarrow A \rightarrow 0.$$

By definition, $\text{Ker } \beta$ consists of those $L \in \text{Pic}^0(X)$ for which $F(\text{id}, L) = (\text{id}, M)$; obviously, F now induces an isomorphism $\text{Ker } \beta \simeq \text{Ker } \alpha$, and therefore $\text{Pic}^0(X)$ and $\text{Pic}^0(Y)$ are isogenous. Now by Rouquier's isomorphism (3.1) and the uniqueness of $\text{Aff}(G)$ in Chevalley's theorem we have $\text{Aff}(G_X) \simeq \text{Aff}(G_Y)$ and

$$\text{Alb}(G_X) \times \text{Pic}^0(X) \simeq \text{Alb}(G_Y) \times \text{Pic}^0(Y).$$

Therefore we also have equivalently that $\text{Alb}(G_X)$ and $\text{Alb}(G_Y)$ are isogenous.

It remains to check part (2). Clearly $a(X) = a(Y)$. If $a(X) = 0$, we obviously have $\text{Pic}^0(X) \simeq \text{Pic}^0(Y)$. On the other hand, if $a(X) > 0$, Lemmas 2.4 and 2.5 show that X can be written as an étale locally trivial fiber bundle over a quotient of $\text{Alb}(G_X)$ by a finite subgroup, so an abelian variety isogenous to $\text{Alb}(G_X)$. The same holds for Y by symmetry. Note that in this case we have $\chi(\mathcal{O}_X) = \chi(\mathcal{O}_Y) = 0$ by Corollary 2.6.

Remark 3.2. Results of Mukai [Mu], §5 and §6, imply that each $H^i(\mathcal{F})$ on $A \times B$ in the proof above has a filtration with simple semi-homogeneous quotients, all of the same slope, associated to the subvariety $\text{Im } \pi$. In line with Orlov's work on derived equivalences of abelian varieties [Or] §5, one may guess that these simple bundles induce derived equivalences between A and B , and that $\text{Im } \pi$ induces an isomorphism between $A \times \widehat{A}$ and $B \times \widehat{B}$, but we have not been able to prove this.

Remark 3.3 (Further numerical applications). In the case of fourfolds, in addition to the Hodge numbers that are equal due to the general invariance of Hochschild homology (namely $h^{3,0}$ and $h^{4,0}$), Corollary B implies:

Corollary 3.4. *Let X and Y be smooth projective fourfolds with $\mathbf{D}(X) \simeq \mathbf{D}(Y)$. Then $h^{2,1}(X) = h^{2,1}(Y)$. If in addition $\text{Aut}^0(X)$ is not affine, then $h^{2,0}(X) = h^{2,0}(Y)$ and $h^{3,1}(X) = h^{3,1}(Y)$.*

Proof. The analogue of (1.1) for fourfolds implies that $h^{2,1}$ is invariant if and only if $h^{1,0}$ is invariant, and $h^{2,0}$ is invariant if and only if $h^{3,1}$ is invariant. On the other hand, if $\text{Aut}^0(X)$ is not affine, then $\chi(\mathcal{O}_X) = 0$ (cf. Lemma 2.6), which implies that $h^{2,0}$ is invariant if and only if $h^{1,0}$ is invariant. We apply Corollary B. \square

It is also worth noting that Corollary B can help in verifying the invariance of classification properties characterized numerically. We exemplify with a quick proof of the following statement ([HN] Proposition 3.1): *If $\mathbf{D}(X) \simeq \mathbf{D}(Y)$, and X is an abelian variety, then so is Y .* Indeed, the derived invariance of the pluricanonical series [Or] Corollary 2.1.9 and Theorem A imply that $P_1(Y) = P_2(Y) = 1$ and $q(Y) = \dim Y$. The main result of [CH] implies that Y is birational, so it actually has a birational morphism, to an abelian variety B . But $\omega_X \simeq \mathcal{O}_X$, so $\omega_Y \simeq \mathcal{O}_Y$ as well (see e.g. [Hu] Proposition 4.1), and therefore $Y \simeq B$.

REFERENCES

- [BK] S. Barannikov and M. Kontsevich, Frobenius manifolds and formality of Lie algebras of polyvector fields, *Internat. Math. Res. Notices* no. 4 (1998), 201–215.
- [Ba] V. Batyrev, Stringy Hodge numbers of varieties with Gorenstein canonical singularities, *Integrable systems and algebraic geometry (Kobe/Kyoto, 1997)*, World Sci. Publ., River Edge, NJ (1998), 1–32.
- [BM] T. Bridgeland and A. Maciocia, Complex surfaces with equivalent derived categories, *Math. Z.* **236** (2001), 677–697.
- [Br1] M. Brion, Some basic results on actions of non-affine algebraic groups, preprint arXiv:math/0702518.
- [Br2] M. Brion, On the geometry of algebraic groups and homogeneous spaces, preprint arXiv:math/09095014.
- [Ca] A. Căldăraru, The Mukai pairing, I: The Hochschild structure, preprint arXiv:math/0308079.
- [CH] J. A. Chen and Ch. Hacon, Characterization of abelian varieties, *Invent. Math.* **143** (2001), 435–447.
- [DL] J. Denef and F. Loeser, Germs of arcs on singular algebraic varieties and motivic integration, *Invent. Math.* **135** (1999), no. 1, 201–232.
- [Hu] D. Huybrechts, *Fourier-Mukai transforms in algebraic geometry*, Oxford 2006.
- [HN] D. Huybrechts and M. Nieper-Wisskirchen, Remarks on derived equivalences of Ricci-flat manifolds, preprint arXiv:0801.4747, to appear in *Math. Z.*
- [Ka] Y. Kawamata, D -equivalence and K -equivalence, *J. Diff. Geom.* **61** (2002), 147–171.
- [Ko] M. Kontsevich, Homological algebra of mirror symmetry, *Proc. Int. Congr. Math. Zürich 1994*, Birkhäuser (1995), 120–139.
- [Ma] H. Matsumura, On algebraic groups of birational transformations, *Atti della Accademia Nazionale dei Lincei. Rendiconti. Classe di Scienze Fisiche, Matematiche e Naturali. Serie VIII*, **34** (1963), 151–155.
- [Mu] S. Mukai, Semi-homogeneous vector bundles on an abelian variety, *J. Math. Kyoto Univ.* **18** (1978), 239–272.
- [Or] D. Orlov, Derived categories of coherent sheaves and the equivalences between them, *Russian Math. Surveys* **58** (2003), 511–591.
- [Ph] T. Pham, in preparation.
- [Ro] R. Rouquier, Automorphismes, graduations et catégories triangulées, available at <http://people.maths.ox.ac.uk/~rouquier/papers/autograd.pdf>, preprint 2009.
- [Se1] J.-P. Serre, Espaces fibrés algébriques, *Séminaire Claude Chevalley (1958)*, Exposé No. 1, *Documents Mathématiques* **1**, Soc. Math. France, Paris, 2001.
- [Se2] J.-P. Serre, Morphismes universels et variété d’Albanese, *Séminaire Claude Chevalley (1958–1959)*, Exposé No. 10, *Documents Mathématiques* **1**, Soc. Math. France, Paris, 2001.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS AT CHICAGO, 851 S. MORGAN STREET,
CHICAGO, IL 60607, USA

E-mail address: mpopa@math.uic.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS AT CHICAGO, 851 S. MORGAN STREET,
CHICAGO, IL 60607, USA

E-mail address: cschnell@math.uic.edu