

KODAIRA-SAITO VANISHING AND APPLICATIONS

MIHNEA POPA

ABSTRACT. The first part of the paper contains a detailed proof of M. Saito's generalization of the Kodaira vanishing theorem, following the original argument and with ample background. The second part contains some recent applications, and a Kawamata-Viehweg-type statement in the setting of mixed Hodge modules.

1. Introduction. This article was originally the outcome of a lecture delivered at the Clay workshop on mixed Hodge modules, held at Oxford University in August 2013. The main goal was to explain in detail the proof of Morihiko Saito's extension of the Kodaira-Nakano vanishing theorem to mixed Hodge modules, and discuss various special cases and recent applications. This is done in the first and main part of the paper, Sections 2–9, which also includes ample background. Since then I have also included some new applications. One is a proof of weak positivity for the lowest graded piece of a Hodge module obtained jointly with C. Schnell (which also appears in [Sch13]). Another is a Hodge module version of the Kawamata-Viehweg vanishing theorem, likely not in its final form.

M. Saito's vanishing theorem is stated and proved as Theorem 8.2 below. It was obtained in [Sai88, §2.g]; the proof provided here is a detailed account of Saito's original argument, which in turn is a generalization of Ramanujam's topological approach to vanishing. C. Schnell [Sch14a] has recently found a different proof of the theorem, this time extending the Esnault-Viehweg approach to vanishing via the degeneration of the Hodge-to-de Rham spectral sequence on cyclic covers.

In order to make the underlying approach of Saito clear, I will first recall the proof of the Kodaira-Nakano vanishing theorem based on the weak Lefschetz theorem, the Hodge decomposition, and cyclic covering constructions. In the proof of Theorem 8.2, the corresponding roles will be played by the Artin-Grothendieck vanishing theorem for constructible sheaves and by M. Saito's generalization of the standard results of Hodge theory to the setting of mixed Hodge modules. There are however significant new difficulties that are resolved with the use of the interaction between the Hodge filtration and the Kashiwara-Malgrange V -filtration established in [Sai88], recalled in the preliminaries; the background discussion will survey this and other facts about filtered \mathcal{D} -modules in Hodge theory, with references for all the statements needed in the paper.

Many of the standard vanishing theorems involving ample line bundles are special cases of Saito vanishing. This will be reviewed in Section 9, where I will also

2010 *Mathematics Subject Classification.* 14F17; 14F10, 14D07.

During the preparation of this paper I was supported by the NSF grant DMS-1101323.

mention its use to generic vanishing theory. When passing to big and nef line bundles however, the situation is more complicated. While further understanding is needed about transversality conditions between Hodge modules and stable base loci of linear series in order to obtain the strongest possible statements, in Section 11 I prove a first version of Kawamata-Viehweg for mixed Hodge modules – it essentially assumes that the Hodge module is a variation of mixed Hodge structure over the augmented base locus of a nef and big line bundle. Another application, provided in Section 10, is a proof together with Schnell of an extension of a weak positivity theorem of Viehweg to the lowest graded piece of the Hodge filtration on a Hodge \mathcal{D} -module. Arguing along the lines of Kollár’s approach to weak positivity provides a very quick argument, once Kodaira vanishing and adjunction have been extended to setting of mixed Hodge modules.

As most of the paper is expository, my main goal is to make these very useful statements and techniques more accessible to algebraic geometers; the viewpoint is that of cohomological methods in birational geometry. The reader interested in a more general overview of the theory of mixed Hodge modules is encouraged to consult the recent [Sch14b], besides of course the original [Sai88] and [Sai90].

Acknowledgements. I am very grateful to Christian Schnell, from whom I learned a lot about Hodge modules, and who made numerous useful comments on this paper. I would also like to thank Nero Budur, Mircea Mustața, Claude Sabbah and Morihiko Saito for answering my questions, and the organizers of the Oxford Clay workshop on mixed Hodge modules (all among the above) for putting together such a valuable event.

2. The topological/Hodge theoretic approach to Kodaira vanishing.

In this section I will recall the approach to the Kodaira vanishing theorem based on topological and Hodge theoretic methods, which also gives the more general Nakano vanishing. It was first observed by Ramanujam that one can use such methods, Kodaira’s original proof being of a differential geometric nature. I will follow the treatment in [Laz04] §4.2; this is intended to be an introduction to the strategy used by Saito in order to prove the more general result for Hodge modules.

Theorem 2.1 (Kodaira-Nakano Vanishing Theorem). *Let X be a smooth complex projective variety, and L an ample line bundle on X . Then*

$$H^q(X, \Omega_X^p \otimes L) = 0 \quad \text{for } p + q > n,$$

or equivalently

$$H^q(X, \Omega_X^p \otimes L^{-1}) = 0 \quad \text{for } p + q < n.$$

Before proving the theorem, let’s review some useful technical tools. First, recall the following well-known cyclic covering construction, needed in order to “take m -th roots” of divisors $D \in |mL|$, with L some line bundle. For a proof of this and other covering constructions see [Laz04] §4.1.B.

Proposition 2.2. *Let X be a variety over an algebraically closed field k , and let L be a line bundle on X . Let $0 \neq s \in H^0(X, L^{\otimes m})$ for some $m \geq 1$, with $D = Z(s) \in |mL|$. Then there exists a finite flat morphism $f : Y \rightarrow X$, where Y is*

a scheme over k such that if $L' = f^*L$, there is a section

$$s' \in H^0(Y, L') \text{ satisfying } (s')^m = f^*s.$$

Moreover:

- if X and D are smooth, then so are Y and $D' = Z(s')$.
- the divisor D' maps isomorphically onto D .
- there is a canonical isomorphism $f_*\mathcal{O}_Y \simeq \mathcal{O}_X \oplus L^{-1} \oplus \dots \oplus L^{-(m-1)}$.

Furthermore, recall that if X is a smooth variety, and D is a smooth effective divisor on X , then the sheaf of 1-forms on X with log-poles along D is

$$\Omega_X^1(\log D) = \Omega_X^1 \left\langle \frac{df}{f} \right\rangle, \quad f \text{ local equation for } D.$$

Concretely, if z_1, \dots, z_n are local coordinates on X , chosen such that $D = (z_n = 0)$, then $\Omega_X^1(\log D)$ is locally generated by $dz_1, \dots, dz_{n-1}, \frac{dz_n}{z_n}$. This is a free system of generators, so $\Omega_X^1(\log D)$ is locally free of rank n . For any integer p , we define

$$\Omega_X^p(\log D) := \bigwedge^p (\Omega_X^1(\log D)).$$

Using local calculations and the residue map, it is standard to verify the following statements (see [EV92, §2] or [Laz04, Lemma 4.2.4]):

Lemma 2.3. *There are short exact sequences:*

$$(i) \quad 0 \longrightarrow \Omega_X^p \longrightarrow \Omega_X^p(\log D) \longrightarrow \Omega_D^{p-1} \longrightarrow 0.$$

$$(ii) \quad 0 \longrightarrow \Omega_X^p(\log D)(-D) \longrightarrow \Omega_X^p \longrightarrow \Omega_D^p \longrightarrow 0.$$

Lemma 2.4. *Let $f : Y \rightarrow X$ be the m -fold cyclic cover branched along D , as in Proposition 2.2. Let D' be the divisor in Y such that $f^*D = mD'$, mapping isomorphically onto D . Then*

$$f^*\Omega_X^p(\log D) \simeq \Omega_Y^p(\log D').$$

Sketch of proof of Theorem 2.1. By Serre duality it suffices to show the second part of the statement. For $m \gg 0$, let $D \in |mL|$ be a smooth divisor. One can assume by induction on $n = \dim X$ that we already know Kodaira-Nakano vanishing on D , so that

$$H^q(D, \Omega_D^{p-1} \otimes L_D^{-1}) = 0 \quad \text{for } p+q < n.$$

Using this and passing to cohomology in the sequence in Lemma 2.3(i), it suffices then to prove that

$$H^q(X, \Omega_X^p(\log D) \otimes L^{-1}) = 0 \quad \text{for } p+q < n.$$

Let now $f : Y \rightarrow X$ be the m -fold cyclic cover branched along D as in Proposition 2.2, with $f^*D = mD'$ and $L' = \mathcal{O}_Y(D')$. Proposition 2.2 says that Y and D' can be chosen to be smooth; also, D' is obviously ample. Since f is a finite cover, using Lemma 2.4 what we want is equivalent to showing that

$$H^q(Y, \Omega_Y^p(\log D') \otimes \mathcal{O}_Y(-D')) = 0 \quad \text{for } p+q < n.$$

One can now appeal to the exact sequence in Lemma 2.3(ii). Using this, our desired statement is equivalent to the fact that the restriction maps

$$r_{p,q} : H^q(Y, \Omega_Y^p) \longrightarrow H^q(D', \Omega_{D'}^p)$$

are isomorphisms for $p+q \leq n-2$, and injective for $p+q = n-1$. But this follows immediately from the weak Lefschetz theorem, as the restriction maps

$$H^i(Y, \mathbb{C}) \longrightarrow H^i(D', \mathbb{C})$$

are morphisms of Hodge structures. \square

Saito's generalization of Theorem 2.1 is stated and proved in Section 8, while important special cases are explained in Section 9. Before being able to do this we need a lengthy review of background material. The reader may already visit those sections however, for a first encounter with the main topic.

3. Filtered \mathcal{D} -modules and de Rham complexes. In this section I will recall some filtered \mathcal{D} -module terminology used in the paper; a comprehensive introduction can be found for instance in the lecture notes of Maisonobe-Sabbah [MS]. In what follows the standard language is that of right \mathcal{D} -modules; as emphasized in [Sai88], this is often more appropriate in the theory of mixed Hodge modules, for instance due to the fact that it is the natural setting for considering direct image or duality functors. Occasionally however left \mathcal{D} -modules will be necessary, in which case I will state explicitly that we are considering that setting and are performing the left-right transformation described below.

Let X be a complex manifold or smooth variety. A filtered right \mathcal{D} -module on X is a \mathcal{D}_X -module with an increasing filtration $F = F_\bullet \mathcal{M}$ by coherent \mathcal{O}_X -modules, bounded from below and satisfying

$$F_l \mathcal{M} \cdot F_k \mathcal{D}_X \subseteq F_{k+l} \mathcal{M} \quad \text{for all } k, l \in \mathbb{Z}.$$

In addition, the filtration is *good* if the inclusions above are equalities for $k \gg 0$. This condition is equivalent to the fact that the total associated graded object

$$\mathrm{Gr}_\bullet^F \mathcal{M} = \bigoplus_k \mathrm{Gr}_k^F \mathcal{M} = \bigoplus_k F_k \mathcal{M} / F_{k-1} \mathcal{M}$$

is finitely generated over $\mathrm{Gr}_\bullet^F \mathcal{D}_X \simeq \mathrm{Sym} T_X$, i.e. induces a coherent sheaf on the cotangent bundle T^*X .

As one often needs to switch between the two, let's recall that there is a one-to-one correspondence between left and right \mathcal{D}_X -modules given by

$$\mathcal{N} \mapsto \mathcal{M} = \mathcal{N} \otimes_{\mathcal{O}_X} \omega_X \quad \text{and} \quad \mathcal{M} \mapsto \mathcal{N} = \mathcal{H}om_{\mathcal{O}_X}(\omega_X, \mathcal{M}).$$

In terms of filtrations, the left-right rule is

$$F_p \mathcal{N} = F_{p-n} \mathcal{M} \otimes_{\mathcal{O}_X} \omega_X^{-1}.$$

While we will consider right \mathcal{D}_X -modules when talking about Hodge modules, one naturally associates the de Rham complex to the corresponding left \mathcal{D}_X -module \mathcal{N} :

$$\mathrm{DR}_X(\mathcal{N}) = \left[\mathcal{N} \rightarrow \Omega_X^1 \otimes \mathcal{N} \rightarrow \cdots \rightarrow \Omega_X^n \otimes \mathcal{N} \right],$$

which is a \mathbb{C} -linear complex placed in degrees $0, \dots, n$, with maps induced by the corresponding integrable connection $\nabla : \mathcal{N} \rightarrow \mathcal{N} \otimes \Omega_X^1$. It turns out that the

natural de Rham complex to consider for the right \mathcal{D} -module \mathcal{M} (sometimes called a Spencer complex; see [MS, 1.4.2]) satisfies

$$\mathrm{DR}_X(\mathcal{M}) \simeq \mathrm{DR}_X(\mathcal{N})[n].$$

By definition the filtration $F_\bullet \mathcal{M}$ is compatible with the \mathcal{D}_X -module structure on \mathcal{M} and therefore, using the left-right rule above, this induces a filtration on the de Rham complex of \mathcal{M} by the formula

$$F_k \mathrm{DR}_X(\mathcal{M}) = \left[\bigwedge^n T_X \otimes F_{k-n} \mathcal{M} \rightarrow \bigwedge^{n-1} T_X \otimes F_{k+1-n} \mathcal{M} \rightarrow \cdots \rightarrow F_k \mathcal{M} \right][n].$$

The associated graded complexes for the filtration above are

$$\mathrm{Gr}_k^F \mathrm{DR}_X(\mathcal{M}) = \left[\bigwedge^n T_X \otimes \mathrm{Gr}_{k-n}^F \mathcal{M} \rightarrow \bigwedge^{n-1} T_X \otimes \mathrm{Gr}_{k+1-n}^F \mathcal{M} \rightarrow \cdots \rightarrow \mathrm{Gr}_k^F \mathcal{M} \right][n],$$

which are now complexes of coherent \mathcal{O}_X -modules in degrees $-n, \dots, 0$, and provide objects in $\mathbf{D}^b(X)$, the bounded derived category of coherent sheaves on X .

We will be particularly interested in the lowest graded piece of a filtered \mathcal{D} -module. For one such right \mathcal{D}_X -module (\mathcal{M}, F) define

$$(3.1) \quad p(\mathcal{M}) := \min \{p \mid F_p \mathcal{M} \neq 0\} \quad \text{and} \quad S(\mathcal{M}) := F_{p(\mathcal{M})} \mathcal{M}.$$

For the associated left \mathcal{D}_X -module we then have

$$p(\mathcal{N}) = p(\mathcal{M}) + n \quad \text{and} \quad S(\mathcal{N}) = S(\mathcal{M}) \otimes \omega_X^{-1}.$$

4. Hodge modules and variations of Hodge structure. In this section I will briefly recall the objects which are the main focus of the paper, while in the next section I will give several important examples. The main two references for the theory of Hodge modules are Morihiko Saito's papers [Sai88] in the pure case, and [Sai90] in the mixed case. A quite gentle but comprehensive overview of the theory was recently provided by Schnell [Sch14b]. Here I will only recall the information necessary for understanding Saito's vanishing theorem (but will discuss the necessary facts on V -filtrations in a later section), indicating what the reader should consult in the references above.¹

Let X be a smooth complex algebraic variety of dimension n , and Z an irreducible closed subset. Let \mathbf{V} be a polarizable variation of \mathbb{Q} -Hodge structure of weight k on an open set U in the smooth locus of Z , so in particular

$$\mathbf{V} = (\mathcal{V}, F^\bullet, \mathbf{V}_{\mathbb{Q}})$$

with $\mathbf{V}_{\mathbb{Q}}$ a local system of \mathbb{Q} -vector spaces on U , $\mathcal{V} = \mathbf{V}_{\mathbb{Q}} \otimes_{\mathbb{Q}} \mathcal{O}_U$, and $F^p = F^p \mathcal{V}$ a decreasing filtration of subbundles of \mathcal{V} inducing a Hodge filtration over each $x \in U$ and satisfying Griffiths transversality with respect to the connection associated to \mathcal{V} , meaning that for each p there is an induced

$$\nabla : F^p \longrightarrow F^{p-1} \otimes \Omega_U^1.$$

To this, according to [Sai88] one associates a pure Hodge module of weight $\dim Z + k$ on U , whose main constituents are:

¹Several of the contributions to the Clay workshop were devoted to providing a detailed introduction to the background theory of mixed Hodge modules, and hopefully will eventually lead to further useful material.

- The right \mathcal{D} -module $\mathcal{M} = \mathcal{V} \otimes \omega_U$ with filtration $F_p \mathcal{M} = F^{-p-n} \mathcal{V} \otimes \omega_U$.
- The \mathbb{Q} -perverse sheaf $P = \mathbf{V}_{\mathbb{Q}}[n]$.

In fact, according to Saito's theory, this extends uniquely to a Hodge module of weight $\dim Z + k$ on X , whose support is Z and whose underlying perverse sheaf is the intersection complex $\mathrm{IC}_Z(\mathbf{V}_{\mathbb{Q}})$ associated to the given local system. The filtration on the corresponding \mathcal{D} -module is (nontrivially) determined by the Hodge filtration on U .

More generally, in [Sai88] Saito introduced an abelian category of $\mathrm{HM}(X, \ell)$ of pure polarizable Hodge modules on X of weight ℓ . The main two constituents of one such are still:

- A filtered regular holonomic \mathcal{D}_X -module (\mathcal{M}, F) , where $F = F_{\bullet} \mathcal{M}$ is a good filtration by \mathcal{O}_X -coherent subsheaves, so that $\mathrm{Gr}_{\bullet}^F \mathcal{M}$ is coherent over $\mathrm{Gr}_{\bullet}^F \mathcal{D}_X$.
- A \mathbb{Q} -perverse sheaf P on X whose complexification corresponds to \mathcal{M} via the Riemann-Hilbert correspondence, so that there is an isomorphism

$$\alpha : \mathrm{DR}_X(\mathcal{M}) \xrightarrow{\sim} P \otimes_{\mathbb{Q}} \mathbb{C}.$$

If X is a point, such a Hodge module is simply a polarizable Hodge structure of weight ℓ , while in general the main examples are those constructed as above. Indeed, the existence of polarizations makes the category $\mathrm{HM}(X, \ell)$ semi-simple: each object admits a decomposition by support, and simple objects with support equal to an irreducible subvariety $Z \subseteq X$ (called pure Hodge modules with strict support Z , which means that they have no nontrivial subobjects or quotient objects whose support is Z) are obtained from polarizable variations of Hodge structure on Zariski-open subsets of Z by the procedure described above; this is one of the fundamental results obtained in [Sai88], [Sai90]. Formally,

$$(4.1) \quad \mathrm{HM}(X, \ell) = \bigoplus_{Z \subseteq X} \mathrm{HM}^Z(X, \ell),$$

with $\mathrm{HM}^Z(X, \ell)$ the subcategory of pure Hodge modules of weight ℓ with strict support Z .

A fundamental point of the definition of $\mathrm{HM}_Z(X, \ell)$ is that it is inductive on the dimension of the support, by means of the use of the monodromy filtration on nearby cycles. I will gloss over this as it will not play a key role here; a nice discussion can be found in [Sch14b, §12].

Furthermore, M. Saito introduced in [Sai90] the abelian category $\mathrm{MHM}(X)$ of (graded-polarizable) mixed Hodge modules on X . In addition to data as in (i) and (ii) above, in this case a third main constituent is:

- A finite increasing weight filtration $W_{\bullet} M$ of M by objects of the same kind, compatible with α , such that the graded quotients $\mathrm{Gr}_{\ell}^W M = W_{\ell} M / W_{\ell-1} M$ are pure Hodge modules in $\mathrm{HM}(X, \ell)$.

Again, if X is a point a mixed Hodge module is a graded-polarizable mixed Hodge structure. These components are subject to several conditions, which are defined by induction on the dimension of the support of \mathcal{M} : roughly speaking, it

is required that the nearby and vanishing cycles of \mathcal{M} with respect to any locally defined holomorphic function be again mixed Hodge modules (now on a variety of dimension $n - 1$). For a further discussion of the definition, see also [Sch14b, §20].

One of the most important results of the theory is M. Saito's theorem on the behavior of direct images of polarizable Hodge modules via projective morphisms. (I am only stating part of it here.)

Theorem 4.2 (Stability Theorem, [Sai88, Théorème 5.3.1]). *Let $f : X \rightarrow Y$ be a projective morphism of smooth complex varieties, and $h = c_1(H)$ for a line bundle H on X which is ample relative to f . If $M \in \text{HM}(X, \ell)$ is a polarizable Hodge module, then*

(i) *The filtered direct image $f_*(\mathcal{M}, F)$ is strict, and $\mathcal{H}^i f_* \mathcal{M}$ underlies a polarizable Hodge module $M_i \in \text{HM}(Y, \ell + i)$.*

(ii) *For every i one has an isomorphism of pure Hodge modules*

$$h^i : M_{-i} \longrightarrow M_i(i).$$

Recall that a complex of filtered \mathcal{D} -modules is strict if its morphisms are strictly compatible with the filtrations. The statement in (i) is a key property of \mathcal{D} -modules underlying Hodge modules that is not shared by arbitrary filtered \mathcal{D} -modules; for more on this see e.g. [Sch14b, §26-28]. One of its important consequence is the commutation of the graded quotients of the de Rham complex with direct images:

$$\mathbf{R}f_* \text{Gr}_k^F \text{DR}_X(\mathcal{M}) \simeq \text{Gr}_k^F \text{DR}_Y(f_* \mathcal{M}).$$

A fundamental consequence of the theorem above deduced in [Sai88] is the analogue of the decomposition theorem for pure polarizable Hodge modules, obtained formally from the above as an application of Deligne's criterion for the degeneration of the Leray spectral sequence in terms of the Lefschetz operator. The result is often stated for the underlying perverse sheaves, extending the well-known BBD-decomposition theorem; here I state the filtered \mathcal{D} -modules version, which is crucial for the applications presented later.

Theorem 4.3 (Saito Decomposition Theorem). *Let $f : X \rightarrow Y$ be a projective morphism of smooth complex varieties, and let $M \in \text{HM}(X, \ell)$, with underlying filtered \mathcal{D} -module (\mathcal{M}, F) . Then*

$$f_*(\mathcal{M}, F) \simeq \bigoplus_{i \in \mathbb{Z}} \mathcal{H}^i f_*(\mathcal{M}, F)[-i]$$

in the derived category of filtered \mathcal{D}_Y -modules.

Remark 4.4. As we are working in the algebraic category and all mixed Hodge modules will be polarizable (cf. [Sai90, §4.2]), I will implicitly assume that all objects are polarizable in what follows and ignore mentioning this condition.

5. Examples. In this section I will review the main examples that will be of interest in view of Saito's vanishing theorem. I will use freely the notation of the previous sections.

Example 5.1 (The canonical bundle). If X is smooth of dimension n and $\mathbf{V} = \mathbb{Q}_X$ is the constant variation of Hodge structure, we have that $P = \mathbb{Q}_X[n]$, $\mathcal{M} = \omega_X$ with the natural right \mathcal{D} -module structure, and $F_k \omega_X = \omega_X$ for $k \geq -n$, while $F_k \omega_X = 0$ for $k < -n$. The associated Hodge module is usually denoted $\mathbb{Q}_X^H[n]$. The de Rham complex of \mathcal{M} is

$$\mathrm{DR}_X(\omega_X) = \mathrm{DR}_X(\mathcal{O}_X)[n] = \left[\mathcal{O}_X \rightarrow \Omega_X^1 \rightarrow \cdots \rightarrow \Omega_X^n \right][n].$$

Note that

$$\mathrm{Gr}_{-k}^F \mathrm{DR}(\omega_X, F) = \Omega_X^k[n-k] \quad \text{for all } k.$$

Finally, in this example we have $p(\mathcal{M}) = -n$ and $S(\mathcal{M}) = \omega_X$.

Example 5.2 (Direct images). Let $f : X \rightarrow Y$ be a projective morphism with X smooth of dimension n and Y of dimension m , and let \mathbf{V} be a polarizable variation of \mathbb{Q} -Hodge structure of weight k on an open dense subset $U \subset X$, inducing a pure Hodge module M of weight $n+k$ on X as in the previous section. If (\mathcal{M}, F) is the underlying filtered \mathcal{D}_X -module, Theorem 4.3 gives a decomposition

$$f_*(\mathcal{M}, F) \simeq \bigoplus_i (\mathcal{M}_i, F)[-i]$$

in the derived category of filtered regular holonomic \mathcal{D}_Y -modules. According to Theorem 4.2, each (\mathcal{M}_i, F) underlies a pure Hodge module $M_i = \mathcal{H}^i f_* M$ on Y , of weight $n+k+i$. Furthermore, $f_*(\mathcal{M}, F)$ satisfies the strictness property, which is essentially the commutation of the associated graded of the filtration with direct images. A particular case of this is

$$(5.3) \quad \mathbf{R}f_* S(\mathcal{M}) \simeq F_{p(\mathcal{M})}(f_* \mathcal{M}) \simeq \bigoplus_i F_{p(\mathcal{M})} \mathcal{M}_i[-i]$$

in the bounded derived category of coherent sheaves on Y .

For instance, in the case when $\mathbf{V} = \mathbb{Q}_X$ is the constant variation of Hodge structure, by Example 5.1 $p(\mathcal{M}) = -n$ and $S(\mathcal{M}) = \omega_X$. This implies for all i that

$$p(\mathcal{M}_i) = -n \quad \text{and} \quad F_{-n} \mathcal{M}_i = R^i f_* \omega_X.$$

Note that for the corresponding left \mathcal{D} -modules \mathcal{N}_i this means

$$p(\mathcal{N}_i) = m-n \quad \text{and} \quad F_{m-n} \mathcal{N}_i = R^i f_* \omega_{X/Y}.$$

Finally, formula (5.3) specializes to

$$\mathbf{R}f_* \omega_X \simeq \bigoplus_i R^i f_* \omega_X[-i],$$

which is the well-known Kollár decomposition theorem [Kol86b]. Moreover, we will see in Corollary 9.1 and Theorem 11.4 below that $R^i f_* S(\mathcal{M})$ satisfy other important properties known from [Kol86a] in the case of canonical bundles, like vanishing and torsion-freeness.

Example 5.4 (Localization). Let \mathcal{M} be a right \mathcal{D}_X -module and D an effective divisor on a complex manifold X , given locally by an equation f . One can define a new \mathcal{D}_X -module $\mathcal{M}(*D)$ by localizing \mathcal{M} in f ; in other words, globally we have

$$\mathcal{M}(*D) = j_* j^{-1} \mathcal{M},$$

where $j : U \hookrightarrow X$ is the inclusion of the complement $U = X - D$.²

A standard characterization of those \mathcal{D} -modules which do not change under localization will be useful later.

Proposition 5.5. *Let X be a smooth complex variety, D an effective divisor in X , and denote $j : U \hookrightarrow X$ the inclusion of the complement $U = X - D$. Then the restriction functor j^* induces an equivalence between the following categories:*

- (i) *Regular holonomic \mathcal{D}_X -modules \mathcal{M} such that the natural morphism $\mathcal{M} \rightarrow \mathcal{M}(*D)$ is an isomorphism.*
- (ii) *Regular holonomic \mathcal{D}_U -modules.*

Proof. A quick argument is to apply the Riemann-Hilbert correspondence for regular holonomic \mathcal{D} -modules, as the condition defining the category in (i) says that for the perverse sheaf K associated to \mathcal{M} one has $K \simeq j_*j^{-1}K$, i.e. K can be recovered from its restriction to U . \square

Assume now that \mathcal{M} underlies a mixed Hodge module M . By the formula above, $\mathcal{M}(*D)$ underlies the corresponding mixed Hodge module $j_*j^{-1}M$, and so continues to carry a natural Hodge filtration F . This is in general very complicated to compute; the case $\mathcal{M} = \omega_X$, where $\omega_X(*D)$ is the sheaf of meromorphic n -forms on X that are holomorphic on U and the corresponding Hodge module is $j_*\mathbb{Q}_U^H[n]$, is already very relevant. I will say a few words below, and more later.

We always have $F_\ell\omega_X(*D) \cdot F_k\mathcal{D}_X \subseteq F_{k+\ell}\omega_X(*D)$, since the filtration is compatible with the order of differential operators, while by [Sai93, Proposition 0.9] we have

$$(5.6) \quad F_k\omega_X(*D) \subseteq P_k\omega_X(*D) = \begin{cases} \omega_X((n+k+1)D) & \text{if } k \geq -n \\ 0 & \text{if } k < -n, \end{cases}$$

i.e. the Hodge filtration is contained in the filtration by pole order. Furthermore, in [Sai93, Corollary 4.3] it is shown that if D is smooth, then

$$F_k\omega_X(*D) = P_k\omega_X(*D) \quad \text{for all } k.$$

We will see in Section 9 that the first nontrivial step in the filtration is always related to the V -filtration along D , and that this provides a useful relationship with multiplier ideals. For this purpose it is more convenient to write things in terms of left \mathcal{D} -modules. In fact, for the left \mathcal{D} -module $\mathcal{O}_X(*D)$ associated to $\omega_X(*D)$ (recall that $F_p\mathcal{O}_X(*D) = F_{p-n}\omega_X(*D) \otimes \omega_X^{-1}$), one has the formula

$$S(\mathcal{O}_X(*D)) = F_0\mathcal{O}_X(*D) = \tilde{V}^1\mathcal{O}_X \cdot \mathcal{O}_X(D).$$

The V -filtration on \mathcal{M} and $\mathcal{M}(*D)$ is discussed in the next section, and provides further insight into the process of localization.

²A detailed discussion of localization can be found throughout [MS], for instance in §5.3 and §7.4.

6. Duality. For later use, a few words are in order about duality for polarized Hodge modules, on a smooth projective variety X of dimension n . Further discussion and references can be found for instance in [Sch14b, §13 and §29].

A polarization on a pure Hodge module $M = (\mathcal{M}, F; P)$ of weight d is an isomorphism $P(d) \simeq \mathbf{D}P$, where $\mathbf{D}P$ is the Verdier dual of the perverse sheaf P , compatible with the filtration F .³ This means that for the dual holonomic right \mathcal{D}_X -module

$$\mathbf{D}\mathcal{M} := \mathcal{E}xt^n(\mathcal{M}, \omega_X \otimes_{\mathcal{O}_X} \mathcal{D}_X)$$

we have $\mathbf{D}\mathcal{M} \simeq \mathcal{M}$, but furthermore the natural induced filtration on $\mathbf{D}\mathcal{M}$ should satisfy

$$(\mathbf{D}\mathcal{M}, F) \simeq (\mathcal{M}, F_{\bullet-d}).$$

It is necessary therefore for the filtration on $\mathbf{D}\mathcal{M}$ to be strict, something that does not in general hold for arbitrary filtered \mathcal{D} -modules. In fact, it is standard that this strictness property is equivalent to $\mathrm{Gr}_{\bullet}^F \mathcal{M}$ being Cohen-Macaulay as a $\mathrm{Gr}_{\bullet}^F \mathcal{D}_X$ -module; this last statement holds by [Sai88, Lemma 5.1.13] for filtered \mathcal{D} -modules underlying Hodge modules. A consequence is that one can define a dual Hodge module $\mathbf{D}M$, and in fact $\mathbf{D}M \simeq M(d)$, with underlying filtered \mathcal{D} -module $(\mathcal{M}, F_{\bullet-d})$.

Moreover, by [Sai88, §2.4.11] the filtered de Rham complex commutes with the duality functor. Given the discussion above, a useful consequence is:

Lemma 6.1. *If X is a smooth projective variety of dimension n and (\mathcal{M}, F) is the filtered \mathcal{D} -module underlying a pure Hodge module $M \in \mathrm{HM}(X, \ell)$, then*

$$\mathbf{R}\Delta \mathrm{Gr}_k^F \mathrm{DR}_X(\mathcal{M}) \simeq \mathrm{Gr}_{-k-\ell}^F \mathrm{DR}_X(\mathcal{M}),$$

where $\mathbf{R}\Delta = \mathbf{R}\mathcal{H}om_{\mathcal{O}_X}(\cdot, \omega_X)[n]$ is the Grothendieck duality functor.

7. The V -filtration. In this section I will recall some key definitions and results regarding the V -filtration with respect to a hypersurface, and its interaction with the Hodge filtration. I am mostly following [Sai88, §3], which is a complete reference for all the definitions and results recalled here.

Let X be a complex manifold or smooth complex variety of dimension n , and let X_0 be an smooth divisor on X defined locally by an equation t . We first consider a rational filtration on \mathcal{D}_X , given by

$$V_\alpha \mathcal{D}_X = \{P \in \mathcal{D}_X \mid P \cdot \mathcal{I}_{X_0}^j \subseteq \mathcal{I}_{X_0}^{j-[\alpha]}\} \text{ for } \alpha \in \mathbb{Q},$$

where \mathcal{I}_{X_0} is the ideal of X_0 in \mathcal{O}_X , with the convention that $\mathcal{I}_{X_0}^j = \mathcal{O}_X$ for $j \leq 0$.

Definition 7.1 (V -filtration). Let \mathcal{M} be a coherent right \mathcal{D}_X -module. A *rational V -filtration* (a slight refinement of the *Kashiwara-Malgrange filtration*) of \mathcal{M} along X_0 is an increasing filtration $V_\alpha \mathcal{M}$ with $\alpha \in \mathbb{Q}$ satisfying the following properties:

- The filtration is exhaustive, i.e. $\bigcup_\alpha V_\alpha \mathcal{M} = \mathcal{M}$, and each $V_\alpha \mathcal{M}$ is a coherent $V_0 \mathcal{D}_X$ -submodule of \mathcal{M} .

³Together of course with further properties compatible with the inductive definition of Hodge modules by dimension of support via modules of nearby cycles.

- $V_\alpha \mathcal{M} \cdot V_i \mathcal{D}_X \subseteq V_{\alpha+i} \mathcal{M}$ for every $\alpha \in \mathbb{Q}$ and $i \in \mathbb{Z}$; furthermore

$$V_\alpha \mathcal{M} \cdot t = V_{\alpha-1} \mathcal{M} \quad \text{for } \alpha < 0.$$

- The action of $t\partial_t - \alpha$ on $\text{Gr}_\alpha^V \mathcal{M}$ is nilpotent for each α , where ∂_t is a vector field such that $[\partial_t, t] = 1$. (One defines $\text{Gr}_\alpha^V \mathcal{M}$ as $V_\alpha \mathcal{M} / V_{<\alpha} \mathcal{M}$, where $V_{<\alpha} \mathcal{M} = \cup_{\beta < \alpha} V_\beta \mathcal{M}$.)

It is known that if a V -filtration exists, then it is unique. In addition, \mathcal{D} -modules underlying mixed Hodge modules also come by definition with a Hodge filtration, and it is important to compare the two. Note first that on each $\text{Gr}_\alpha^V \mathcal{M}$ one considers the filtration induced by that on \mathcal{M} , i.e.

$$F_p \text{Gr}_\alpha^V \mathcal{M} := \frac{F_p \mathcal{M} \cap V_\alpha \mathcal{M}}{F_p \mathcal{M} \cap V_{<\alpha} \mathcal{M}}.$$

Definition 7.2 (Regular and quasi-unipotent). In the situation above, assume that \mathcal{M} is endowed with a good filtration F . We say that (\mathcal{M}, F) is *quasi-unipotent* (or *strictly specializable*) along X_0 if \mathcal{M} admits a rational V -filtration along X_0 and the following conditions are satisfied:

- $(F_p V_\alpha \mathcal{M}) \cdot t = F_p V_{\alpha-1} \mathcal{M}$ for $\alpha < 0$.
- $(F_p \text{Gr}_\alpha^V \mathcal{M}) \cdot \partial_t = F_{p+1} \text{Gr}_{\alpha+1}^V \mathcal{M}$ for $\alpha > -1$.

One says that (\mathcal{M}, F) is *regular and quasi-unipotent along* X_0 if in addition the filtration $F_\bullet \text{Gr}_\alpha^V \mathcal{M}$ is a good filtration for $-1 \leq \alpha \leq 0$.

Let now $f : X \rightarrow \mathbb{C}$ be a holomorphic function, and denote by

$$i = i_{\Gamma_f} : X \hookrightarrow X \times \mathbb{C} = Y$$

the embedding of X as the graph of f . Denote by t the coordinate on \mathbb{C} , so that in the notation above we have $X_0 = X \times \{0\} = t^{-1}(0)$. If \mathcal{M} is a coherent right \mathcal{D}_X -module, denote $(\tilde{\mathcal{M}}, F) = i_*(\mathcal{M}, F)$. One says that (\mathcal{M}, F) is *strictly specializable along* f if $(\tilde{\mathcal{M}}, F)$ is so along X_0 , and the same for *regular and quasi-unipotent along* f . One important feature of mixed Hodge module theory is that all \mathcal{D} -modules underlying Hodge modules are required to satisfy this last property with respect to any holomorphic function.

The following technical result on the behavior of regular and quasi-unipotent filtered \mathcal{D} -modules is a key step in extending Kashiwara's theorem on closed embeddings to the setting of Hodge \mathcal{D} -modules. This will be very useful when stating Saito's vanishing theorem on singular varieties in Section 8.

Lemma 7.3 ([Sai90, Lemma 3.2.6]). *Let $f : X \rightarrow \mathbb{C}$ be a holomorphic function, and (\mathcal{M}, F) a filtered coherent \mathcal{D}_X -module. Assume that $\text{Supp}(\mathcal{M}) \subseteq f^{-1}(0)$. Then the following are equivalent:*

- (i) (\mathcal{M}, F) is regular and quasi-unipotent along f .
- (ii) $\text{Gr}_p^F \mathcal{M} \cdot f = 0$ for all p .
- (iii) $(\tilde{\mathcal{M}}, F) \simeq j_*(\mathcal{M}, F)$, where $j : X \times \{0\} \hookrightarrow X \times \mathbb{C}$.

We will also need a transversality notion for a filtered \mathcal{D} -module with respect to a morphism (or a submanifold) introduced in [Sai88, 3.5.1], under which filtered inverse images become particularly simple.

Definition 7.4 (Non-characteristic morphism). Let $f : X \rightarrow Y$ be a morphism of complex manifolds, and let (\mathcal{M}, F) be a filtered coherent \mathcal{D}_Y -module. One says that f is *non-characteristic* for (\mathcal{M}, F) if the following two conditions are satisfied:

- $\mathcal{H}^i(f^{-1} \mathrm{Gr}^F \mathcal{M} \otimes_{f^{-1}\mathcal{O}_Y} \mathcal{O}_X) = 0$ for $i \neq 0$.⁴
- The natural morphism

$$df^* : p_2^{-1}(\mathrm{Char}(\mathcal{M})) \rightarrow T^*X$$

is finite, where $p_2 : X \times_Y T^*Y \rightarrow T^*Y$ is the second projection and

$$df^* : X \times_Y T^*Y \rightarrow T^*X, \quad (x, \omega) \mapsto df^*\omega \quad \text{for all } x \in X, \omega \in T^*Y.$$

If f is a closed immersion, we say that X is non-characteristic for (\mathcal{M}, F) if f is so.

If f is non-characteristic for (\mathcal{M}, F) and $d = \dim X - \dim Y$, then as in [Sai88, §3.5] one has the filtered pullback $\mu^*(\mathcal{M}, F) = (\tilde{\mathcal{M}}, F)[-d]$ given by the formula

$$\tilde{\mathcal{M}} = \mu^{-1}\mathcal{M} \otimes_{\mu^{-1}\mathcal{O}_X} \omega_{Y/X} \quad \text{and} \quad F_p\tilde{\mathcal{M}} = \mu^{-1}F_{p+d}\mathcal{M} \otimes_{\mu^{-1}\mathcal{O}_X} \omega_{Y/X}.$$

In other words we can define the inverse image to be, up to shift, the naive filtration on the naive pullback, and this again gives a holonomic \mathcal{D}_X -module if \mathcal{M} is so; see [Sai88, Lemma 3.5.5]. If (\mathcal{M}, F) underlies a pure Hodge module M of weight m on Y , $\mu^*(\mathcal{M}, F)$ then underlies a pure Hodge module μ^*M of weight $m + d$ on X .⁵

Example 7.5. The most basic examples of this notion are:

- If f is smooth, then it is non-characteristic for any (\mathcal{M}, F) , as df^* is injective and f is flat.
- If (\mathcal{M}, F) underlies a variation of Hodge structure, any f is non-characteristic for it, as $\mathrm{Char}(\mathcal{M})$ is the zero section, while each $\mathrm{Gr}_k^F \mathcal{M}$ is locally free.

As a combination of the two, if f is smooth outside of the locus where (\mathcal{M}, F) underlies a variation of Hodge structure, then f is non-characteristic for (\mathcal{M}, F) .

The following lemmas are important in what follows; they show that under the non-characteristicity assumption one can perform concrete calculations with the V -filtration.

Lemma 7.6 ([Sai88, Lemme 3.5.6]). *Let $i : D \hookrightarrow X$ be an inclusion of a smooth hypersurface in a smooth complex variety. Let (\mathcal{M}, F) be a filtered coherent right \mathcal{D}_X -module for which D is non-characteristic. Then*

- (1) (\mathcal{M}, F) is regular and quasi-unipotent along D .

⁴In terms of the individual graded pieces, which are coherent sheaves of Y , this simply says that $L^i f^* \mathrm{Gr}_k^F \mathcal{M} = 0$ for all $i \neq 0$ and all k .

⁵This is a non-trivial result, using the fact that pure Hodge modules with strict support come from generic variations of Hodge structure; see e.g. [Sch14b, §30] for an explanation.

(2) The V -filtration on M is given by

$$V_\alpha \mathcal{M} = \mathcal{M} \cdot \mathcal{O}_X(-iD) \text{ for } -i-1 \leq \alpha < -i, i \geq 0 \text{ and } V_\alpha \mathcal{M} = \mathcal{M} \text{ for } \alpha \geq 0.$$

Lemma 7.7 ([Sai88, Lemme 3.5.7]). *With the notation of Lemma 7.6, we have that*

(1) The V -filtration on $\mathcal{M}(*D)$ satisfies

$$V_\alpha \mathcal{M}(*D) = \mathcal{M} \cdot \mathcal{O}_X(-iD) \text{ for } -i-1 \leq \alpha < -i.$$

(2) There is a filtration F on $\mathcal{M}(*D)$ which makes it a filtered coherent right \mathcal{D}_X -module, such that there is an exact sequence of filtered \mathcal{D} -modules

$$0 \longrightarrow (\mathcal{M}, F) \longrightarrow (\mathcal{M}(*D), F) \longrightarrow i_* i^1(\mathcal{M}, F)[1] \longrightarrow 0.$$

In addition, $(\mathcal{M}(*D), F)$ is regular and quasi-unipotent along D .

It will also be crucial, under suitable hypotheses, to be able to recover the Hodge filtration from its restriction over the complement of a hypersurface. This is one of the key points of the interaction between the Hodge filtration and the V -filtration in the case of filtered \mathcal{D} -modules underlying Hodge modules.

Lemma 7.8 ([Sai88, Proposition 3.1.8]). *With the notation of Lemma 7.6, and $U = X - D$, let \mathcal{M}' be the smallest sub-object of \mathcal{M} such that $\mathcal{M}|_U = \mathcal{M}'|_U$. Then:*

(1) $\mathcal{M}' = V_\alpha \mathcal{M} \cdot \mathcal{D}_X$ for $\alpha < 0$.

(2) $\mathcal{M}/\mathcal{M}' \simeq i_* \text{Coker}(\text{can} = \partial_t : \text{Gr}_{-1}^V \mathcal{M} \rightarrow \text{Gr}_0^V \mathcal{M})$.

In particular, $\mathcal{M} = V_\alpha \mathcal{M} \cdot \mathcal{D}_X$ for $\alpha < 0$ if can is surjective.

Lemma 7.9 ([Sai88, Proposition 3.2.2]). *With the notation of Lemma 7.8, and $j : U \rightarrow X$ the natural inclusion, we have that:*

(1) The first condition in Definition 7.2 is equivalent to

$$F_p V_{<0} \mathcal{M} = V_{<0} \mathcal{M} \cap j_* j^{-1} F_p \mathcal{M} \text{ for all } p.$$

(2) If $\mathcal{M} = V_{<0} \mathcal{M} \cdot \mathcal{D}_X$, or equivalently if $\text{can} = \partial_t : \text{Gr}_{-1}^V \mathcal{M} \rightarrow \text{Gr}_0^V \mathcal{M}$ is surjective, the second condition in Definition 7.2 for $\alpha \geq -1$ ⁶ is equivalent to

$$F_p \mathcal{M} = \sum_{i \geq 0} (F_{p-i} V_{<0} \mathcal{M}) \cdot \partial_t^i \text{ for all } p.$$

8. Kodaira-Saito vanishing. We now come to the main goal, M. Saito's vanishing theorem. Before stating and proving the theorem, it is important to emphasize the following point: this is a result that works on *singular* varieties by embedding them into smooth ambient spaces. It is known that the objects considered are independent of the embedding.

⁶There is an extra point here, for which I am grateful to C Sabbah: in Definition 7.2 one only considers $\alpha > -1$, while in the lemma $\alpha = -1$ appears as well. However, the property we want for $\alpha = -1$ follows from Hodge theory conditions on Gr_{-1}^V and Gr_0^V ; in our application they will be trivially satisfied since both terms will be 0.

It is therefore important to have a way of thinking about mixed Hodge modules and filtered \mathcal{D} -modules on singular varieties, compatible with the material developed for smooth varieties. In general this can only be done by locally embedding X into smooth ambient spaces, and then using a gluing procedure (see [Sai90, §2.1]).

However, on projective varieties we can use the embedding of X into some \mathbf{P}^N . If $X \hookrightarrow \mathbf{P}^N$ is one such, then one defines the category of mixed Hodge modules on X to be that of mixed Hodge modules on \mathbf{P}^N with support contained in X , i.e.

$$\mathrm{MHM}(X) = \mathrm{MHM}_X(\mathbf{P}^N).$$

One can do the same with any embedding $X \subset Z$ into a smooth variety; at least when Z is projective, the fact that the resulting $\mathrm{MHM}(X)$ is independent of the embedding follows by extending Kashiwara's equivalence theorem for closed embeddings to the setting of Hodge modules.

Indeed, recall that Kashiwara's theorem says that for a closed embedding $h : Z \hookrightarrow W$ one has

$$\mathrm{Mod}_{\mathrm{coh}}(\mathcal{D}_Z) \simeq \mathrm{Mod}_{\mathrm{coh}, Z}(\mathcal{D}_W),$$

where the category on the right is that of coherent \mathcal{D}_W -modules with support contained in Z . This correspondence restricts on both sides to the subcategories of objects with support contained in X . The equivalence does not extend in general to filtered \mathcal{D} -modules; however, those underlying mixed Hodge modules are regular and quasi-unipotent (Definition 7.2) along the zero-locus of any holomorphic function.

In the regular and quasi-unipotent case, one can use Lemma 7.3 for each local defining equation f for Z inside W (or global equations when $W = \mathbf{P}^N$) in order to deduce that for every (\mathcal{M}, F) on W with support in Z , there exists (\mathcal{M}_Z, F) on Z such that $(\mathcal{M}, F) \simeq h_*(\mathcal{M}_Z, F)$. Thus Kashiwara's theorem extends to these special filtered holonomic \mathcal{D} -modules, which is the key step in extending it to mixed Hodge modules. Once this is established, it is not too hard to deduce that $\mathrm{MHM}(X)$ is independent of the embedding; formally

$$(8.1) \quad \mathrm{HM}(X, \ell) = \mathrm{HM}_X(Z, \ell) \quad \text{and} \quad \mathrm{MHM}(X) = \mathrm{MHM}_X(Z)$$

for any smooth Z containing X . Further details can be found in [Sai88, Lemme 5.1.9] and [Sai90, 2.17.5]; see also [Sch14a, §6 and 7].

Theorem 8.2 (M. Saito, [Sai90, §2.g]). *Let X be a complex projective variety, and L an ample line bundle on X . Consider an integer $m > 0$ such that $L^{\otimes m}$ is very ample and gives an embedding $X \subseteq \mathbf{P}^N$. Let (\mathcal{M}, F) be the filtered \mathcal{D} -module underlying a mixed Hodge module M on \mathbf{P}^N with support contained in X , i.e. an object in $\mathrm{MHM}(X)$. Then:*

(1) $\mathrm{Gr}_k^F \mathrm{DR}_{\mathbf{P}^N}(\mathcal{M})$ is an object in $\mathbf{D}^b(X)$ for each k , independent of the embedding of X in \mathbf{P}^N .⁷

(2) We have the hypercohomology vanishing

$$H^i(X, \mathrm{Gr}_k^F \mathrm{DR}_{\mathbf{P}^N}(\mathcal{M}) \otimes L) = 0 \quad \text{for all } i > 0.$$

⁷In fact, based on the discussion above it can be shown that each $\mathrm{Gr}_k^F \mathrm{DR}_{\mathbf{P}^N}(\mathcal{M})$ is independent of the embedding of X into any smooth complex variety.

and

$$H^i(X, \mathrm{Gr}_k^F \mathrm{DR}_{\mathbf{P}^N}(\mathcal{M}) \otimes L^{-1}) = 0 \quad \text{for all } i < 0.$$

Proof. Step 1. This step addresses (1) and a number of useful reductions towards (2). For the first statement in (1), due to the definition of $\mathrm{Gr}_k^F \mathrm{DR}_{\mathbf{P}^N}(\mathcal{M})$, it is enough to have that each $\mathrm{Gr}_k^F \mathcal{M}$ is an \mathcal{O}_X -module. But note that by Lemma 7.3, if for a holomorphic function f the support of \mathcal{M} is contained in $f^{-1}(0)$, the condition of (\mathcal{M}, F) being regular and quasi-unipotent along f is equivalent to having

$$\mathrm{Gr}_p^F \mathcal{M} \cdot f = 0 \quad \text{for all } p.$$

Now our (\mathcal{M}, F) satisfies this for any f , as it underlies a Hodge module, and applying it for the defining equations of X inside \mathbf{P}^N we obtain the conclusion.

Note that the independence on the embedding of the definition $\mathrm{MHM}(X) = \mathrm{MHM}_X(\mathbf{P}^N)$ follows from the discussion preceding the statement of the theorem. However here strictly speaking one only needs to know independence of embeddings $X \hookrightarrow \mathbf{P}^N$ by various powers $L^{\otimes m}$. Thus the Kashiwara-type statement (8.1) actually suffices, as any two such can be compared inside a common Veronese embedding.

Along the same lines, the independence of the embedding for complex of \mathcal{O}_X -modules $\mathrm{Gr}_k^F \mathrm{DR}_{\mathbf{P}^N}(\mathcal{M})$ follows then from the remark above and the fact that if $h : Z \hookrightarrow W$ is a closed embedding of two smooth varieties containing X , and $(\mathcal{M}, F) \simeq h_*(\mathcal{M}_Z, F)$ on W , then one has the easily checked formula

$$\mathrm{Gr}_k^F \mathrm{DR}_W(\mathcal{M}, F) \simeq h_* \mathrm{Gr}_k^F \mathrm{DR}_Z(\mathcal{M}_Z, F).$$

Based on the fact that our objects do not depend on the embedding $X \subseteq \mathbf{P}^N$, to attack (2) we may assume furthermore that $m \geq 2$. This will come up later, as we will need to produce non-integral rational numbers with denominator m .

A standard reduction is that it is enough to assume that M is a polarized pure Hodge module with strict support X , of some weight d . First, once we have reduced to the case of pure Hodge modules, we can apply the strict support direct sum decomposition (4.1) to reduce to this case. On the other hand, if M is in $\mathrm{MHM}(X)$, recall that it has a finite weight filtration $W_\bullet M$ by objects in $\mathrm{MHM}(X)$, such that the graded quotients $\mathrm{Gr}_\ell^W M = W_\ell M / W_{\ell-1} M$ are in $\mathrm{HM}(X, \ell) = \mathrm{HM}_X(\mathbf{P}^N, \ell)$. To reduce to the pure case, we simply use the fact that the functor $\mathrm{Gr}_k^F \circ \mathrm{DR}$ is exact by construction.

Given this last reduction, we also see that it is enough to check only the second statement in (2). This follows from Grothendieck-Serre duality and Lemma 6.1.

Step 2. Let Y be a general hyperplane in \mathbf{P}^N , chosen to be non-characteristic for (\mathcal{M}, F) . Denote $D = X \cap Y$, the zero locus of some section $s \in H^0(X, L^{\otimes m})$. Let $f : \tilde{X} \rightarrow X$ be the m -fold cyclic cover branched along D as in Proposition 2.2, with $f^*D = mD'$ and $L' = \mathcal{O}_{\tilde{X}}(D')$.

Denote now

$$U = \mathbf{P}^N - Y \quad \text{and} \quad j : U \hookrightarrow \mathbf{P}^N$$

the natural inclusion of the (affine) complement of Y . Denoting also by $i : Y \hookrightarrow \mathbf{P}^N$ the inclusion of Y , by Lemma 7.7 there is a filtered short exact sequence

$$(8.3) \quad 0 \longrightarrow (\mathcal{M}, F) \longrightarrow (\mathcal{M}(*Y), F) \longrightarrow (\mathcal{H}^1 i^! \mathcal{M}, F) \longrightarrow 0$$

(Note that here $\mathcal{H}^1 i^! \mathcal{M}$ simply means $\mathcal{M} \otimes \omega_{Y/\mathbf{P}^N}$.)

For each k , we apply the exact functor $\mathrm{Gr}_k^F \circ \mathrm{DR}_{\mathbf{P}^N}$ to (8.3) to obtain a distinguished triangle of complexes of coherent sheaves on X :

$$\begin{aligned} \mathrm{Gr}_k^F \mathrm{DR}_{\mathbf{P}^N}(\mathcal{M}) \otimes L^{-1} &\longrightarrow \mathrm{Gr}_k^F \mathrm{DR}_{\mathbf{P}^N}(\mathcal{M}(*Y)) \otimes L^{-1} \longrightarrow \\ &\longrightarrow \mathrm{Gr}_k^F \mathrm{DR}_{\mathbf{P}^N}(\mathcal{H}^1 i^! \mathcal{M}) \otimes L^{-1} \longrightarrow \mathrm{Gr}_k^F \mathrm{DR}_{\mathbf{P}^N}(\mathcal{M}) \otimes L^{-1}[1]. \end{aligned}$$

The claim is that

$$(8.4) \quad H^i(X, \mathrm{Gr}_k^F \mathrm{DR}_{\mathbf{P}^N}(\mathcal{M}(*Y)) \otimes L^{-1}) = 0 \quad \text{for all } i \neq 0.$$

This will be proved in Step 4. Assuming it for now, by the long exact sequence on cohomology we are reduced to showing

$$H^i(X, \mathrm{Gr}_k^F \mathrm{DR}_{\mathbf{P}^N}(\mathcal{H}^1 i^! \mathcal{M}) \otimes L^{-1}) = 0 \quad \text{for all } i < -1.$$

But in fact the statement is true even for $i < 0$ by induction on $n = \dim X$, since $(\mathcal{H}^1 i^! \mathcal{M}, F)$ is supported on D and, again by non-characteristic pullback as in Section 7 it underlies a Hodge module in $\mathrm{HM}_D(Y, d+1)$.

Step 3. Note first that we can extend the cover $f : \tilde{X} \rightarrow X$ ramified over D to a cover still denoted $f : \tilde{\mathbf{P}}^N \rightarrow \mathbf{P}^N$, ramified over Y ; it is enough to do this locally since Hodge modules are local by construction. Fix a point $x \in X$. The claim is that there exists a neighborhood $x \in U_x \subset \mathbf{P}^N$ such that the restriction of $f : \tilde{X} \rightarrow X$ over $U_x \cap X$ can be extended to a finite cover $f_x : V_x \rightarrow U_x$, ramified over $Y \cap U_x$. If $x \notin Y$, it is clear that there is such an extension. On the other hand, if $x \in Y$, then we use a local holomorphic trivialization $(U_x, U_x \cap X) \simeq (U_x \cap Y, U_x \cap D) \times \mathbb{D}^2$, considering a contractible neighborhood of x in Y such that the contraction is compatible with D .

This new f is non-characteristic for (\mathcal{M}, F) by our choice of Y , and so the filtered pullback $(f^* \mathcal{M}, F)$ on $\tilde{\mathbf{P}}^N$ can be defined as in the remarks after Definition 7.4. It underlies a pure Hodge module $f^* M$ of weight d , as the relative dimension is zero. By Theorem 4.2 we then obtain $f_* f^* M \in \mathrm{HM}_X(\mathbf{P}^N, d)$; note that this is a single Hodge module since f is finite. There is a natural monomorphism $M \rightarrow f_* f^* M$, and we define \tilde{M} as its cokernel, so that there is an exact sequence

$$(8.5) \quad 0 \longrightarrow M \longrightarrow f_* f^* M \longrightarrow \tilde{M} \longrightarrow 0,$$

in the abelian category $\mathrm{HM}_X(\mathbf{P}^N, d)$, i.e. \tilde{M} is a new pure polarized Hodge module of weight d with support contained in X . Note that by Saito's fundamental result mentioned in Section 4, all the Hodge modules in the exact sequence above are uniquely extended from the open subset of U on which they are variations of Hodge structure; in particular they coincide with the strict support extension of their restriction to U .

If we denote by \tilde{P} the \mathbb{Q} -perverse sheaf associated to \tilde{M} , so that $\mathrm{DR}_{\mathbf{P}^N}(\tilde{\mathcal{M}}) \simeq \tilde{P}_{\mathbb{C}} := \tilde{P} \otimes \mathbb{C}$. Since as mentioned above \tilde{M} is the unique extension with strict

support X of its restriction to U , we have

$$\tilde{P} \simeq j_* j^{-1} \tilde{P},$$

i.e. \tilde{P} is the extension of its restriction to the affine open set U as well. By the Artin-Grothendieck vanishing theorem (see e.g. [Laz04, Theorem 3.1.13]), we then have

$$H^i(X, \tilde{P}_{\mathbb{C}}) \simeq H^i(U, j^{-1} \tilde{P}_{\mathbb{C}}) = 0 \text{ for all } i > 0.$$

Since \tilde{M} is polarized, as in Section 6.1 we have that $\mathbf{D}\tilde{P} \simeq \tilde{P}(d)$, where $\mathbf{D}\tilde{P}$ is the Verdier dual. By Verdier duality we then also get

$$H^i(X, \tilde{P}_{\mathbb{C}}) = 0 \text{ for all } i < 0.$$

In conclusion, we have verified that

$$(8.6) \quad H^i(X, \mathrm{DR}_{\mathbf{P}^N}(\tilde{\mathcal{M}})) = 0 \text{ for all } i \neq 0.$$

The main assertion in this step is that

$$(8.7) \quad H^i(X, \mathrm{Gr}_k^F \mathrm{DR}_{\mathbf{P}^N}(\tilde{\mathcal{M}})) = 0 \text{ for all } k \text{ and all } i \neq 0.$$

To this end we need to use stability under projective morphisms, Theorem 4.2; applied to the absolute case $\mathbf{P}^N \rightarrow \mathrm{pt}$, the strictness in the statement amounts to the degeneration at E_1 of the natural Hodge-to-de Rham spectral sequence

$$E_1^{p,q} = H^{p+q}(X, \mathrm{Gr}_{-q}^F \mathrm{DR}_{\mathbf{P}^N}(\tilde{\mathcal{M}})) \implies H^{p+q}(X, \mathrm{DR}_{\mathbf{P}^N}(\tilde{\mathcal{M}})).$$

Note that here we are using the identification $f_* \tilde{\mathcal{M}} \simeq \mathbf{R}\Gamma \mathrm{DR}_{\mathbf{P}^N}(\tilde{\mathcal{M}})$ (which is a special case of the definition of push-forward via smooth morphisms). Given (8.6), this degeneration immediately implies (8.7).

Step 4. We are left with proving (8.4), which will be done in this step. More precisely, for each k we will prove the isomorphism

$$(8.8) \quad \mathrm{Gr}_k^F \tilde{\mathcal{M}} \simeq \mathrm{Gr}_k^F \mathcal{M}(*Y) \otimes \tilde{L},$$

where

$$\tilde{L} := \mathrm{Coker}(\mathcal{O}_X \rightarrow \pi_* \mathcal{O}_{\tilde{X}}) \simeq L^{-1} \oplus \dots \oplus L^{-(m-1)},$$

the last isomorphism coming from Proposition 2.2. The isomorphism (8.8) implies what we want in combination with (8.7); it is proved using the interaction between the Hodge filtration and the V -filtration along Y .

To this end, note first that by definition there is a canonical isomorphism of filtered right \mathcal{D}_U -modules

$$(\tilde{\mathcal{M}}, F)|_U \simeq (\mathcal{M}, F) \otimes \tilde{L}|_U.$$

Indeed, this follows from (8.5) and the definition of the filtration on $f^* \mathcal{M}$ given after Definition 7.4; passing to the filtration on the \mathcal{D} -modules underlying (8.5) is, on the open set U on which f is étale, the same as the split short exact sequence

$$0 \longrightarrow F_p \mathcal{M} \longrightarrow F_p \mathcal{M} \otimes f_* \mathcal{O}_{\mathbf{P}^N} \longrightarrow F_p \mathcal{M} \otimes \tilde{L} \longrightarrow 0.$$

Here we consider both $f_* \mathcal{O}_{\mathbf{P}^N}$ and \tilde{L} as left \mathcal{D} -modules (integrable connections) with the trivial filtration, so that the tensor products $\mathcal{M} \otimes f_* \mathcal{O}_{\mathbf{P}^N}$ and $\mathcal{M} \otimes \tilde{L}$ are again filtered right \mathcal{D} -modules.

The statement follows if we show that the isomorphism on U above can be extended uniquely to an isomorphism of filtered \mathcal{D} -modules

$$(8.9) \quad (\tilde{\mathcal{M}}, F) \simeq (\mathcal{M}(*Y), F) \otimes_{\mathcal{O}_{\mathbb{P}^N}} \tilde{L}.$$

Here again \tilde{L} is considered as a left \mathcal{D} -module associated to an integrable connection. Both sides of (8.9) are regular holonomic; moreover, they are isomorphic to their localization along Y , i.e. a local equation of Y acts on them bijectively. Forgetting about the filtration, the isomorphism in (8.9) then follows from Proposition 5.5.

As for the filtration F , we need to compare it to the V -filtration along the divisor Y . The first claim is that $(\mathcal{M}(*Y), F) \otimes \tilde{L}$ is regular and quasi-unipotent along Y ; see Definition 7.2. To this end, note first that the summand $\mathcal{M}(*Y) \otimes L^{-i}$ of $\mathcal{M}(*Y) \otimes \tilde{L}$, with i ranging from 1 to $m-1$, coincides with $\mathcal{M}(*Y)$ on U , and so their V -filtrations along Y are the same for $\alpha < 0$. On the other hand, if t is a local equation for Y , then multiplication by L^{-i} coincides with the action of $t^{i/m}$, and so by the definition of the V -filtration we obtain for each α :

$$(8.10) \quad V_\alpha(\mathcal{M}(*Y) \otimes L^{-i}) = V_{\alpha+i/m}\mathcal{M}(*Y) \otimes L^{-i}.$$

This gives in particular

$$F_p V_\alpha(\mathcal{M}(*Y) \otimes L^{-i}) = (F_p V_{\alpha+i/m}\mathcal{M}(*Y)) \otimes L^{-i}$$

for all possible indices. Using this, the fact that $(\mathcal{M}(*Y), F) \otimes \tilde{L}$ is regular and quasi-unipotent is an immediate consequence of the fact that $\mathcal{M}(*Y)$ is so (as it underlies a mixed Hodge module), together with Lemma 7.7(1).

From (8.10) we also obtain that

$$\mathrm{Gr}_\alpha^V(\mathcal{M}(*Y) \otimes L^{-i}) \simeq \mathrm{Gr}_{\alpha+i/m}^V(\mathcal{M}(*Y)) \otimes L^{-i}.$$

We see however from Lemma 7.7(1) that

$$\mathrm{Gr}_\alpha^V \mathcal{M}(*Y) = 0 \quad \text{for } \alpha \notin \mathbb{Z},$$

and therefore

$$\mathrm{Gr}_\alpha^V(\mathcal{M}(*Y) \otimes L^{-i}) = 0 \quad \text{for } \alpha + i/m \notin \mathbb{Z}.$$

The bottom line is that in order to have $\mathrm{Gr}_\alpha^V(\mathcal{M}(*Y) \otimes \tilde{L}) \neq 0$, one must have $\alpha + i/m \in \mathbb{Z}$ for all $1 \leq i \leq m-1$, and consequently α cannot be an integer (recall that we are working with $m \geq 2$).

Let's now denote $\mathcal{M}' = \mathcal{M}(*Y) \otimes \tilde{L}$ for simplicity. Using this last remark, by Lemma 7.8 we deduce that \mathcal{M}' is generated as a \mathcal{D} -module by the negative part of its V -filtration, i.e.

$$\mathcal{M}' \simeq V_{<0}\mathcal{M}' \cdot \mathcal{D}_{\mathbb{P}^N}.$$

The next thing to note is that, again since the jumps in the V -filtration do not happen at integers, according to Lemma 7.9(2) the second condition in Definition 7.2 is equivalent to the fact that

$$F_p \mathcal{M}' = \sum_{i \geq 0} (F_{p-i} V_{<0} \mathcal{M}') \cdot \partial_t^i$$

for all p . Consequently, the Hodge filtration is determined by its restriction to the negative part of the V -filtration. Finally, this restriction is determined by

the corresponding filtration on the open complement U since according to Lemma 7.9(1) for all p we have

$$F_p V_{<0} \mathcal{M}' = V_{<0} \mathcal{M}' \cap j_* j^{-1} F_p \mathcal{M}'.$$

As $(\tilde{\mathcal{M}}, F)$ and $(\mathcal{M}(*Y), F) \otimes \tilde{L}$ coincide on U , and as $(\tilde{\mathcal{M}}, F)$ was defined by extension from U , the two filtered \mathscr{D} -modules must then agree everywhere. \square

9. Particular cases. In this section I will explain how M. Saito's vanishing theorem can be used to deduce many of the standard vanishing theorems. In the next sections I will make the point however that the abstract version is equally valuable for concrete applications.

Kodaira-Nakano vanishing. Let X be a smooth projective complex variety of dimension n . We consider the trivial Hodge module $M = \mathbb{Q}_X^H[n]$. According to Example 5.1, the corresponding right \mathscr{D} -module is ω_X , with filtration $F_p \omega_X = \omega_X$ if $p \geq -n$ and $F_p \omega_X = 0$ if $p < -n$, so that

$$\mathrm{Gr}_{-p}^F \mathrm{DR}_X(\omega_X) = \Omega_X^p[n-p] \quad \text{for all } p.$$

Theorem 8.2 gives

$$H^q(X, \Omega_X^p \otimes L) = 0 \quad \text{for } p + q > n$$

and the dual statement, for any L ample, i.e. Kodaira-Nakano vanishing.

If we restrict to the Kodaira vanishing theorem, which corresponds to the lowest non-zero piece of the filtration on ω_X , then we can see it as an example of the following more easily stated special case of Theorem 8.2; it is useful to record this for applications.

Corollary 9.1. *If (\mathcal{M}, F) is a filtered \mathscr{D} -module underlying a mixed Hodge module M on a projective variety X , and L is an ample line bundle on X , then*

$$H^i(X, S(\mathcal{M}) \otimes L) = 0 \quad \text{for all } i > 0.$$

Kollár vanishing. The following theorem of Kollár is a natural generalization of Kodaira vanishing to higher direct images of canonical bundles.

Theorem 9.2 ([Kol86a, Theorem 2.1(iii)]). *Let $f : X \rightarrow Y$ be a morphism between complex projective varieties, with X smooth, and let L be an ample line bundle on Y . Then*

$$H^i(X, R^j f_* \omega_X \otimes L) = 0 \quad \text{for all } i > 0 \quad \text{and all } j.$$

To deduce the statement from Theorem 8.2, we consider the push-forward $M = f_* \mathbb{Q}_X^H[n]$ of the trivial Hodge module on X , with $n = \dim X$. According to Example 5.2, for the underlying \mathscr{D} -modules we have

$$f_*(\omega_X, F) \simeq \bigoplus_i (\mathcal{M}_i, F)[-i]$$

in the derived category of filtered \mathscr{D}_Y -modules (so compatible with inclusions into smooth varieties), and for each i we have $S(\mathcal{M}_i) = R^i f_* \omega_X$. Theorem 9.2 then follows from Corollary 9.1. More generally, the same argument shows the following vanishing theorem due to Saito: the statement of Theorem 9.2 holds for $R^i f_* S(\mathcal{M})$,

where \mathcal{M} corresponds to the unique pure Hodge module with strict support X extending a polarized variation of Hodge structure on an open set $U \subseteq X$.

Nadel vanishing. To deduce Nadel vanishing, one needs a more subtle relationship between multiplier ideals, the V -filtration on the structure sheaf, and the Hodge filtration on localizations, combining results of [BS05] and [Sai09]. As mentioned in Example 5.4, this is one place where it is more convenient to have the initial discussion in terms of left \mathcal{D} -modules.

Let X be a smooth projective variety, and D an effective Cartier divisor on X . Recall that $\mathcal{O}_X(*D)$ is equipped with a natural Hodge filtration F , as the left \mathcal{D} -module associated to the Hodge module $j_*\mathbb{Q}_U^H[n]$, where $j : U = X - D \hookrightarrow X$ is the inclusion; see Example 5.4. Looking at the first step in this filtration, one can recognize multiplier ideals from the formula

$$(9.3) \quad F_0\mathcal{O}_X(*D) \simeq \mathcal{J}((1 - \varepsilon)D) \cdot \mathcal{O}_X(D),$$

where $0 < \varepsilon \ll 1$, and in general $\mathcal{J}(B)$ stands for the multiplier ideal of a \mathbb{Q} -divisor B (see [Laz04, Ch.9]). Indeed, [Sai09, Theorem 0.4] says that

$$F_0\mathcal{O}_X(*D) \simeq \tilde{V}^1\mathcal{O}_X \cdot \mathcal{O}_X(D),$$

while [BS05, Theorem 0.1] says that for any $\alpha \in \mathbb{Q}$ one has

$$\tilde{V}^\alpha\mathcal{O}_X \simeq \mathcal{J}((\alpha - \varepsilon)D).$$

Here the V -filtration notation means the following: assume that D is given locally by an equation f , and consider the graph embedding $i_f : X \rightarrow X \times \mathbb{C}$. One can consider the V -filtration on the left \mathcal{D} -module $i_{f*}\mathcal{O}_X = \mathcal{O}_X \otimes_{\mathbb{C}} \mathbb{C}[\partial_t]$ along $X_0 = X \times \{0\}$. The notation $\tilde{V}^\alpha\mathcal{O}_X$ stands for the filtration induced on $\mathcal{O}_X = \mathcal{O}_X \otimes 1$.

This allows us to deduce the Nadel vanishing theorem (see e.g. [Laz04, Theorem 9.4.8]), at least when D is a Cartier divisor.

Theorem 9.4. *With the notation above, if L is a line bundle on X such that $L - D$ is ample, then*

$$H^i(X, \omega_X \otimes L \otimes \mathcal{J}((1 - \varepsilon)D)) = 0 \quad \text{for all } i > 0.$$

Proof. From the discussion above it follows that for the left \mathcal{D} -module $\mathcal{O}_X(*D)$ the lowest graded piece for the filtration F is

$$\mathrm{Gr}_0^F \mathcal{O}_X(*D) = \mathcal{J}((1 - \varepsilon)D) \cdot \mathcal{O}_X(D),$$

so for the associated right \mathcal{D} -module we have

$$S(\mathcal{M}) = \omega_X \otimes \mathcal{O}_X(D) \otimes \mathcal{J}((1 - \varepsilon)D).$$

Corollary 9.1 implies that if A is ample, then

$$H^i(X, \omega_X \otimes A \otimes \mathcal{O}_X(D) \otimes \mathcal{J}((1 - \varepsilon)D)) = 0 \quad \text{for all } i > 0.$$

But by assumption we can write $L \simeq A \otimes \mathcal{O}_X(D)$ with A ample. \square

Remark 9.5 (Arbitrary \mathbb{Q} -divisors). The Nadel vanishing theorem for arbitrary \mathbb{Q} -divisors B is not in general a vanishing theorem for the lowest graded piece of the Hodge filtration corresponding to a mixed Hodge module; it is however a consequence of the same result. Roughly speaking one can reduce to the situation studied above after performing a Kawamata covering construction to arrive at a

Cartier divisor, using a bistrictness property of Hodge modules which allows us to deduce vanishing for the push-forward to the base, and finally passing to an eigensheaf of the push-forward. In other words multiplier ideals are naturally direct summands of Hodge theoretic objects, while Theorem 8.2 also applies to filtered direct summands of \mathcal{D} -modules underlying mixed Hodge modules, again since the functor $\mathrm{Gr}_k^F \circ \mathrm{DR}$ is exact. I thank N. Budur for this observation.

On the other hand, it is perhaps most natural to try and prove an analogue of the Kawamata-Viehweg vanishing theorem for \mathbb{Q} -divisors in the context of mixed Hodge modules. This will be done in Theorem 11.1 below. An analogous extension of Nadel vanishing is then an immediate consequence; see Corollary 11.3.

Abelian varieties. In the case of abelian varieties it turns out that Theorem 8.2 holds directly for the graded pieces of a filtered \mathcal{D} -module (\mathcal{M}, F) underlying a Hodge module itself, rather than those of its de Rham complex.

Proposition 9.6 ([PS13, Lemma 2.5]). *Let A be a complex abelian variety, (\mathcal{M}, F) the filtered \mathcal{D} -module underlying a mixed Hodge module on A , and let L be an ample line bundle. Then for each $k \in \mathbb{Z}$, we have*

$$H^i(A, \mathrm{Gr}_k^F \mathcal{M} \otimes L) = 0 \quad \text{for all } i > 0.$$

Proof. Denote $g = \dim A$. Consider for each $k \in \mathbb{Z}$ the complex of coherent sheaves

$$\mathrm{Gr}_k^F \mathrm{DR}_A(\mathcal{M}) = \left[\mathrm{Gr}_k^F \mathcal{M} \rightarrow \Omega_A^1 \otimes \mathrm{Gr}_{k+1}^F \mathcal{M} \rightarrow \cdots \rightarrow \Omega_A^g \otimes \mathrm{Gr}_{k+g}^F \mathcal{M} \right],$$

supported in degrees $-g, \dots, 0$. According to Theorem 8.2, this complex has the property that, for $i > 0$,

$$H^i(A, \mathrm{Gr}_k^F \mathrm{DR}_A(\mathcal{M}) \otimes L) = 0.$$

Using the fact that $\Omega_A^1 \simeq \mathcal{O}_A^{\oplus g}$, one can deduce the asserted vanishing theorem for the individual sheaves $\mathrm{Gr}_k^F \mathcal{M}$ by induction on k . Indeed, since $\mathrm{Gr}_k^F \mathcal{M} = 0$ for $k \ll 0$, inductively one has for each k a distinguished triangle

$$E_k \rightarrow \mathrm{Gr}_k^F \mathrm{DR}_A(\mathcal{M}) \rightarrow \mathrm{Gr}_{k+g}^F \mathcal{M} \rightarrow E_k[1],$$

with E_k an object satisfying $H^i(A, E_k \otimes L) = 0$. \square

This observation is one of the key points towards showing that, under the above assumptions, all graded pieces $\mathrm{Gr}_k^F \mathcal{M}$ satisfy the analogues of the generic vanishing theorems of [GL87], [GL91], [Hac04]. In view of the examples in Section 5, besides recovering these results the statement leads to new applications, for instance to Nakano-type generic vanishing (see [PS13, Theorem 1.2]).

Theorem 9.7 ([PS13], Theorem 1.1). *Under the assumptions of Proposition 9.6, each $\mathrm{Gr}_k^F \mathcal{M}$ is a GV-sheaf on A , i.e.*

$$\mathrm{codim}_{\mathrm{Pic}^0(A)} \{ \alpha \in \mathrm{Pic}^0(A) \mid H^i(A, \mathrm{Gr}_k^F \mathcal{M} \otimes \alpha) \neq 0 \} \geq i, \quad \text{for all } i \geq 0.$$

A stronger generic vanishing statement was proved in [PS13] for the total associated graded object

$$\mathrm{Gr}_\bullet^F \mathcal{M} = \bigoplus_k \mathrm{Gr}_k^F \mathcal{M},$$

seen as a coherent sheaf on $T^*A \simeq A \times H^0(A, \Omega_A^1)$; this was useful in proving that all holomorphic 1-forms on varieties of general type have zeros [PS14a].

10. Weak positivity. This section contains a proof of an extension of Viehweg’s weak positivity theorem for direct images of relative canonical sheaves, based on Theorem 8.2 and found jointly with C. Schnell; see also [Sch13]. The general strategy follows Kollár’s approach to semipositivity via vanishing theorems in [Kol86a, §3]. The shortness of the proof is due to the fact that one can apply the machinery of vanishing theorems to abstract Hodge modules.

Definition 10.1. A torsion-free coherent sheaf \mathcal{F} on a (quasi-)projective variety X is *weakly positive* on a non-empty open set $U \subseteq X$ if for every ample line bundle A on X and every $a \in \mathbb{N}$, the sheaf $\hat{S}^{ab} \mathcal{F} \otimes A^{\otimes b}$ is generated by global sections at each point of U for b sufficiently large. (Here $\hat{S}^p \mathcal{F}$ denotes the reflexive hull of the symmetric power $S^p \mathcal{F}$.)

Before proving the main result, let’s record a standard global generation consequence of Theorem 8.2.

Corollary 10.2. *Let X be a smooth projective complex variety of dimension n , and (\mathcal{M}, F) a filtered \mathcal{D} -module on X underlying a mixed Hodge module M . Then for any ample and globally generated line bundle L on X , the sheaf*

$$S(\mathcal{M}) \otimes L^{\otimes(n+1)}$$

is globally generated.

Proof. Denoting $\mathcal{F} = S(\mathcal{M}) \otimes L^{\otimes(n+1)}$, Corollary 9.1 implies that

$$H^i(X, \mathcal{F} \otimes L^{\otimes -i}) = 0 \text{ for all } i > 0.$$

The result is then an immediate consequence of the Castelnuovo-Mumford Lemma; see [Laz04, Theorem 1.8.5]. \square

We also need the following simplification of what is needed in order to check weak positivity under our hypotheses.

Lemma 10.3. *Let \mathcal{F} be a torsion-free sheaf on a smooth (quasi-)projective variety X , and L a line bundle on X . Then \mathcal{F} is weakly positive on an open set $U \subseteq X$ on which \mathcal{F} is locally free if $\mathcal{F}^{\otimes a} \otimes L$ is generated by global sections over U for all $a > 0$.*

Proof. This is well known, so I will only sketch the proof. First, it is standard that one can reduce to checking the definition for only one (not necessarily ample) line bundle L , and all $a > 0$; see [Vie83, Remark 1.3(ii)]. Now a torsion-free sheaf is locally free and therefore coincides with its reflexive hull outside of a closed set of codimension at least 2. On the other hand, its global sections inject into those of the reflexive hull. So it is enough to reduce the definition to the usual symmetric powers, which in turn are quotients of the tensor powers. \square

Viehweg's theorem in [Vie83] saying that $f_*\omega_{Z/X}$ is weakly positive for any surjective morphism $f : Z \rightarrow X$ of smooth projective varieties is a special case of the following result.⁸ The statement and proof are more conveniently phrased in terms of left \mathcal{D} -modules.

Theorem 10.4. *Let X be a smooth projective complex variety, and (\mathcal{N}, F) the filtered left \mathcal{D} -module on X underlying a mixed Hodge module M which is a variation of mixed Hodge structure on a non-empty open set $U \subseteq X$. Then $S(\mathcal{N})$ is weakly positive over U .*

Proof. Step 1. First, as M is a variation of mixed Hodge structure generically over X , it is well known that $S(\mathcal{N})$ is a torsion-free sheaf on X . Fix now a positive integer m , and consider the diagonal embedding

$$i : X \hookrightarrow X \times \cdots \times X,$$

where the product is taken m times. On this product, consider the box product mixed Hodge module

$$M^{\boxtimes m} := M \boxtimes \cdots \boxtimes M.$$

As the filtration on $M^{\boxtimes m}$ is the convolution of the filtrations on the individual factors, it is not hard to see that $p(\mathcal{N}^{\boxtimes m}) = m \cdot p(\mathcal{N})$ and moreover

$$i^* S(\mathcal{N}^{\boxtimes m}) = S(\mathcal{N})^{\otimes m}.$$

Denoting by $r = (m-1)n$ the codimension of X via the diagonal embedding, in the derived category of coherent sheaves on X we have a natural morphism

$$(10.5) \quad F_{m \cdot p(\mathcal{N}) - r} i^! (\mathcal{N}^{\boxtimes m}, F) \longrightarrow \mathbf{L}i^* S(\mathcal{N}^{\boxtimes m})[-r],$$

which is an isomorphism over the open set U where M is a variation of mixed Hodge structure. This follows for instance from [Sch12, Lemma 2.17] (see also [Sch13, Lemma 3.2]).

Step 2. We can specialize formula (10.5) by passing to the cohomology sheaves in degree r , in order to obtain a natural sheaf homomorphism

$$(10.6) \quad S(\mathcal{Q}) = F_{m \cdot p(\mathcal{N}) - r} \mathcal{Q} \longrightarrow S(\mathcal{N})^{\otimes m}$$

which is an isomorphism on U ; here (\mathcal{Q}, F) is another filtered left \mathcal{D} -module on X , underlying the object $i^* M^{\boxtimes m}$ in $\text{MHM}(X)$.

Fix now a very ample line bundle L on X . In order to deduce that $S(\mathcal{N})$ is weakly positive over U , using Lemma 10.3 it suffices then to show that $S(\mathcal{Q}) \otimes \omega_X \otimes L^{\otimes(n+1)}$ is globally generated, where $n = \dim X$. But this is a consequence of Corollary 10.2, recalling that $S(\mathcal{Q}) \otimes \omega_X$ is the lowest non-zero graded piece of the right \mathcal{D} -module associated to \mathcal{Q} . \square

In [Vie83], Viehweg proved that if $f : Z \rightarrow X$ is a surjective morphism of smooth projective varieties, then $f_*\omega_{Z/X}^{\otimes m}$ is weakly positive for $m \geq 2$ as well. A natural question to ask in this direction is the following:

⁸We apply it to the left \mathcal{D} -modules \mathcal{N}_i corresponding to \mathcal{M}_i in the decomposition $f_*(\omega_Z, F) \simeq \bigoplus_i (\mathcal{M}_i, F)[-i]$; see Example 5.2.

Question 10.7. Let $f : Z \rightarrow X$ be a surjective morphism of smooth projective varieties, and (\mathcal{M}, F) the filtered left \mathcal{D} -module underlying a mixed Hodge module M which is a variation of mixed Hodge structure on a non-empty open set in Z . Is

$$f_* \left(S(\mathcal{N}) \otimes \omega_{Z/X}^{\otimes m} \right)$$

weakly positive for all $m \geq 1$?

Assuming a positive answer to this question, the exact same method of proof as in Theorem 10.4 would imply for all $m \geq 2$ the weak positivity of

$$f_* \left(S(\mathcal{N})^{\otimes m} \otimes \omega_{Z/X}^{\otimes m} \right).$$

It is worth noting that it is indeed now possible to give a proof of Viehweg's statement on $f_* \omega_{Z/X}^{\otimes m}$ using cohomological methods à la Kollár; see [PS14b].

11. Kawamata-Viehweg-type vanishing. In this section I will show that the Kawamata-Viehweg vanishing theorem for \mathbb{Q} -divisors continues to hold for the lowest graded piece of a mixed Hodge module as long as its singular locus does not intersect the augmented base locus $\mathbf{B}_+(L)$ of a big and nef line bundle (in particular always for variations of mixed Hodge structure). The proof follows quite closely the original one, with modifications permitted by Saito's study of non-characteristic pullbacks. I expect a stronger version to hold, at least under certain non-characteristicity hypotheses with respect to $\mathbf{B}_+(L)$.⁹

Theorem 11.1. *Let (\mathcal{M}, F) be the filtered right \mathcal{D} -module underlying a mixed Hodge module M on a smooth projective variety X , and let L be a line bundle on X with $L \sim_{\mathbb{Q}} A + \Delta$, where A is a big and nef \mathbb{Q} -divisor on X and (X, Δ) is a klt pair. Assume that $\mathbf{B}_+(A) \cup \text{Supp } \Delta$ is contained in the smooth locus of M . Then*

$$H^i(X, S(\mathcal{M}) \otimes L) = 0 \quad \text{for all } i > 0.$$

Remark 11.2. In particular we have the vanishing above if L is a big and nef line bundle such that $\mathbf{B}_+(L)$ is contained in the smooth locus of M . Note that one does not have a similar statement for other associated graded pieces $\text{Gr}_k^F \text{DR}(\mathcal{M})$ of the filtered de Rham complex, as in the case of Kodaira-Saito vanishing. This is already well known for the trivial Hodge module $M = \mathbb{Q}_X^H[n]$. In this case, by Example 5.1 the graded pieces are $\Omega_X^k[n-k]$ with $n = \dim X$. Simple examples show however that for $k < n$ the Nakano extension of Kodaira vanishing does not usually hold for twists by big and nef line bundles; see [Laz04, Example 4.3.4].

In order to understand the statement and proof, we need to review a few more definitions and results. Before doing this, let's note that an immediate consequence of the theorem above is the following generalization of the Nadel vanishing theorem; see also Section 9.

Corollary 11.3. *Let X be a smooth projective variety, and D an effective \mathbb{Q} -divisor on X with associated multiplier ideal $\mathcal{I}(D)$. Let L be a line bundle in X such that $L - D$ is big and nef, and assume that $\mathbf{B}_+(L - D) \cup \text{Supp } D$ is contained*

⁹This is pursued by one of my Ph.D. students as part of his thesis work.

in the smooth locus of a mixed Hodge module M with underlying filtered \mathcal{D} -module (\mathcal{M}, F) . Then

$$H^i(X, S(\mathcal{M}) \otimes L \otimes \mathcal{I}(D)) = 0 \text{ for all } i > 0.$$

Higher direct images of the lowest Hodge piece. Let X be a smooth variety. Recall that according to M. Saito's theory [Sai90], for a mixed Hodge module M with strict support equal to X , there exists a maximal non-empty open set $U \subseteq X$ on which M is variation of mixed Hodge structure, denoted say by \mathbf{V} ; we call this the *smooth locus* of M . Note that the lowest Hodge piece $S(\mathcal{M})$ is a locally free sheaf on U .

As the functor $S(\cdot)$ is exact, we can often restrict our study to the case when M is a pure Hodge module which is a polarized variation of Hodge structure on U . In this case, in response to a conjecture of Kollár, Saito proved (among other things) the following, the second part of which can be seen as a generalization of the Grauert-Riemenschneider vanishing theorem.

Theorem 11.4 (Saito, [Sai91]). *Let $f : X \rightarrow Y$ be a surjective projective morphism (with Y possibly singular), and let (\mathcal{M}, F) be the filtered \mathcal{D} -module underlying a pure Hodge module with strict support X that is generically a polarized variation of Hodge structure \mathbf{V} . For each $i \geq 0$, one has*

$$R^i f_* S(\mathcal{M}) = S(Y, \mathbf{V}^i),$$

the lowest Hodge piece of the variation of Hodge structure \mathbf{V}^i on the intersection cohomology of \mathbf{V} along the fibers of f . Consequently, $R^i f_ S(\mathcal{M})$ are torsion-free, and in particular*

$$R^i f_* S(\mathcal{M}) = 0 \text{ for } i > \dim X - \dim Y.$$

Augmented base loci. We start by recalling the definition and some basic results on augmented base loci of divisors.

Definition 11.5 ([ELM⁺06, §1]). Let D be a \mathbb{Q} -divisor on a normal complex projective variety X . The *augmented base locus* of L is

$$\mathbf{B}_+(D) := \mathbf{B}(D - \epsilon H),$$

where H is any ample divisor on X , $0 < \epsilon \ll 1$ is rational, and $\mathbf{B}(D - \epsilon H)$ denotes the stable base locus of the \mathbb{Q} -divisor $D - \epsilon H$, i.e. the base locus of $|m(D - \epsilon H)|$ for $m \gg 0$. If L is a line bundle, we define $\mathbf{B}_+(L)$ similarly. It is not hard to check (see [ELM⁺06, Proposition 1.5]) that equivalently one has

$$(11.6) \quad \mathbf{B}_+(D) = \bigcap_{D=A+E} \text{Supp } E,$$

where the intersection is taken over all \mathbb{Q} -divisor decompositions of D such that A is ample and E is effective.

We have that $\mathbf{B}_+(L) \neq X$ if and only if L is big. When L is a big and nef, according to Nakamaye's theorem [Nak00], one has the following description

$$\mathbf{B}_+(L) = \text{Null}(V),$$

where $\text{Null}(V)$ is the union of all subvarieties $V \subset X$ such that $L^{\dim V} \cdot V = 0$, or equivalently $L|_V$ is not big.

We will need birational interpretations of the augmented base locus. The following results are standard, but seem hard to locate in the literature.

Lemma 11.7. *If D is a \mathbb{Q} -divisor on X , then*

$$\mathbf{B}_+(D) = \bigcap_{f,A,E} f(\text{Supp } E),$$

where the intersection is taken over all projective birational morphisms $f : Y \rightarrow X$ with Y normal, and all decompositions $f^*D = A + E$, with A ample and E effective.

Proof. We need to show that for any decomposition $f^*D = A + E$ as in the statement we have $\mathbf{B}_+(D) \subseteq f(\text{Supp } E)$. Let H be an ample Cartier divisor on X . Consider $0 < \epsilon \ll 1$ rational such that $\mathbf{B}_+(D) = \mathbf{B}(D - \epsilon H)$ and $A - \epsilon f^*H$ is ample. Writing

$$f^*(D - \epsilon H) = (A - \epsilon f^*H) + E,$$

we deduce that $f^{-1}(\mathbf{B}(D - \epsilon H)) = \mathbf{B}(f^*(D - \epsilon H)) \subseteq \text{Supp } E$, which implies what we want. \square

Lemma 11.8. *Let $f : Y \rightarrow X$ be a birational morphism of smooth projective varieties, and $\text{Exc}(f) \subseteq Y$ its exceptional locus. If D is an \mathbb{Q} -divisor on X , then*

$$\mathbf{B}_+(f^*(D)) = f^{-1}(\mathbf{B}_+(D)) \cup \text{Exc}(f).$$

Proof. Assume first that V is an irreducible component of $\text{Exc}(f)$. If $W = f(V)$, then $\dim W < \dim V$, and so $f^*D|_V$ is not big. This implies $V \subseteq \mathbf{B}_+(f^*(D))$, essentially by definition. Assume now that $x \in f^{-1}(\mathbf{B}_+(D)) \setminus \text{Exc}(f)$. If $x \notin \mathbf{B}_+(f^*(D))$, then by (11.6) we can write $f^*D = A + E$ with A ample, E effective, and $x \notin \text{Supp } E$. But by Lemma 11.7, this gives contradicts the fact that $f(x) \in \mathbf{B}_+(D)$.

For the other inclusion, we show that if $x \in \mathbf{B}_+(f^*(D)) \setminus \text{Exc}(f)$, then $f(x) \in \mathbf{B}_+(D)$. If this were not the case, then we would find A ample and E effective on X such that $D = A + E$ and $f(x) \notin \text{Supp } E$. Now by the Negativity Lemma (see e.g. [Laz04, Corollary 4.1.4]), there exists an effective \mathbb{Q} divisor F supported on $\text{Exc}(f)$ such that $f^*A - F$ is ample. Using the decomposition

$$f^*D = (f^*A - F) + (f^*E + F),$$

given that $x \notin \text{Supp}(f^*E + F)$ we deduce that $x \notin \mathbf{B}_+(f^*(D))$, a contradiction. \square

Proof of Theorem 11.1. First, just as in the proof of Saito's vanishing theorem, due to the exactness of the functor $S(\cdot)$ we can reduce to assuming that M is a pure Hodge module. I will divide the proof into a few steps which loosely follow the standard steps in the proof of the Kawamata-Viehweg theorem. In the first three steps we will assume that L is a big and nef line bundle, and $\Delta = 0$. The last two will deal with the general case.

The line bundle case. Note to begin with that since L is big, in general there exist an $m > 0$, an ample line bundle A , and an effective divisor E , such that

$$(11.9) \quad L^{\otimes m} \simeq A \otimes \mathcal{O}_X(E).$$

Step 1. This is a Norimatsu-type statement (see [Laz04, Lemma 4.3.5]): we show that if A is an ample line bundle, and $E \subset X$ is a reduced simple normal crossings divisor on X contained in the smooth locus of M , then

$$H^i(X, S(\mathcal{M}) \otimes A \otimes \mathcal{O}_X(E)) = 0 \text{ for all } i > 0.$$

Let's assume first that E is a smooth divisor. As $S(\mathcal{M})$ is locally free in a neighborhood of E , we have a short exact sequence

$$0 \longrightarrow S(\mathcal{M}) \otimes A \longrightarrow S(\mathcal{M}) \otimes A \otimes \mathcal{O}_X(E) \longrightarrow S(\mathcal{M})|_E \otimes A|_E \otimes \mathcal{O}_E(E) \longrightarrow 0$$

Passing to cohomology and applying Corollary 9.1, we see that it is enough to show that

$$H^i(E, S(\mathcal{M})|_E \otimes \mathcal{O}_E(E) \otimes A|_E) = 0 \text{ for all } i > 0.$$

Again by Corollary 9.1, it suffices then to note that $S(\mathcal{M})|_E \otimes \mathcal{O}_E(E) \simeq S(\mathcal{M}')$, for some filtered \mathcal{D} -module underlying a mixed Hodge module M' on E . We can in fact take

$$(\mathcal{M}', F) := (\mathcal{H}^1 i^! \mathcal{M}, F).$$

On one hand, this filtered \mathcal{D} -module underlies a Hodge module, as

$$\mathcal{H}^1 i^! \mathcal{M}, F) \simeq i^!(\mathcal{M}, F)[1]$$

by [Sai88, Lemme 3.5.6]. On the other hand, since E is contained in the smooth locus of M , using [Sch12, Lemma 2.17] (as in the proof of Theorem 10.4) we see that there is an isomorphism $S(\mathcal{N}') \simeq S(\mathcal{N})|_E$, where \mathcal{N} is again notation for the associated left \mathcal{D} -modules. This is equivalent to what we want by adjunction.

In general we have $E = E_1 + \dots + E_k$, where E_j are smooth divisors with transverse intersections. The statement can be easily proved by induction on k , using exact sequences similar to the one above, and the fact that M continues to be a variation of mixed Hodge structure when restricted to the log-canonical centers of E .

Step 2. In this step we show that we can reduce the general statement to the case where in (11.9) we have that E has simple normal crossings support, and this support is contained in the smooth locus of M . Consider the notation of Definition 11.5, so that

$$\mathbf{B}_+(L) = \mathbf{B}(L - \epsilon H) = \text{Bs}(L^{\otimes k} \otimes \mathcal{O}_X(-k\epsilon H)),$$

for k sufficiently large and divisible, and $\text{Bs}(\cdot)$ stands for the usual base locus. We consider $\mu : Y \rightarrow X$ a log-resolution of the linear series $|L^{\otimes k} \otimes \mathcal{O}_X(-k\epsilon H)|$, so that

$$\mu^*(L^{\otimes k} \otimes \mathcal{O}_X(-k\epsilon H)) \simeq M_k \otimes \mathcal{O}_Y(F_k),$$

where M_k is the moving part of the pullback, a big and basepoint-free line bundle, and F_k is its fixed divisor. From Lemma 11.8 we have that

$$\mathbf{B}_+(\mu^*L) = \mu^{-1}(\mathbf{B}_+(L)) \cup \text{Exc}(\mu) = \text{Supp}(F_k) \cup \text{Exc}(\mu),$$

which is a divisor with simple normal crossings support on Y .

By assumption $\mathbf{B}_+(L)$ is contained in the smooth locus of M . Choosing the log-resolution to be an isomorphism outside of $\mathbf{B}_+(L)$, by Example 7.5 we have

that μ is non-characteristic for (\mathcal{M}, F) . Recall that this implies that the filtered inverse image $\mu^*(\mathcal{M}, F) = (\tilde{\mathcal{M}}, F)$ is given by the formula

$$\tilde{\mathcal{M}} = \mu^{-1}\mathcal{M} \otimes_{\mu^{-1}\mathcal{O}_X} \omega_{Y/X} \quad \text{and} \quad F_p \tilde{\mathcal{M}} = \mu^* F_p \mathcal{M} \otimes \omega_{Y/X},$$

and this underlies the Hodge module $\mu^* M$. We see then that $S(\mu^* \mathcal{M}) \simeq \mu^* S(\mathcal{M}) \otimes \omega_{Y/X}$, and so

$$\mu_* S(\mu^* \mathcal{M}) \simeq S(\mathcal{M}),$$

as $\mu_* \omega_{Y/X} \simeq \mathcal{O}_X$. Assuming that we proved that

$$(11.10) \quad H^i(Y, S(\mu^* \mathcal{M}) \otimes \mu^* L) = 0 \quad \text{for all } i > 0,$$

this implies the vanishing we want on X as $R^i \mu_* S(\mu^* M) = 0$, which is a consequence of Theorem 11.4.

Let's now write

$$F_k = \sum_j a_j E_j,$$

with the convention that $a_j \geq 0$, so that we may assume that the sum contains all the exceptional divisors of μ among the E_j . By construction we have that $\mathbf{B}_+(\mu^* L)$ is contained in the smooth locus of $\mu^* M$; equivalently, this statement holds for all E_j in the sum above.

Finally, note that by construction we have

$$\mu^* L^{\otimes k} \simeq \mu^* \mathcal{O}_X(k\epsilon H) \otimes M_k \otimes \mathcal{O}_Y(F_k),$$

and the line bundle $\mu^* \mathcal{O}_X(k\epsilon H) \otimes M_k$ is still big and nef. To conclude, one appeals to a version of the Negativity Lemma, stating that for such a $k \gg 0$, there exist $b_j \geq 0$ such that

$$\mu^* \mathcal{O}_X(k\epsilon H) \otimes \mathcal{O}_Y(-\sum_j b_j F_j)$$

is ample, where the sum runs over the exceptional divisors of μ (and so with the same convention as above we can assume that it runs over all E_j); see e.g. [Laz04, Corollary 4.1.4]. But now we can write

$$\mu^* L^{\otimes k} \simeq (\mu^* \mathcal{O}_X(k\epsilon H) \otimes \mathcal{O}_Y(-\sum_j b_j F_j)) \otimes \mathcal{O}_Y(\sum_j (a_j + b_j) F_j),$$

which is of the form required at the beginning of this reduction step.

Step 3. In this last step we conclude the proof assuming that E in (11.9) has simple normal crossings support contained in the smooth locus of M , which is the outcome of Step 2. By standard arguments using Kawamata covers, it is known that there exists a finite cover $f : Y \rightarrow X$ with Y smooth projective, such that

$$f^* L \simeq A' \otimes \mathcal{O}_Y(E'),$$

with A' ample and E' a reduced simple normal crossings divisor; see e.g. [Laz04, p.255]. Moreover, f can be chosen to be non-characteristic with respect to (\mathcal{M}, F) .

This last statement requires some discussion; recall that Kawamata covers can be constructed in two steps (see [Laz04, Proposition 4.1.12]). The first is a Bloch-Gieseker type cover $g : Z \rightarrow X$, where for some component E_1 of E one can write $g^* E = kE_1$, for a given k and some E_1 not necessarily effective. In this step one can assume that E is very ample by writing it as the difference of two very ample

line bundles, and then g can be constructed so as to be ramified along a generic union of hyperplane sections of X in the embedding given by E ; see the proof of [Laz04, Theorem 4.1.10]. From this genericity it follows that g is non-characteristic with respect to (\mathcal{M}, F) . On the other hand, the second step is to consider a cyclic cover $h : Y \rightarrow Z$, which is ramified along f^*E_1 ; since this is contained in the smooth locus of f^*M , this cover is also non-characteristic. One then applies this procedure inductively for all components of E .

Going back to the proof, we can now consider the filtered inverse image $f^*(\mathcal{M}, F)$ underlying the pullback Hodge module just as in Step 2. Note that we have $E' = f^{-1}(\text{Supp } E)$, and so E' is contained in the smooth locus of f^*M . By Step 1, we then have

$$H^i(X, S(f^*\mathcal{M}) \otimes f^*L) = 0 \quad \text{for all } i > 0.$$

But precisely as in Step 2 we have that

$$f_*S(f^*\mathcal{M}) \simeq S(\mathcal{M}) \otimes f_*\omega_{Y/X}.$$

As \mathcal{O}_X is a direct summand of $\omega_{Y/X}$ via the trace map, we obtained the desired vanishing using the projection formula.

The \mathbb{Q} -divisor case. We do this in two further steps which reduce us to the line bundle case discussed above. We first reduce to the case when $\text{Supp } \Delta$ is a simple normal crossings divisor.

Step 4. Let $\mu : Y \rightarrow X$ be a log-resolution of (X, Δ) , and write

$$K_Y - \mu^*(K_X + \Delta) = P - N,$$

where P and N are effective \mathbb{Q} -divisors with simple normal crossings support, without common components, and such that P is exceptional and all the coefficients in N are strictly less than 1. We then have

$$K_Y + N + [P] - P + \mu^*A = \mu^*(K_X + \Delta + A) + [P],$$

and so there exists a line bundle L' on Y such that $L' \sim_{\mathbb{Q}} \mu^*A + \Delta'$, where $\Delta' = N + [P] - P$, a strictly boundary divisor with normal crossings support. Note that μ^*A is still big and nef, and in fact by Lemma 11.8 we have

$$\mathbf{B}_+(\mu^*A) = \mu^{-1}(\mathbf{B}_+(A)) \cup \text{Exc}(\mu).$$

We can choose μ such that it is an isomorphism outside the support of Δ . It follows that both $\mathbf{B}_+(\mu^*A)$ and $\text{Supp } \Delta'$ are contained in the smooth locus of μ^*M . Note finally that it is enough to show that

$$H^i(Y, S(\mu^*\mathcal{M}) \otimes L') = 0 \quad \text{for all } i > 0.$$

Indeed, we have observed before that

$$\mu_*S(\mu^*\mathcal{M}) \simeq S(\mathcal{M}) \quad \text{and} \quad R^i\mu_*S(\mu^*\mathcal{M}) = 0 \quad \text{for } i > 0.$$

Step 5. It is enough to assume then that Δ is a divisor with simple normal crossings support, say $\Delta = \sum_{i=1}^k a_i D_i$, with $0 < a_i < 1$ and D_i smooth.

The strategy is to prove the statement by induction on k . The case $k = 0$ is the line bundle case proved above. Assume now that $k > 0$, and let's write $a_1 = \frac{p}{q}$. Note that $0 < p \leq q - 1$. Just as in Step 3, one considers a Kawamata cover associated to the divisor D_1 ; concretely, there exists a finite morphism $f : Y \rightarrow X$,

with Y smooth projective, such that on Y the divisor D_1 becomes divisible by d . In other words, we have

$$L' := f^*L \sim_{\mathbb{Q}} A' + cD'_1 + \sum_{i=2}^k a_i D'_i,$$

where $A' = f^*A$ and $D'_i = f^*D_i$, still satisfying the fact that $\sum D'_i$ has simple normal crossings. Moreover, this morphism can be chosen to be non-characteristic for (\mathcal{M}, F) , so we can deal with f^*M as in the previous proof.

By induction we can now assume that the line bundle $L' \otimes \mathcal{O}_Y(-cD'_1)$ satisfies

$$H^i(X', S(f^*\mathcal{M}) \otimes L' \otimes \mathcal{O}_Y(-cD'_1)) = 0 \text{ for all } i > 0.$$

Recall that due to the definition of the filtration under non-characteristic inverse image we have $S(f^*\mathcal{M}) \simeq f^*S(\mathcal{M}) \otimes \omega_{Y/X}$. On the other hand, it is standard that in the covering construction above we have that $f_*(L' \otimes \mathcal{O}_Y(-cD'_1) \otimes \omega_{Y/X})$ contains L as a direct summand. The desired vanishing follows from the projection formula.

REFERENCES

- [BS05] N. Budur and M. Saito, *Multiplier ideals, V-filtration, and spectrum*, J. Algebraic Geom. **14** (2005), no. 2, 269–282.
- [ELM⁺06] L. Ein, R. Lazarsfeld, M. Mustață, M. Nakamaye, and M. Popa, *Asymptotic invariants of base loci*, Ann. Inst. Fourier **56** (2006), 1701–1734.
- [EV92] H. Esnault and E. Viehweg, *Lectures on vanishing theorems*, DMV Seminar, vol. 20, Birkhäuser Verlag, Basel, 1992.
- [GL87] M. Green and R. Lazarsfeld, *Deformation theory, generic vanishing theorems, and some conjectures of Enriques, Catanese and Beauville*, Invent. Math. **90** (1987), no. 2, 389–407.
- [GL91] ———, *Higher obstructions to deforming cohomology groups of line bundles*, J. Amer. Math. Soc. **1** (1991), no. 4, 87–103.
- [Hac04] C. Hacon, *A derived category approach to generic vanishing*, J. Reine Angew. Math. **575** (2004), 173–187.
- [Kol86a] J. Kollár, *Higher direct images of dualizing sheaves I*, Ann. of Math. **123** (1986), no. 1, 11–42.
- [Kol86b] ———, *Higher direct images of dualizing sheaves II*, Ann. of Math. **124** (1986), 171–202.
- [Laz04] R. Lazarsfeld, *Positivity in algebraic geometry I & II*, Ergebnisse der Mathematik und ihrer Grenzgebiete, Springer-Verlag, Berlin, 2004.
- [MS] P. Maisonobe and C. Sabbah, *Aspects of the theory of \mathcal{D} -modules (Kaiserslautern 2002)*, at <http://www.math.polytechnique.fr/cmat/sabbah/livres.html>.
- [Nak00] M. Nakamaye, *Stable base loci of linear series*, Math. Ann. **318** (2000), 837–847.
- [PS13] M. Popa and C. Schnell, *Generic vanishing theory via mixed Hodge modules*, Forum of Math., Sigma **1** (2013), 1–60.
- [PS14a] ———, *Kodaira dimension and zeros of holomorphic one-forms*, Ann. of Math. **179** (2014), 1–12.
- [PS14b] ———, *On direct images of pluricanonical bundles*, preprint arXiv:1405.6125 (2014).
- [Sai88] M. Saito, *Modules de Hodge polarisables*, Publ. RIMS **24** (1988), 849–995.
- [Sai90] ———, *Mixed Hodge modules*, Publ. Res. Inst. Math. Sci. **26** (1990), no. 2, 221–333.
- [Sai91] ———, *On Kollár’s conjecture*, Several complex variables and complex geometry, Part 2 (Santa Cruz, CA, 1989), Proc. Sympos. Pure Math. **52** (1991), 509–517.
- [Sai93] ———, *On b-function, spectrum and rational singularity*, Math. Ann. **295** (1993), no. 1, 51–74.
- [Sai09] ———, *On the Hodge filtration of Hodge modules*, Mosc. Math. J. **9** (2009), no. 1, 161–191, back matter (English, with English and Russian summaries).

- [Sch12] C. Schnell, *Complex analytic Néron models for arbitrary families of intermediate Jacobians*, *Invent. Math.* **188** (2012), no. 1, 1–81.
- [Sch13] ———, *Weak positivity via mixed Hodge modules*, dedicated to Herb Clemens, to appear (2013).
- [Sch14a] ———, *On Saito's vanishing theorem*, preprint (2014).
- [Sch14b] ———, *An overview of Morihiko Saito's theory of mixed Hodge modules*, preprint arXiv:1405.3096 (2014).
- [Vie83] E. Viehweg, *Weak positivity and the additivity of the Kodaira dimension of certain fiber spaces*, *Adv. Studies PureMath.* **1** (1983), 329–353.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS AT CHICAGO, 851 S. MORGAN STREET,
CHICAGO, IL 60607, USA

E-mail address: `mpopa@uic.edu`